

## Original Research Article

### **Bacterial vaginosis: Prevalence and risk factors among women in Dakar, Senegal**

#### **Abstract:**

**Introduction:** Bacterial vaginosis (BV) is an imbalance of the vaginal flora that most commonly affects women of childbearing age. It is characterised by discomfort due to vaginal discharge and is associated with serious complications in women. We aimed to determine the prevalence of BV and its risk factors.

**Methodology:** We conducted a prospective cross-sectional descriptive study in November 2020 at the Pasteur Institute of Dakar among women who were seen for vaginal sampling. We collected the following parameters: age, parity, history of infection, pregnancy and contraception. VB was diagnosed using the Amsel criteria and the Nugent score. Scoring is based on the presence of different bacterial morphotypes where a score  $\geq 7$  indicates the presence of BV, 4-6 indicates intermediate flora and 0- 3 normal flora. Analysis was performed using SPSS version 20.0. A statistically significant difference between the occurrence of bacterial vaginosis and a risk factor was retained for a p value  $< 0.05$ .

**Results:** A total of 236 patients were included with a median age of 38 years. The prevalence of BV was 20.3%. It was more prevalent in women aged between [30-40 years] (28/101, 27.7%) followed by the age group [20-30 years] (13/51, 25.5%) with a statistically significant difference ( $p=0.0001$ ). VB was more prevalent in nulliparous or primiparous women (28/118, 23.7%) and in those with no history of vaginal infections (20/83, 24.1%). Pregnant women and women without contraception were more affected with respectively (12/42, 28.6%) and (43/199, 21.6%). No relationship was found between BV and these previous potential risk factors.

**Conclusion:** BV is a very common condition in women of childbearing age. Its complexity motivates numerous researches in order to elucidate its real implication in the occurrence of adverse gynaecological conditions.

**Keywords:** bacterial vaginosis, prevalence, risk factors.

#### **I. Introduction**

Bacterial vaginosis (BV) is defined as a vaginal dysbiosis resulting in an imbalance of the vaginal flora due to an increase in anaerobic bacteria and the disappearance of lactobacilli, which are considered to be protective, and are replaced by commensal organisms: *Mobiluncus*

*spp.*, *Gardnerella vaginalis*, *Mycoplasma hominis*, etc., which are responsible for a number of symptoms (1). It is manifested by a grey, homogeneous, abundant, adherent and foul-smelling vaginal discharge, an increase in vaginal pH and the presence of bacteria adhering to the walls of the vaginal cells (clue-cells), which can be detected by microscopic examination of a slide of vaginal secretions (2). It is a common condition and is thought to be the leading cause of vaginitis, ahead of vaginal *Candida* spp infections, affecting mainly young sexually active women, but can occur in the absence of sexual intercourse (3,4).

Thus, the prevalence of BV is generally estimated to be between 15 and 30%, but some studies show higher prevalences (61% in an STI clinic) or sometimes much lower (4.9% to 20% in pregnant women) (5). However, its prevalence varies considerably according to ethnic and/or geographical origin (4-58%), with women of African origin being associated with the highest prevalence (6). Moreover, 20-50% of women in sub-Saharan Africa suffer from bacterial vaginosis, with a predominance of sexually active women, in whom almost one-third are affected (7).

Serious consequences may be associated with an increased risk of sexually transmitted infections, including human immunodeficiency virus, and complications of the upper genital tract especially in pregnancy with intra-amniotic infections, infections of the uterus after delivery or after abortion (8,9). Indeed, numerous retrospective studies, mainly from the United States and Scandinavian countries, suggest an association between VB and an increased risk of preterm delivery, chorioamnionitis, premature rupture of membranes (PMR) and even first trimester abortions (10).

After several decades of research on VB, the pathophysiology and sequelae associated with VB remain poorly understood, thus accentuating the enigma that this condition represents (1,11). It is in this context that we conducted this study to determine the prevalence of BV and to investigate potential risk factors.

## **II. Material and method:**

### **1. Type, setting and period of study**

This was a prospective cross-sectional study with a descriptive aim conducted in November 2020 at the Pasteur Institute in Dakar.

### **2. Study population**

- **Inclusion criteria:** the patients included were those received at the laboratory during the study period on whom a vaginal swab was taken after consent was obtained.

- **Non-inclusion criteria:** patients who did not give consent were not included.

### 3. Methods

- **Parameters studied:** we collected the following parameters: age, parity, history of vaginal infections, pregnancy and contraception.


- **Diagnosis of BV:** BV is diagnosed using the Amsel criteria and the Nugent score (12-16). In the laboratory, a vaginal swab was taken from the ectocervix (posterior cul de sac). A Gram stain was performed on this specimen to determine the Nugent score. The scoring is based on the presence of different bacterial morphotypes where a score  $\geq 7$  indicates the presence of BV, 4-6 indicates intermediate flora and 0-3 indicates normal flora (12,15,17).

### 4. Statistical analysis

Data entry was performed with Excel version 2010 and analysis with SPSS version 20.0. The Chi-square test was performed to see if there was a statistically significant difference between the occurrence of bacterial vaginosis and other risk factors. The level of significance was set at a value of  $p < 0.05$ .

### III. Results

A total of 236 patients were included with a median age of 38 years with a predominance of the age group [30-40 years] with 42.80% followed by the age groups [20-30 years], [40-50 years] with 21.60% and 6.80% respectively. The number of nulliparous or primiparous women was equal to multiparous women. The notion of using contraception and a history of infections were reported in 15.7% and 35% respectively. We found a prevalence of BV of 20.3% (Figure 1).



**Fig 1 : Distribution of patients according to vaginal flora**

The vaginal flora of women with BV was composed of *Gardenerella vaginalis* (GV) in all cases of BV, which may be associated with *Corynebacteria* (Coryne) (29%) and/or *Mobiluncus* (Mob) (44%). We investigated potential factors that may influence the occurrence of bacterial vaginosis (Table 1). With regard to age group, BV was more common

in women between the ages of [30-40 years] (28/101, 27.7%) followed by the age group [20-30 years] (13/51, 25.5%). We found a statistically significant difference ( $p=0.0001$ ) between VB and age group. For parity, VB was more noted in nulliparous or primiparous women (28/118, 23.7%). No significant difference was observed between VB and parity ( $p=0.131$ ). Regarding history of infection, BV occurred in women with no history (20/83, 24.1%) with no statically significant difference ( $p=0.318$ ).

VB was more frequent in pregnant women with only (12/42, 28.6%) compared to others without any significant difference ( $p=0.315$ ). BV was more common in women without contraception (43/199, 21.6%) compared to women on contraception (5/37, 13.5%) and no relationship was found between BV and contraception ( $p=0.532$ ).

UNDER PEER REVIEW

**Table 1: Relationship between flora type and potential risk factors**

Caractéristiques	Type of flora			Total N	p-value
	Normal n (%)	Intermediate n (%)	Vaginosis n (%)		
<b>Age groups (year)</b>					
[20-30[	23 (45,0)	15 (29,4)	13 (25,5)	51	0,0001
[30-40[	52 (51,5)	21 (20,8)	28 (27,7)	101	
[40-50[	26 (51,0)	24 (47,0)	1 (2,0)	51	
[50-60[	2 (13,3)	10 (66,6)	3 (20,0)	15	
[60-70[	3 (18,7)	10 (62,5)	3 (18,7)	16	
[70 - 80[	0 (0,0)	2 (100,0)	0 (0,0)	2	
<b>Parity</b>					
≤ 1	56 (47,4)	34 (28,8)	28 (23,7)	118	0,131
≥ 2	50 (42,4)	48 (40,6)	20 (17,0)	118	
<b>History of infections</b>					
Yes	74 (48,3)	51 (33,3)	28 (18,3)	153	0,318
No	32 (38,5)	31 (37,3)	20 (24,1)	83	
<b>Pregnancy</b>					
Yes	18 (42,8)	12 (28,6)	12 (28,6)	42	0,315
No	88 (45,3)	70 (36,0)	36 (18,5)	194	
<b>Contraception</b>					
Yes	18 (48,6)	14 (37,8)	5 (13,5)	37	0,532
No	88 (44,2)	68 (34,1)	43 (21,6)	199	
<b>Total</b>	<b>106 (45,0)</b>	<b>82 (34,7)</b>	<b>48 (20,3)</b>	<b>236</b>	

#### **IV. Discussion**

BV is a very common vaginal dysbiosis in women of childbearing age (18,19). Then, we found a prevalence of BV of 20.3%. The prevalence observed in our study is within the prevalence range (6%-32%) documented by most studies in Africa (1,20). Indeed, prevalences of 14.2% and 30.3% have been reported in Nigeria and Zimbabwe respectively (11). The same is true in other geographical areas, notably Bangladesh and the United States, where prevalence was 23.2% and 29.2% respectively (21,22).

In contrast, very high prevalences have been reported in South Africa. Indeed, the prevalence of BV is higher in South East Africa than in West Africa (21,23). A study by Myer et al. in South Africa reported that the prevalence of BV was 58.3% in 2002 (21).

Regions such as Australia and New Zealand and Western Europe have the lowest prevalence of BV. The prevalence of BV in Australia was determined to be 4.7% in 2008 (24); a relatively low prevalence of BV was also reported in Finland for the year 2008, at 8.6% (21,25).

The literature has not yet explored factors that predict intranational and international differences in the prevalence of BV. Cultural factors may play a role in the observed differences. In addition, there may be differences in the surveillance techniques used and BV may not be a notifiable disease in all countries. Diagnostic techniques vary according to the availability of resources. Approximately 50% of cases of BV are asymptomatic (13). Due to variations in clinical guidelines, BV may go undetected and the actual prevalence of BV within a country and the reported prevalence would not be the same (11).

Cases of BV were more frequent among women in the [30-40] and [20-30] age groups, with 27.7% and 25.5% respectively, and a statistically significant difference ( $p=0.0001$ ) was found. These results are in agreement with those found by some authors such as Ranjit, et al. in 2018 who found the highest number of BV cases in the 30-40 year age group (8.8%) and the lowest number of BV cases was observed in patients in the 10-20- and 50-60-year age groups (1.3%) (19). The results of this study are also similar to those reported in the studies by Asiegbu et al. in 2018, who found the highest rate of BV (69.1%) in the 25-34 age group (20).

Indeed, BV is a condition that mainly affects sexually active young women, but can occur in the absence of sexual intercourse (4).

This is not consistent with the study by Adinma et al (Nigeria) who found a prevalence of 17.0% for 16-20 year olds (26) and Larsson (Sweden) 18-25 year olds (27).

The highest prevalence of BV has been observed in women who are not on contraception compared to women on contraception. Many studies have shown that the use of hormonal

contraceptives is associated with a reduced likelihood of developing BV (19). These studies demonstrate an association between hormonal contraceptive use and the risk of BV, and raise the role of steroid hormones in influencing the vaginal environment in order to protect against the development of BV. The association between BV and contraception varies according to the methods used. Studies have shown that combined oral contraceptives, progestin-only contraceptives and condom use protect against BV (15,19,24,27,28). They postulated that the difference observed between women who used oral contraceptives and those who did not could be attributed to the effect of increased levels of oestrogen, which could potentially promote the growth of specific bacteria responsible for reducing the risk of BV (19). Of note, Ranjit et al. reported that the risk of BV was higher in those who used condoms daily compared to those who sometimes used condoms, but this difference was not statistically significant. They also found that oral contraceptives reduced the risk of BV (19). Conversely, consistent condom use has been reported in other studies to significantly reduce the risk of recurrent BV (1,27). The relationship between BV and intrauterine devices (IUDs) is unclear (29). Some studies have determined that IUDs increase the risk of BV (24) while others have determined that there is a reduced risk (15). However, intravaginal bleeding in the first six months in IUD users was found to be associated with twice the risk of BV (15). Conversely, Joesoef found that BV was significantly associated with IUDs (1). It is important to consider that there may be differences in the type of IUD used in each study (hormone-loaded versus copper-bearing IUDs). The above studies did not distinguish between the types of IUD used. In our study, the association between vaginosis and pregnancy was found to be 28.6% and 18.5% in non-pregnant women. In Senegal, a similar prevalence (28.0%) was found by Ngom-Cisse et al (30). In a study conducted in Lagos, Nigeria, a similar prevalence (26.0%) was observed among pregnant women by Afolabi BB et al (31). Similarly, Gözde İşik observed a prevalence of 27.9% in Turkey (32).

Other studies have reported lower prevalences in pregnant women, notably in Nigeria and Thailand, where prevalences of 16.6% and 19.3% respectively have been found (33,34). According to some authors, BV increases the risk of miscarriage, preterm labour, premature delivery, chorioamnionitis and postpartum complications such as endometritis (33).

VB was more frequent in nulliparous and primiparous women (23.7%). This was reported in Senegal by Ngom-Cissé who found more VB in primiparous women (31.25%) than in multiparous women (25.45%) without any statistically significant difference (30). These results are corroborated by other studies, notably by Achondou AE et al. in 2016 in Cameroon (35) and by Konadu DG et al. in 2015 in Ghana (36).

This high rate of VB in nulliparous and uniparous women can also be attributed to the increased frequency of intercourse and douching for these women (37). This is not in line with most authors. Indeed, the epidemiological profile of women with BV is similar to that of women with STIs: BV is significantly associated with having a recent sexual partner or multiple sexual partners (38). In the study by Ghattargi et al, BV was more common in multiparous women with a statistically significant difference (39).

Indeed, other previous studies have reported that multiparous women are the group most affected by bacterial vaginosis (39,40). Although multiparous women are normally more sexually active, with social change the age of first intercourse is becoming earlier and young people are engaging in more sexual activity with increasingly sophisticated methods of protection.

We found a higher frequency of BV in women without a history of genital infections with no statistically significant difference.

BV is associated with severe upper genital tract sequelae. Women who are more sexually active and shower more often have more vaginal infections that leave a "breeding ground" for bacteria. The carriage of BV is also favoured by sexual contact with penetration, but also by digital-genital contact, which are practices that favour infections (41).

While contributing to the strengthening of biological and epidemiological knowledge about BV, our study nevertheless has some limitations. Indeed, the clinical information of the patients as well as their intimate habits could help to better understand the occurrence of BV.

## **V. Conclusion**

This work has allowed us to find a fairly high prevalence of BV and to study the potential involvement of certain factors in its occurrence. Among the factors suspected in our study, only the association between age and bacterial vaginosis was found. Knowledge of the factors that contribute to BV and an understanding of its pathophysiology will help to raise women's awareness in order to limit its occurrence and avoid complications in women at risk.

## **VI. Ethical approval**

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

## **References**

1. Makella S Coudray ,Purnima Madhivanan. Bacterial Vaginosis - A Brief Synopsis of the Literature, Eur J Obstet Gynécol Reprod Biol. 2020 février ; 245 : 143–148.
2. J.-P. Menard, F. Bretelle. Gynécologie Obstétrique & Fertilité 40 (2012) 48–54.
3. Eckert LO. Clinical practice. Acute vulvovaginitis., N Engl J Med 2006;355:1244–52.

4. J D Sobel. Bacterial vaginosis, *Annu Rev Med.* 2000;51:349-56.
5. Bohbot JM, Lepargneur JP. La vaginose en 2011 : encore beaucoup d'interrogations, *Gynecologie Obstetrique & Fertilité.* 2012 ; 40 31–36.
6. G. Brabant. Vaginose bactérienne et prématurité spontanée. *Journal de Gynecologie Obstetrique et Biologie de la Reproduction* ,2016 ;45,1247 —1260.
7. Allsworth JE, Peipert JF. « Prevalence of bacterial vaginosis: 2001-2004 National Health and Nutrition Examination Survey data », *Obstet Gynecol.* 2007; vol. 109, n° 1, p. 114-20. (PMID 17197596, DOI 10.1097/01.AOG.0000247627.84791.91, lire en ligne [archive] [html]).
8. Herrero DR, Domingo AA. Bacterial vaginosis, *Enferm Infecc Microbiol Clin.* 2016 Jul;34 Suppl 3:14-8. doi: 10.1016/S0213-005X(16)30214-2.
9. Ellington K, Saccomano SJ. Recurrent bacterial vaginosis, *Nurse Pract.* 2020 Oct;45(10):27-32. doi: 10.1097/01.NPR.0000696904.36628.0a.
10. Questea EB, Montaigne KH, Canis F, Noulard MN, Plennevaux JL, Tilloy E and al. La vaginose bactérienne est-elle facteur de risque de prématurité ? Étude d'une cohorte de 1336 patientes au centre hospitalier d'Arras. *Journal de Gynecologie Obstetrique et Biologie de la Reproduction.* 2012 ; 41, 262—270.
11. Coudray MS, Madhivanan P. Bacterial vaginosis—A brief synopsis of the literature. *Eur J Obstet Gynecol Reprod Biol.* 2020 Feb;245:143-148.
12. Ménard JP. Traitement antibactérien de la vaginose bactérienne : thérapies actuelles et émergentes . *Int J Santé des femmes.* 2011; 3: 295–305.
13. Hay P. Vaginose bactérienne . *Médecine .* 2014; 42 (7):359–63.
14. Schwebke JR. Vaginose bactérienne asymptomatique : réponse au traitement . *Suis J Obstet Gynecol .* 2000 ; 183 ( 6 ):1434–9.
15. Madden T, Grentzer JM, Secura GM, Allsworth JE, Peipert JF. Risque de vaginose bactérienne chez les utilisatrices du dispositif intra-utérin : une étude longitudinale . *Maladies sexuellement transmissibles.* 2012; 39 (3):217–22.
16. Marshall AO. Gestion de la vaginose bactérienne récurrente : informations pour les prestataires occupés . *Sex Med Rev.* 2015; 3 (2):88–92.
17. Martin DH, Marrazzo JM. Le microbiome vaginal : compréhension actuelle et orientations futures. *Le Journal des maladies infectieuses .* 2016 ; 214 Supplément 1 :S36–41.
18. Schwebke JR, Desmond R. Natural history of asymptomatic bacterial vaginosis in a high-risk group of women. *Sexually transmitted diseases.* 2007;34(11):876-7.

19. Ranjit E, Raghubanshi BR, Maskey S, Parajuli P. Prevalence of Bacterial Vaginosis and Its Association with Risk Factors among Nonpregnant Women: A Hospital Based Study. *Int J Microbiol.*2018;2018:8349601.
20. Asiegbu OG, Asiegbu UV, Onwe B, Iwe ABC. Prevalence of Bacterial Vaginosis among Antenatal Patients at Federal Teaching Hospital Abakaliki, South East Nigeria, *Open Journal of Obstetrics and Gynecology* > Vol.8 No.1, January 2018. DOI: 10.4236/ojog.2018.81010
21. Kenyon C, Colebunders R, Crucitti T. The global epidemiology of bacterial vaginosis: a systematic review. *Am J Obstet Gynecol.* 2013;209(6):505-23.
22. Mohammadzadeh F, Dolatian M, Jorjani M, Alavi Majd H. Diagnostic value of Amsel's clinical criteria for diagnosis of bacterial vaginosis. *Glob J Health Sci.* 2014;7(3):8-14.
23. Bautista CT, Wurapa E, Sateren WB, Morris S, Hollingsworth B, Sanchez JL. Bacterial vaginosis: a synthesis of the literature on etiology, prevalence, risk factors, and relationship with chlamydia and gonorrhoea infections. *Mil Med Res.* 2016 Feb 13;3:4. doi: 10.1186/s40779-016-0074-5. eCollection 2016.
24. Brotman RM. Microbiome vaginal et infections sexuellement transmissibles : une perspective épidémiologique. *Le Journal d'investigation clinique.* 2011;121(12):4610–7.
25. Fethers KA, Fairley CK, Morton A, Hocking JS, Hopkins C, Kennedy LJ, et al. Early sexual experiences and risk factors for bacterial vaginosis. *The Journal of infectious diseases.*2009;200(11):1662-70.
26. Adinma JIB, Okoroli NR, Agbai A, Unaeze N. Prévalence de Gardnerella vaginalis chez les femmes nigérianes enceintes. *Journal africain de la santé reproductive.*2001; 5, 50-55.
27. Larsson PG, Fahraeus L, Carlsson B, Jakobsson T, Forsum U. (2007) Facteurs prédisposants à la vaginose bactérienne, efficacité du traitement et résultat de la grossesse parmi les accouchements à terme ; Résultats d'une étude sur l'accouchement prématuré. *Santé des femmes BMC.*2007 ;7, 20.
28. Ménard JP. Traitement antibactérien de la vaginose bactérienne : thérapies actuelles et émergentes. *Int J Santé des femmes.* 2011;3:295–305.
29. Cook RL, Redondo-Lopez V, Schmitt C, Meriwether C, Sobel JD. Facteurs cliniques, microbiologiques et biochimiques de la vaginose bactérienne récurrente. *Journal de microbiologie clinique.*1992;30(4):870–7.
30. Ngom-Cisse S, Dieng A, Gassama O, Diagne-Samb H, Cisse M, Ndiaye SML et al. Risk Factors Associated with Bacterial Vaginosis in Pregnant Women in Senegal. *Microbiol. Res. J. Int.*, vol. 32, no. 8, pp. 30-39, 2022.

31. Afolabi BB, Moses OE, Oduyebo OO. Bacterial vaginosis and pregnancy outcome in Lagos, Nigeria. *Open Forum Infect Dis.* 2016; 3(1): ofw030. PubMed | Google Scholar.
32. Isik G, Demirezen S, Dönmez HG, Beksaç MS. Bacterial vaginosis in association with spontaneous abortion and recurrent pregnancy losses. *J Cytol.* 2016; 33(3): 135-40. PubMed | Google Scholar.
33. Aduloju OP, Akintayo AA, Aduloju T. Prevalence of bacterial vaginosis in pregnancy in a tertiary health institution, south western Nigeria. *Pan African Medical Journal.* 2019; 33:9. doi:10.11604/pamj.2019.33.9.17926.
34. Tachawatcharapunya S, Chayachinda C, Parkpinyo N. The Prevalence of Bacterial Vaginosis in Pregnant Women During Early Third Trimester and the Pregnancy Complications. *Thai J Obstet Gynaecol.* 2017; 25, 95-102.
35. Achondou AE, Fumoloh FF, Aseneck AC, Awah AR, Utokoro AM. Prevalence of bacterial vaginosis among sexually active women attending the CDC central clinic tiko, South West Region, Cameroon. *African J Infect Dis.* 2016;10: 96–101.
36. Konadu DG, Owusu-Ofori A, Yidana Z, , Boadu F, Iddrisu LF, Gyasi DA et al. Prevalence of vulvovaginal candidiasis, bacterial vaginosis and trichomoniasis in pregnant women attending antenatal clinic in the middle belt of Ghana. *BMC Pregnancy Childbirth.* 2019 Sep 23;19(1):341. doi: 10.1186/s12884-019-2488-z.
37. Omole-Ohonsi A, Nwokedi EE. Sociodemographic Characteristics and Aetiological Factors of Vaginal Discharge in Pregnancy. *Jos Journal of Medicine,* 2011. Vol. 5 No. 2. DOI: 10.4314/jjm.v5i2.70690.
38. Fethers KA, Fairley CK, Hocking JS, Gurrin LC, Bradshaw CS. Sexual risk factors and bacterial vaginosis: a systematic review and meta-analysis. *Clin Infect Dis* 2008;47(11):1426–35.
39. Ghattargi S, Sheela N, Dias M. Prevalence and risk factors for bacterial vaginosis in sexually active females in age group 20-45 years and comparison of Amsel's criteria with Nugent's score. *Int J Reprod Contracept Obstet Gynecol.* 2018; 7(9): 3478-3484.
40. Ibrahim SM, Bukar M, Galadima GB, Audu BM, Ibrahim HA. Prevalence of bacterial vaginosis in pregnant women in Maiduguri, North-Eastern Nigeria. *Nigerian Journal of Clinical Practice* • Mar-Apr 2014 • Vol 17 • Issue 2.
41. Verstraelen H, Verhelst R, Vaneechoutte M, Temmerman M. The epidemiology of bacterial vaginosis in relation to sexual behaviour. *BMC Infect Dis.* 2010; 10, 81. <https://doi.org/10.1186/1471-2334-10-81>.