

Standardization of plant growing media and seasons for cuttings of Kusum [*Schleichera oleosa* (Lour.) Oken]: A major challenge

Comment [f1]: Under line and Italic

Comment [f2]: Not necessary in here

ABSTRACT

Comment [f3]: Under line

Growing cuttings of Kusum (*Schleichera oleosa*), is a major hurdle for the Lac growers in India. This species is primarily known as the lac host tree for lac production besides other multiple uses. Due to high population pressure and market demand, the natural population and genetic diversity of this species is declining day-by-day. Maintaining the selected best germplasm through clonal propagation is the need of the hour. Mass multiplication through cuttings could only be the right choice of the grower if found successful. But, no successful cuttings of Kusum have been reported till date. Therefore, an experiment was carried out during 2018-21 following Completely Randomized Design (CRD) to standardize the plant growing in here media and the seasons for growing cuttings at the very initial stages of the experiment. The result revealed that the root, shoot and leaf growths in cuttings of Kusum are significantly influenced by the different plant growing media. The maximum rooting initiation ($43.33 \pm 0.96\%$) was observed in the cuttings grown in media of Coco peat: FYM: Vermicompost (2:2:1). The maximum rooting ($44.50 \pm 1.85\%$) in the cuttings prepared from seedlings of less than 5 years old and $33.75 \pm 1.49\%$ rooting in the cuttings prepared from 10-20 years old plantation, were recorded in the month of May. Based on the present investigation, it is concluded that the plant growing media having the composition of Coco peat: FYM: Vermicompost (2:2:1) is more appropriate and recommended for growing cuttings of Kusum. Moreover, the month of May is highly recommended for growing cuttings of Kusum for maximum success. Regarding selection of mother plant, it is advised to select the Kusum plant from the plantation aged less than five years old.

Comment [f4]: Coco peat. FYM and vermicompost

Comment [f5]: Like a above

Key words: Kusum; *Schleichera oleosa*; clonal propagation; vegetative propagation; cuttings; plant growing media; season

Comment [f6]: delet

Comment [f7]: Key word: The keywords section is inconsistent with the text and it is better to replace the keywords

INTRODUCTION

In India, the species *Schleichera oleosa* (Lour.) Oken (syn., *Pistacia oleosa* Lour., *S. trijuga* Willd., *Cussambium oleosum* Kuntze), belongs to the family Sapindaceae, locally known as Kusum, is primarily known as the lac host tree for lac production, which is also known to be exploited for multiple uses including ethnobotanical purpose (Guleria and Vaidya, 2015; Goswami and Singh, 2017; Sarkar et al., 2017a). Growing cuttings of Kusum, is a major hurdle for the Lac growers in India. The natural population and genetic diversity of this species is declining day-by-day due to high population pressure, industrialization, modernization and market demand (Sarkar et al., 2017b). Conservation of such demanding species and also for maintaining the selected best germplasm through clonal propagation is the need of the hour. Mass production of seedlings of this species is a major challenge as it is a

Comment [f8]: Under line

Comment [f9]: ???

Comment [f10]: It's not necessary

39 cross pollinated species that leads to heterozygosity in the plants (Radhamani *et al.*, 2003).
40 Genetic improvement through this auto-generated heterozygous plants may call for long term
41 experiments and may require high monetary investments.

42 Mass multiplication through the most common method like cuttings could only be the
43 most alternative, reliable and quicker approaches for the lac growers if found successful. Bhatt
44 and Todaria (1990a & b), Chauhan *et al.* (1993) and Chauhan *et al.* (1997) reported that,
45 different concentration of auxins and seasonal variation play a crucial role in root initiation in
46 branch cuttings of different woody species. Moreover, the level of organic matter content,
47 water holding capacity (Sabir *et al.*, 2004) and the level of nitrogen and potassium in rooting
48 media (Sengel *et al.*, 2012) effect the growth of cuttings. Palanisamy and Kumar (1997)
49 reported that exogenously applied auxins are sensitive to activate the cambium probably in the
50 active period of cambium resulting significant root formation. Smith and Wareing (1972)
51 stated that the endogeneous IAA, photoperiod and temperature apparent to control the vascular
52 tissue more particularly the cambial activity.

Comment [f11]: From

53 The growing media is generally considered to be one of the vital inputs for rooting
54 initiation (Dolor *et al.*, 2009). Rooting media should be selected on the basis of criteria, i.e.,
55 availability and cost of rooting media components (Hartmann *et al.*, 2007). Peat moss is now a
56 commonly used media, due to its properties of homogeneity and quality added to media, but an
57 expensive media too (Abdel-Mohsen, 2015). Many literatures revealed that, growing media
58 has an important role on success of cuttings and thereby influences the survival, shooting,
59 rooting and other growth characteristics of any plant (Aamir *et al.*, 2019). Shah *et al.* (2013)
60 suggested sawdust as an important growing media for quicker sprouting in the cuttings of *Ficus*
61 *binnendijkii*. Experiment on effect of different media of cuttings on rooting of guava (*Psidium*
62 *guajava* L.) revealed that soil loam and sawdust, except for shoot number and root diameter,
63 had the lowest positive effect on guava rooting (Sardoei, 2014). In another experiment, carried
64 out by Mabizela *et al.* (2017) revealed that the maximum rooting and survival percentage, root
65 number and root length in the cuttings of honeybush (*Cyclopia subternata*) was due to the
66 treatments of bark mix and 3mix as rooting media and Seradix B2 and Seradix B3 as plant
67 growth regulators.

Comment [f12]: Under line

68 Though there are ample successful methods of growing cuttings in other woody species
69 available, but no successful cuttings have been reported so far in this Kusum tree. The problem
70 of rooting in cuttings might be due to various internal and external factors like **type of rooting**

71 substrate or plant growing media, growth regulator treatments, age of the source plant,
72 pre-treatments and the season during which the cuttings are collected (Hartmann *et al.*,
73 2002; Soundy *et al.*, 2008). Considering the above all facts, an experiment was conducted to
74 standardize the plant growing media and the seasons for growing cuttings of Kusum at the very
75 initial stages of the experimental investigation.

76 2. Methodology

77 The experiment was conducted at research fields of ICAR Research Complex for
78 Eastern Region, Farming System Research Centre for Hill and Plateau Region, Plandu, Ranchi
79 (located at 23° 17' 17.1" North latitude and 85° 24' 34.79" East longitude) and Genetics and
80 Tree Improvement Division, Institute of Forest Productivity (IFP), Lalgutwa, Ranchi (located at
81 23° 21' 28.37" North latitude and 85° 14' 41.43" East longitude). The study area comes under
82 dry and humid tropical climate, received a maximum average rainfall of 401.95 mm during the
83 month of August with a mean monthly highest temperature ranged from 24.0 °C in January to
84 37.5 °C in May and the average monthly lowest temperature ranged from 7.4 °C in January to
85 27.6 °C in May. The relative humidity of the experimental area increased with the South-west
86 monsoon's onset and it became more than 80% in July. The relative humidity was lowest
87 during summer (*i.e.*, < 50%).

88 2.1. Vegetative propagation through cuttings

89 The cuttings were collected throughout the year (during each month) from the
90 available trees and from selected candidate plus trees (CPTs) of Institute of IFP, Ranchi and
91 ICAR Indian Institute of Natural Resins and Gums (IINRG), Namkum, Ranchi during March,
92 2018 to August, 2021. The defoliation of shoots was carried out at least seven days before the
93 collection of cuttings. The cuttings about 6-12 cm length and 0.5 – 2.5 cm in diameter having
94 2-3 buds were collected by using sharp Secateurs. The cuttings were then washed in 0.2%
95 bavistin solution prior to hormonal treatment [dipped in 0.1 % Ascorbic Acid for 1 hour
96 followed by IBA (2000 ppm) + 5 % Sucrose for 1 hour] to prevent attack of pathogens.

Comment [f13]: Available and selected candidate plus trees

Comment [f14]: Is it fungicide and if yes it's better to write scientific name and where manufacture in parenthesis

97 The cuttings were planted either in the hycopots or root trainers filled with rooting
98 medium (sand) or in sand bed immediately after treatment with different plant growth
99 hormones. Two-third length of the cuttings were inserted in the rooting medium and arranged
100 in the mist chamber or shade-net or poly-tunnel according to Completely Randomized Design
101 (CRD). The planted cuttings were irrigated and weeded as and when required, until the

Comment [f15]: ????

102 termination of the experiment. The procedure for cuttings in Kusum followed in this
103 experiment has been described under the following sub-heads:

104 **2.1.1. Standardization of plant growing media**

105 The plant growing media was standardized using the comparative test of plant growth
106 performance based on the:

107 *(i) Effect of different plant growing media on shoots and leaf growth in cuttings of Kusum,*
108 *and*

109 *(ii) Effect of different plant growing media on root growth and survival of cuttings of Kusum*

110 The treatments details are given in **Table 1**. A total of 8 treatments with three replications
111 consisting of 100 cuttings per replication per treatment following Completely Randomized
112 Design (CRD) were considered. The parameters like **days at first bud initiation, number of**
113 **shoots per cutting, number of leaves per cutting, average shoot length (cm), average leaf**
114 **length (cm), average leaf width (cm), leaf length-width ratio, rooting initiation (%)**,
115 **number of secondary roots per cutting, root spread (cm), total root length (cm) and plant**
116 **survival period (in days) were recorded.**

Comment [f16]: You can write just about a treat
you were measured in this paper and the other
parameter not necessary

117 **Table 1: Different proportion of plant growing media**

Treatment details	Treatment code
Soil : Sand : FYM (1:1:1)	T ₁
Soil : Sand : Vermicompost (2:2:1)	T ₂
Soil : Sand : FYM : Vermicompost (4:4:2:1)	T ₃
Coco peat : Vermicompost (8:1)	T ₄
Coco peat : FYM : Vermicompost (2:2:1)	T ₅
Sand	T ₆
Coco peat	T ₇
Sand : Soil rite (2:1)	T ₈

118

119 **2.1.2 Seasons of growing cuttings**

120 The cuttings of young shoots of Kusum were collected during each month from
121 seedlings of less than 5 years old as well as plantations of 10-20 years old and planted in the
122 sand bed under poly-tunnel maintaining the temperature as 28 - 30 °C and humidity as 70 – 80
123 % by manual spraying. The parameters like rooting initiation (%), rooting success (%) and

124 survival (%) were recorded. A total of 12 treatments with three replications consisting of 30
125 cuttings per replication per treatment following Completely Randomized Design (CRD) were
126 considered.

127 **2.2. Data analysis**

128 The data were subjected to analysis of variance (ANOVA) to quantify the differences
129 among applied treatments by using OPSTAT software (Sheoran *et al.*, 1998). Data
130 transformation was also carried out as and when required to satisfy the ANOVA requirements.
131 The data that have been transformed were expressed in original units for presentation in the
132 Tables. Treatment means were compared from the estimated values of least significant
133 difference (L.S.D) at 5 per cent level of significance for the error degree of freedom and
134 coefficient of variation between different treatments was also calculated. Moreover, the
135 Duncan's Multiple Range (DMR) Test was also done to compare larger pairs of means.

Comment [f17]: If you have a control plot its benefit Otherwise, it will not make a difference

136 **3. Results and Discussion**

137 The results have been explained under the following sub-heads:

138 **3.1. Effect of different plant growing media on shoots and leaf growth in cuttings of Kusum**

139 Data in **Table 2** evinced that the shoots and leaf growth in cuttings of Kusum are
140 significantly ($P \leq 0.05$) influenced by the different plant growing media (**Fig. 1 & 2**). The
141 maximum day required for first bud initiation (7.33 ± 0.34) was observed in the cuttings grown
142 in Coco peat: Vermicompost (8:1) media, whereas the minimum as 3.33 ± 0.28 and 3.33 ± 0.29
143 were observed under the plant growing media of Coco peat and Sand: Soilrite (2:1),
144 respectively. The number of shoots per cutting ranged from 1.13 ± 0.13 to 1.86 ± 0.11 . The
145 maximum number of shoots (1.86 ± 0.11) was observed in the plant growing media of Soil:
146 Sand: Vermicompost (2:2:1), whereas the minimum (1.13 ± 0.13) was recorded in case of the
147 cuttings grown in Coco peat media. Similarly, the number of leaves per cutting was recorded
148 maximum as 5.08 ± 0.11 in case of cuttings grown in Coco peat : FYM : Vermicompost (2:2:1)
149 media and the minimum as 2.45 ± 0.24 was recorded in the cuttings grown in Soil : Sand :
150 FYM (1:1:1). The maximum average shoot length (12.35 ± 0.88 cm) value was observed in the
151 cuttings grown in Coco peat: FYM: Vermicompost (2:2:1) media, whereas the minimum
152 average shoot length (3.42 ± 0.30 cm) value was recorded in the cuttings grown in Soil : Sand :
153 FYM (1:1:1) media. Similarly, the maximum average leaf length (11.33 ± 0.18 cm) was

154 recorded in the cuttings grown in Coco peat : FYM : Vermicompost (2:2:1) media, whereas the
155 minimum average leaf length (3.67 ± 0.19 cm) was observed in the cuttings grown in Soil :
156 Sand : FYM (1:1:1). In case of average leaf width, the maximum (5.63 ± 0.16 cm) was
157 recorded in the cuttings grown in Coco peat : FYM : Vermicompost (2:2:1) media, but the
158 minimum (1.96 ± 0.03 cm) was recorded in the cuttings grown in Soil : Sand : FYM (1:1:1)
159 media. Data further revealed that, the leaf width ratio was found to be maximum as 2.21 ± 0.01
160 in the cuttings grown in Coco peat media, whereas the minimum as 1.55 ± 0.10 in case of the
161 cuttings grown in Soil : Sand : Vermicompost (2:2:1). Similar findings in regards to the effect
162 and influence of plant growing media on shoot and leaf growth of other woody plants were also
163 being reported by [Aamir et al. \(2019\)](#). The influence of different plant growing media on
164 growth of cuttings might be due to the level of organic matter content, water holding capacity
165 ([Sabir et al., 2004](#)) and the level of nitrogen and potassium in rooting media ([Sengel et al.,](#)
166 [2012](#)). Earliness in sprouting and increase in number of sprouts may be due to better utilization
167 of stored carbohydrates, nitrogen and other factors with the help of growth regulators ([Wright,](#)
168 [1975](#); [Chandramouli, 2001](#); [Sahoo et al., 2014](#)). While propagating *Cordia africana* species
169 through cuttings, [Ambebe et al. \(2018\)](#) suggested the plant growing media as sand and
170 sawdust: sand (1:1) as the best suitable alternative for getting maximum spouting and growth of
171 cuttings.

Comment [f18]:

Comment [f19R18]: Under line


Comment [f20]: were

Table 2: Effect of different plant growing media on shoots and leaf growth in cuttings of Kusum

Plant growing media	Days at first bud initiation	No. of shoots per cutting	No. of leaves per cutting	Average shoot length (cm)	Average leaf length (cm)	Average leaf width (cm)	Leaf length width ratio
Soil : Sand : FYM (1:1:1)	6.00 ± 0.58 ^b	1.17 ± 0.11 ^b	2.45 ± 0.24 ^c	3.42 ± 0.30 ^c	3.67 ± 0.19 ^g	1.96 ± 0.03 ^t	1.87 ± 0.07 ^b
Soil : Sand : Vermicompost (2:2:1)	6.00 ± 0.01 ^b	1.86 ± 0.11 ^a	4.64 ± 0.32 ^a	5.52 ± 0.29 ^b	4.73 ± 0.43 ^t	3.04 ± 0.10 ^d	1.55 ± 0.10 ^c
Soil : Sand : FYM : Vermicompost (4:4:2:1)	5.33 ± 0.33 ^{bc}	1.29 ± 0.13 ^b	2.76 ± 0.14 ^{bc}	5.60 ± 0.30 ^b	6.06 ± 0.17 ^d	3.64 ± 0.17 ^c	1.67 ± 0.04 ^c
Coco peat : Vermicompost (8:1)	7.33 ± 0.34 ^a	1.33 ± 0.08 ^b	2.67 ± 0.20 ^c	4.22 ± 0.42 ^c	5.44 ± 0.30 ^e	2.85 ± 0.11 ^e	1.91 ± 0.04 ^b
Coco peat : FYM : Vermicompost (2:2:1)	3.67 ± 0.33 ^d	1.85 ± 0.10 ^a	5.08 ± 0.11 ^a	12.35 ± 0.88 ^a	11.33 ± 0.18 ^a	5.63 ± 0.16 ^a	2.01 ± 0.02 ^b
Sand	4.33 ± 0.32 ^{cd}	1.67 ± 0.07 ^a	3.33 ± 0.17 ^b	5.98 ± 0.19 ^b	8.35 ± 0.32 ^b	4.28 ± 0.12 ^b	1.95 ± 0.03 ^b
Coco peat	3.33 ± 0.28 ^d	1.13 ± 0.13 ^b	2.50 ± 0.10 ^c	5.67 ± 0.17 ^b	6.89 ± 0.06 ^c	3.12 ± 0.03 ^d	2.21 ± 0.01 ^a
Sand : Soil rite (2:1)	3.33 ± 0.29 ^d	1.22 ± 0.07 ^b	2.82 ± 0.09 ^{bc}	6.65 ± 0.19 ^b	6.91 ± 0.07 ^c	3.56 ± 0.06 ^c	1.94 ± 0.02 ^b
F- ratio	18.64	7.57	29.33	40.20	276.41	396.11	21.43
d.f.	14	14	14	14	14	14	14
S.E. (± d)	0.48	0.16	0.27	0.60	0.20	0.08	0.06
L.S.D. (P ≤ 0.05)	1.05	0.34	0.58	1.30	0.44	0.17	0.13
C.V. (%)	12.05	13.25	9.92	11.92	3.70	2.71	4.01

Data is shown as mean values of variables ± SE. The Means in a column followed by different superscript letters are significantly ($P \leq 0.05$) different using Duncan's multiple range test (DMRT).

Comment [f21]: It's not necessary to be Italic



a. Performances of cuttings planted on sand bed under shade-net

b. Performances of cuttings planted on root trainers under manually operated mist chamber

c. Performances of cuttings planted on sand bed under closed poly-tunnel

Fig. 1: Growth performances of cuttings (a – c)



a. Shoot initiation on cuttings under different plant growing media



b. Rooting in cuttings of Kusum

Fig. 2: Shoot and root growth performances of cuttings (a & b)

3.2. Effect of different plant growing media on root growth in cuttings of Kusum

The root growth in cuttings of Kusum was influenced significantly by different plant growing media (**Table 3, Fig. 2**). The maximum rooting initiation (43.33 ± 0.96 %) was observed in the cuttings grown in media of Coco peat : FYM : Vermicompost (2:2:1), whereas no root initiation was observed in the cuttings grown in Soil : Sand : FYM (1:1:1) media. The number of secondary roots per plant ranged from 0 to 4.67 ± 0.31 . The maximum number of secondary roots (4.67 ± 0.31) was observed in the plant growing media of Coco peat : FYM : Vermicompost (2:2:1), whereas no secondary roots were observed in case of the cuttings grown in Soil : Sand : FYM (1:1:1) media. In case of root spread, the maximum as 2.82 ± 0.07 cm was recorded in the cuttings grown in Coco peat : FYM : Vermicompost (2:2:1) media, followed by 2.22 ± 0.07 cm in case of cuttings grown in sand media. The total root length was found to be maximum as 10.45 ± 0.46 cm in the cuttings grown in Coco peat : FYM : Vermicompost (2:2:1) media, followed by 10.21 ± 0.25 cm in Sand media and 6.14 ± 0.46 cm in the Coco peat media. The plant survival period was also recorded and found to be varied significantly in all the plant growing media. The maximum plant survival period (90.00 ± 6.08 days) was recorded in the cuttings grown in Coco peat : FYM : Vermicompost (2:2:1) media, whereas the minimum plant survival period (42.33 ± 1.45 days) was recorded in the cuttings grown in Soil : Sand : FYM (1:1:1) media. Similar findings in regards to the effect and influence of plant growing media on root growth and survival of any plant were also being reported by [Ambebe et al. \(2018\)](#) and [Aamir et al. \(2019\)](#). [Sardoei \(2014\)](#) also had reported the highest rooting success in Guava when grown in sand. Similarly, the combination of plant growing media like garden soil, sand, FYM and peat moss, was reported as the best in terms of survival and rooting of cuttings in Olive trees ([Aamir et al., 2019](#)).

Table 3: Effect of different plant growing media on root growth in cuttings of Kusum

Plant growing media	Rooting initiation ^a (%)	No. of secondary roots per cutting	Root spread (cm)	Total root length (cm)	Plant survival period (in day)
Soil : Sand : FYM (1:1:1)	0.00 ± 0.00 (0.00 ± 0.00) ^f	0.00 ± 0.00 ^e	0.00 ± 0.00 ^f	0.00 ± 0.00 ^e	42.33 ± 1.45 ^e
Soil : Sand : Vermicompost (2:2:1)	3.33 ± 0.38 (10.48 ± 0.62) ^e	1.67 ± 0.33 ^d	1.33 ± 0.06 ^e	2.39 ± 0.54 ^d	47.33 ± 3.18 ^{de}
Soil : Sand : FYM : Vermicompost (4:4:2:1)	10.00 ± 0.77 (18.40 ± 0.74) ^d	2.67 ± 0.32 ^{bc}	1.61 ± 0.03 ^d	4.53 ± 0.5 ^c	55.33 ± 2.03 ^{cd}
Coco peat : Vermicompost (8:1)	3.33 ± 0.19 (10.51 ± 0.31) ^e	2.33 ± 0.33 ^{cd}	1.81 ± 0.06 ^e	4.48 ± 0.13 ^c	63.00 ± 4.04 ^{bc}
Coco peat : FYM : Vermicompost (2:2:1)	43.33 ± 0.96 (41.15 ± 0.56) ^a	4.67 ± 0.31 ^a	2.82 ± 0.07 ^a	10.45 ± 0.46 ^a	90.00 ± 6.08 ^a
Sand	16.67 ± 1.02 (24.07 ± 0.78) ^c	4.33 ± 0.30 ^a	2.22 ± 0.07 ^b	10.21 ± 0.25 ^a	71.33 ± 2.03 ^b
Coco peat	20.00 ± 1.02 (26.54 ± 0.74) ^b	3.33 ± 0.29 ^b	1.75 ± 0.07 ^{cd}	6.14 ± 0.46 ^b	58.00 ± 3.22 ^{cd}
Sand : Soilrite (2:1)	20.00 ± 1.73 (26.51 ± 1.24) ^b	2.33 ± 0.33 ^{cd}	1.67 ± 0.05 ^{cd}	4.10 ± 0.46 ^c	50.67 ± 1.45 ^{de}
F- ratio	328.31	26.05	197.36	99.59	19.74
d.f.	14	14	14	14	14
S.E. (± d)	0.99	0.41	0.08	0.51	4.85
L.S.D. (P ≤ 0.05)	2.15	0.90	0.18	1.10	10.49
C.V. (%)	6.17	18.97	6.02	11.81	9.93

Data is shown as mean values of variables ± SE. The figures in parentheses are arcsine transformed values which were compared based on its derived *F-ratio*, *d.f.*, *S.E.* (± *d*), *L.S.D.* ($P \leq 0.05$) and *C.V.* (%) as mentioned in its corresponding column. Means in a column followed by different superscript letters are significantly ($P \leq 0.05$) different using DMR test.

Comment [f22]: Why use both two number

3.3. Seasonal effect on rooting success and per cent survival in cuttings

A perusal of data presented in **Table 4** revealed that there were significant ($P \leq 0.05$) effects of different seasons or months on rooting success in the cuttings prepared from less than 5 years old trees. Season in which cuttings are taken, can also effect the rooting of stem cuttings (Mabizela *et al.*, 2017). In the present investigation, there was significant effect of different seasons on per cent rooting success in the cuttings in Kusum. The maximum rooting initiation (48.50 ± 2.10 %) and rooting success (44.50 ± 1.85 %) in the cuttings prepared from seedlings of less than 5 years old and 10 - 20 years old Kusum plantation were recorded in the month of May. Similar effect was also reported by Bhatt and Todaria (1990a & b), Chauhan *et al.* (1993), Chauhan *et al.* (1997), Agbo and Obi (2008) and Khosla and Pant (2009) who stated that, different concentration of auxins and seasonal variation play a crucial role in root initiation in branch cuttings of different woody species. Naveen (2002) also reported a better sprouting and rooting success in *Hippophae rhamnoides* when planted in August than in March. Shamet and Naveen (2005) suggested rainy season (July - August) as the best time for maximum cutting success in *Celtis australis*. But, spring season (March – April) was the best time for the cuttings of *Acer acuminatum* to obtain maximum rooting success (Kumar *et al.*, 2018).

Comment [f23]: Under line

Comment [f24]: underline

Comment [f25]: under line

Table 4: Effect of different seasons on rooting success and survival in cuttings of Kusum

Months	Cuttings from seedlings of less than 5 years old		Cuttings from 10-20 years old plantation		Nil
	Rooting initiation (%)	Survival (%)	Rooting initiation (%)	Survival (%)	
January	2.00 ± 0.71 (7.02 ± 2.38) ^g	1.00 ± 0.41 (4.90 ± 1.73) ^{gh}	0.50 ± 0.29 (2.87 ± 1.66) ^f		
February	5.50 ± 0.65 (13.49 ± 0.82) ^f	4.00 ± 0.41 (11.49 ± 0.60) ^f	3.50 ± 0.29 (10.75 ± 0.45) ^e		
March	22.75 ± 1.38 (28.45 ± 0.94) ^d	20.00 ± 0.71 (26.54 ± 0.51) ^d	14.75 ± 0.48 (22.57 ± 0.39) ^d		
April	30.00 ± 1.47 (33.18 ± 0.92) ^c	27.50 ± 1.44 (31.59 ± 0.93) ^c	28.25 ± 1.25 (32.08 ± 0.80) ^b		
May	48.50 ± 2.10 (44.12 ± 1.21) ^a	44.50 ± 1.85 (41.82 ± 1.07) ^a	33.75 ± 1.49 (35.49 ± 0.91) ^a		
June	40.25 ± 1.70 (39.35 ± 0.99) ^b	32.75 ± 1.93 (34.87 ± 1.18) ^b	31.50 ± 1.32 (34.12 ± 0.81) ^{ab}		
July	37.50 ± 2.10 (37.73 ± 1.25) ^b	30.75 ± 1.65 (33.64 ± 1.03) ^{bc}	23.00 ± 0.91 (28.63 ± 0.62) ^c		
August	18.75 ± 1.38 (25.61 ± 1.01) ^d	13.00 ± 0.71 (21.11 ± 0.59) ^e	13.25 ± 0.85 (21.31 ± 0.73) ^d		
September	9.75 ± 0.85 (18.14 ± 0.82) ^e	6.25 ± 0.48 (14.44 ± 0.58) ^f	4.50 ± 0.29 (12.22 ± 0.40) ^e		

October	3.50 ± 0.65 (10.64 ± 1.03) ^f	1.50 ± 0.65 (5.96 ± 2.17) ^g	0.50 ± 0.29 (2.87 ± 1.66) ^f	
November	0.50 ± 0.29 (2.87 ± 1.66) ^h	0.25 ± 0.24 (1.43 ± 1.42) ^h	0.00 ± 0.00 (0.00 ± 0.00) ^f	
December	1.50 ± 0.65 (5.96 ± 2.17) ^{gh}	0.50 ± 0.29 (2.87 ± 1.66) ^{gh}	0.00 ± 0.00 (0.00 ± 0.00) ^f	
F- ratio	144.12	182.59	241.49	-
d.f.	33	33	33	-
S.E. (± d)	1.70	1.49	1.25	-
L.S.D. (P ≤ 0.05)	3.47	3.04	2.56	-
C.V. (%)	10.80	10.94	10.48	-

Data is shown as mean values of variables ± SE. The figures in parentheses are arcsine transformed values which were compared based on its derived *F- ratio*, *d.f.*, *S.E. (± d)*, *L.S.D. (P ≤ 0.05)* and *C.V. (%)* as mentioned in its corresponding column. Means in a column followed by different superscript letters are significantly ($P \leq 0.05$) different using DMR test.

4. Conclusion

The plant growing media and the season of growing cuttings significantly affected the rooting success in cuttings of Kusum. Based on the present investigation, it is concluded that the plant growing media having the composition of Coco peat: FYM : Vermicompost (2:2:1) is more appropriate and recommended for growing cuttings of Kusum. Season also play a key role for successful cuttings. The month of May is highly recommended for growing cuttings of Kusum for maximum success in India, though it may vary from place to place. Regarding selection of mother plant, it is suggested to select the Kusum plant from the plantation aged less than five years old.

6. References

- Aamir SS, Nadeem I, Khan TN, Ullah MA, Ikram Z, Waris A, Ahmed A.** Effect of various growth media on success of Olive cuttings. *Science, Technology and Development*. 2019; 38(1): 25-28. DOI: 10.3923/std.2019.25.28.
- Abdel-Mohsen MA.** Compost as peat substitute in olive cutting media. *Journal of Plant Production*. 2015; 6(8): 1443-1450.
- Agbo CU, Obi IU.** Patterns of vegetative propagation of stem cuttings of three physiological ages of *Gongronema latifolia* Benth over two seasons in Nsukka. *Journal of Tropical Agriculture Food Environment and Extension*. 2008; 7(3): 193-198.
- Ambebe TF, Agbor AEW, Siohdjie CHS.** Effect of different growth media on sprouting and early growth of cutting-propagated *Cordia africana* Lam. *International Journal of Forest, Animal and Fisheries Research*. 2018; 2(1): 28-33.

Comment [f26]: The references part should be written in a format of mag and re check a gain

- Bhatt BP, Todaria NP.** Seasonal rooting behaviour of some agroforestry tree species of Garhwal Himalaya. *Indian Journal of Forestry*. 1990a; 13: 362-364.
- Bhatt BP, Todaria NP.** Vegetative propagation of tree species of Social forestry value in Garhwal Himalaya. *Journal of Tropical Forest Science*. 1990b; 2: 195-210.
- Chandramouli H.** Influence of growth regulators on the rooting of different types of cuttings in *Bursera penicillatai* (DC) Engl. M.Sc. (Agril.) Thesis, University of Agricultural Sciences, Bangalore. 2001; 103 pp.
- Chauhan DS, Bhatt BP, Todaria NP.** Vegetative propagation studies in some tree and shrub species of Garhwal Himalaya. *Indian Journal Plant Physiology*. 1993; XXXVI(2): 112-114.
- Chauhan DS, Bhatt BP, Todaria NP.** Vegetative propagation response of three leguminous taxa of Garhwal Himalaya. *Indian Journal of Forestry*. 1997; 19(4): 377-380.
- Dolor DE, Lkie FO, Nnaji GU.** Effect of propagation media on the rooting of leafy stem cuttings of *Irvingia wombolu* (Vermoesen). *Research Journal of Agriculture and Biological Sciences*. 2009. 5(6): 1146-1152.
- Goswami S, Singh RP.** Ayurvedic, phytochemical and pharmacological review of *Schleichera oleosa* (Lour.) Oken: a traditional plant with enormous biological activity. *World Journal of Pharmaceutical Research*. 2017; 6(10): 295-309. DOI:10.20959/wjpr201710-9370.
- Guleria H, Vaidya M.** Anatomical studies of *Schleichera oleosa* (Lour.) Oken. *World Journal of Pharmaceutical Research*. 2015; 4(12): 1178-1188.
- Hartmann HT, Kester DE, Davies FT, Genev RI.** Plant propagation, principles and practices. Seventh edition. Prentice-hall of India private limited. 2007; p. 880.
- Khosla V, Pant KS.** Seasonal variation of the effect of auxins on rooting of different stem cutting portion of *Jatropha curcas* Linn. *Indian Forester*. 2009; 1: 1185-1192.
- Kumar S, Shamet GS, Kumari N, Hegde N.** Rooting response of *Acer acuminatum* cuttings to IBA, girdling and season. *Indian Journal of Ecology*. 2018; 45(4): 806-809.
- Mabizela GS, Slabbert MM, Bester C.** The effect of rooting media, plant growth regulators and clone on rooting potential of honeybush (*Cyclopia subternata*) stem cuttings at different planting dates. *South African Journal of Botany*. 2017; 110: 75-79. <http://dx.doi.org/10.1016/j.sajb.2016.02.200>.
- Mabizela GS, Slabbert MM, Bester C.** The effect of rooting media, plant growth regulators and clone on rooting potential of honeybush (*Cyclopia subternata*) stem cuttings at different planting dates. *South African Journal of Botany*. 2017; 110: 75-79. <http://dx.doi.org/10.1016/j.sajb.2016.02.200>.

- Naveen CR.** Vegetative propagation of Khirk (*Celtis australis* Linn) and Seabuck thorn (*Hippophae rhamnoides* Linn and *H salicifolia* D) through cuttings. M.Sc. Thesis, Dr .Y. S. Parmar University of Horticulture and Forestry, Solan. 2002; 226 p.
- Palanisamy K, Kumar P.** Seasonal variation on adventitious rooting in branch cuttings of *Pongamia pinnata* (Pierre). Indian Forester. 1997; 122(3): 236-239.
- Radhamani J, Singh AK, Sharma A.** Exploring conventional seed storage in some tropical species. *In: seed conservation: turning science into practice*; Smith, R.D., Dickie, J.B., Linington, S.H., Pritchard, H.W. and Probert, R.J. (eds.). Royal Botanic Gardens, Kew. 2003; pp. 777-784.
- Sabir A, Kara Z, Küçükbasmaç F, Yücel NK.** Effects of different rooting media and auxin treatments on the rooting ability of Rupestris du Lot (*Vitis rupestris*) rootstock cuttings. Food, Agriculture and Environment. 2004; 2(2): 307-309.
- Sahoo PK, Behera LK, Nayak S.** Vegetative propagation of Physic nut (*Jatopha curcas* L.) through stem cuttings. Journal of Applied and Natural Science. 2014; 6 (2):467-472.
- Sardoei AS.** Effect of different media of cuttings on rooting of guava (*Psidium guajava* L.). European Journal of Experimental Biology. 2014; 4(2): 88-92.
- Sarkar PK, Bishnoi SK, Shinde R.** Under-exploited fruits of the forests: Eastern Indian Plateau is a potential hub. *In: Fruits for livelihood: Production technology and management practices*; Dutta, A.K .and Mondal, B. (eds.). Published by Agrobios (India), Jodhpur, India. 2017b; pp. 261-278.
- Sarkar PK, Bishnoi SK, Shinde R, Das B.** Prevalent agroforestry systems of Jharkhand state of India: A livelihood option. Rashtriya Krishi. 2017a; 12(1): 87-89.
- Sengel E, Isci B, Ahmet A.** Effects of different culture media on rooting in grafted grapevine. Ege Üniversitesi Ziraat Fakültesi Derg. 2012; 49(2): 143-148.
- Shah M, Khattak AM, Amin NU.** Effect of different growing media on the rooting of *Ficus binnendijkii* 'Amstel Queen' cuttings. Journal of agricultural and biological science. 2013; 1(3): 15-17.
- Shamet GS, Naveen CR.** Study of rooting in stem cuttings of Khirk (*Celtis australis* Linn.). Indian Journal of Forestry. 2005; 28(4): 363-369.
- Sheoran OP, Tonk DS, Kaushik LS, Hasija RC, Pannu RS.** Statistical software package for agricultural research workers. *In: Recent advances in information theory, statistics & computer applications*; Hooda, D.S. and Hasija, R.C. (eds.). Published by Department of Mathematics Statistics, Chaudhary Charan Singh Haryana Agricultural University (CCS HAU), Hisar. 1998; pp. 139-143.

Smith NG, Wareing PE. The rooting of actively growing and dormant leafy cuttings in relation to endogenous hormone levels and photoperiod. *New Phyto.* 1972; 71: 483-500.

Wright RCM. The complete hand book of plant propagation. Macmillan, New York. 1975.

UNDER PEER REVIEW