

Persistent Organic Pollutants- A Silent Threat to the Agro-ecosystem and Surrounding Environment

ABSTRACT

Persistent organic pollutants (POPs) are a class of hazardous, bio-accumulative, biomagnified, and extremely persistent chemicals with plausible extended off-site mobility in the environment or agro-ecosystem. These constitute a wide variety of chemicals, their sources may be either natural or anthropogenic. These are contaminants of emerging concerns for researchers as well as environmentalists. These contaminants are supposed to cause toxicity in terms of their carcinogenic, genotoxic, neurotoxic, and endocrine disruptors nature. Therefore, it is very important to discuss various kinds of POPs present in the environment and their detection methods. Various extraction and analytical techniques have been strategized to determine the POPs in soil and food samples either quantitatively or qualitatively. The literature lacks a comprehensive review over various POPs present in the environment, different techniques of their extraction and analysis, along with their toxicity in agro-ecosystem. The present review is expected to fulfill this gap by considering all the necessary aspects discussed above.

Keywords: Persistent organic pollutants, Agro-ecosystem, Pollutants, Carcinogenic, extraction techniques

1. Introduction

With the rapid urbanization and industrialization, a plethora of environmental contamination issues have been unveiled and have become increasingly serious around the world [1,2]. A variety of pollutants present in the environment and identified by the scientific community are considered highly toxic to living bodies and environment. Some of these are resistant to environmental degradation (physical, biochemical and biological phenomena) and can exist for a very long time [3]. Persistent organic pollutants (POPs) are the pollutants that are existing in our environment for an extended period of time. POPs include polychlorinated biphenyls (PCBs), polychlorinated dioxins and dibenzofurans (PCDFs- as a single entry), organochlorine insecticides, polycyclic aromatic hydrocarbons (PAHs), and many more. These pollutants have been present for a long time and travel from one location to another. They have also been reported in places where they have never been used, like the earth's polar regions [4].

Many reports are available in the literature which justified that POPs can disrupt the endocrine system and are often called as endocrine disruptors. Different hazardous outcomes have been observed such as population declines, reproductive impairment, embryonic deformities, and metabolic and behavioral changes in various species including eagles, alligators, and cormorants during exposure to POPs. Therefore, there is a big challenge for the scientific community to successfully detect POPs in agro-ecosystem and environmental samples, so that better strategies could be designed to remove them from the environment. Hence, the present review will focus all types of POPs observed till date, their toxicity behaviors, extraction techniques and the methods of their analysis.

2. Persistent organic pollutants

Various pollutants are constantly contaminating the earth's ecosystem and primarily agro-ecosystem. POPs have drawn significant attention over the past several decades. These are mainly featured by their abilities to resist degradation, persist longer and bioaccumulate. These molecules travel great distances, withstand degradation, and bioaccumulate. owing to their unique chemical structures and physicochemical properties. Twelve POPs (commonly called the original 'dirty dozens' or 'legacy POPs') were initially identified during the Stockholm Convention and proposed ban on their usage or source [5]. Later, certain non-chlorinated compounds were also added to the list such as perfluorooctanesulfonate (PFOS), per and poly-fluoroalkyl substances (PFASs), brominated flame retardants (BFRs), and other perfluorinated compounds (PFCs).

2.1. Classification and types of POPs

There are two main categories of POPs on the basis of the source from which they originate. One is intentionally produced and the other one is unintentionally produced (Fig. 1). Considering the main group of unique 12 POPs include 10 intentionally generated pollutants i.e. aldrin, chlordane, DDT, dieldrin, endrin, heptachlor, hexachlorobenzene (HCB), mirex, toxaphene, and polychlorinated biphenyls (PCBs) and 2 unintentionally generated contaminants- polychlorinated dibenzo-p-dioxins (PCDDs) and polychlorinated dibenzofurans (PCDFs) [6].




Fig. 1: Types of POPs with examples**2.1.1. Intentionally generated POPs**


These are mostly chlorinated compounds and have highly lipophilic organic moieties linked with chlorine atoms. These compounds are generally known as organochlorine compounds (OCs) and are produced as a result of different chemical reactions that involve chlorine. For example, dichlorodiphenyltrichloroethane (DDT) and polychlorinated biphenyls (PCBs). They can be classified into two types that are organochlorine pesticides and industrial chemicals [7].









2.1.2. Organochlorine pesticides (OCPs):





OCPs are a class of substances that are commonly employed as pesticides and due to their prolonged persistence, these are also categorized under POPs. They may be insecticides, fungicides, herbicides, rodenticides, nematocides, and acaricides. For example, aldrin, dieldrin, heptachlor, and endrin. They are mostly man-made synthetic organic compounds but some are natural say for example nicotine. Some OCPs along with their chemical structure, usage, toxicity values and WHO recommendations based on the toxicity values are listed in Table 1. These are considered as the most widely used pesticides in developing countries of Asia [8].

Approximately 80% of pesticides used in agriculture are released into the environment by volatilization, runoff, infiltration, transit down the food chain, and other ways. Therefore, the use of OCPs has been prohibited in many countries for a considerable period, but their residues are still causing a significant impact on the environment and the ecosystems. These may be neurotoxic, genotoxic, and carcinogenic in nature. OCPs when applied in the fields, get strongly adsorbed to the soil particles as these are hydrophobic in nature with very high adsorption coefficients. Their residues can be detected on the surface layers of soil after adsorption without draining into the soil profile, allowing them to survive in the soil for months to years [9].

Table 1: Common OCPs along with their chemical structures, usage, half-life, toxicity values, and recommendations by WHO

Sr No	Chemical Name	Structure	Use	Persistence Half-life	Toxicity in rats (mg/kg)	WHO Recommendation on the basis of LD50
1	Aldrin		Insecticide	Moderate 4–7 years	Oral: 39 to 60 Dermal:	Highly hazardous

100						
2	Benzene hexachloride (BHC)		Acaricide Insecticide Rodenticide	High 3 – 6 years	Oral: 10,000	Moderately hazardous
3	Chlordane		Insecticide	High 10 years	Oral: 200-700 Dermal: 530–690	Moderately hazardous
4	DDT		Acaricide Insecticide	High 2–15 years	Oral: 113–130 Dermal: 2510	Moderately hazardous
5	Dieldrin		Insecticide	High 9 months	Oral: 46 Dermal: 50–120	Highly hazardous
6	Endosulfan		Insecticide	Moderate 35-150 days	Oral: 18 to 220	Highly hazardous
7	Endrin		Avicide insecticide	High 1-12 Years	Oral: 3 Dermal: 15	Highly hazardous
8	Heptachlor		Insecticide	High 2 years	Oral: 40– 220 Dermal: 119–320	Highly – Moderately hazardous
9	Isobenzan		Insecticide	High 2.8 years	Oral: 4.8	Highly hazardous

11	Lindane		Rodenticide Acaricide Insecticide	High 15 months	Oral: 88 – 270	Moderately hazardous
12	Methoxychlor		Insecticide	High < 120 Days	Oral: 5000– 6000	Less hazardous
13	Mirex		Insecticide	High 10 years	Oral: 600– 740	Less hazardous
14	Toxaphene (Camphchlor)		Acaricide Insecticide	Moderate 11 Years	Oral: 80– 293	Slightly hazardous

Source: Pesticides properties database (PPDB)

2.1.3. Industrial chemicals:

Thousands of chemicals are being produced in the industries and are used worldwide in large quantities for various preparations such as dyes, resins, plasticizers, antioxidants, surfactants, food preservatives, sanitizers, and detergents. As a result, these harmful compounds are releasing in the environment continuously, especially in wastewater discharged from industries or domestic outlets. Till date, these pollutants have been reported in surface water and wastewater. Some of these have been grouped into the priority pollutants as they cause serious health issues to aquatic organisms. Polychlorinated biphenyls and per and poly-fluoroalkyl substances (PFASs) are the most important types of industrial chemicals found extensively in the agro-ecosystem and causing health hazards.

2.1.4. Polychlorinated biphenyls (PCBs)

PCBs have been discovered in water, sediments, avian tissue, and fish tissue all across the world. These substances are classified as special wastes because they include 2-10 chlorine atoms linked to the biphenyl molecule. This class also includes monochlorinated biphenyls (biphenyl molecules with one chlorine atom added). The basic chemical skeleton of chlorinated biphenyls and some commonly found PCBs are shown in Table 2. PCBs were having numerous industrial applications and were used extensively around 1960-80 as insulating condensers, heat exchangers, plasticizers, flame retardants, fire-resistant transformers, papers, asbestos, etc. Their remains are still present at industrial sites and have

been found in building doors and windows, as well as in paints used in swimming pools and garages [10]. It is believed that over 200 commercial PCB-containing products are now accessible on the market.

Table 2: Some previously reported PCBs along with their chemical structures

PCB 10 ^a	PCB 11 ^b	PCB 24 ^a
PCB 30 ^a	PCB 62 ^a	PCB 65 ^a
PCB 77 ^b	PCB 95 ^b	PCB 116 ^a
PCB 118 ^b	PCB 126 ^b	PCB 153 ^b






a[11], b[12]





2.1.5. Per and poly-fluoroalkyl substances

For more than 50 years, PFASs have been produced and are widely present in the environment. These are non-stick, non-wetting, non-toxic, fire-resistant, and heat resistant. As a result, they have been used in nearly every aspect of labor, from cookware to firefighting foams, paper coatings, and textiles, and so on [13]. These are organofluoride compounds made up of oligomers and polymers in which the hydrogen atoms in the hydrocarbon skeleton have been replaced by fluorine atoms. Per fluorinated sulfonic acids (PFSAs), perfluoro carboxylic acids (PFCAs), high-molecular-weight fluoropolymers, and low-molecular-weight perfluoro alkanolamines are the four categories. Out of these, PFSAs and PFOA, have received the most attention. The most commonly discovered PFASs are mentioned in Table 3. They are

both hydrophobic and lipophobic in nature, and they contain a CF bond, which is one of the strongest chemical connections and provides these molecules with extremely high chemical and thermal stability. Due to these reasons, they are resistant to the typical environmental degradation processes like photolysis, hydrolysis, and atmospheric photooxidation, thereby persisting and bioaccumulating in the environment for years [14]. The National Toxicology Program has also conducted research on PFOA, other PFCAs and PFSA's in order to better understand their toxicity and persistence in human blood [15]. PFOA-contaminated water has been shown to have a negative impact on mammary gland development in mice [16]. Another study found that PFOS exposure can influence the nervous and endocrine system in rats; however, their mechanisms are yet to be established [17].

Table 3: Different types of well-known PFASs, their chemical structures, and sources.

Sr No.	Molecular Structure	Sources	References
1	 <p>Perfluorobutanoic Acid (PFBA)</p>	Stain repellants, food packaging carpets	[18]
2	 <p>Perfluorobutane sulfonic Acid (PFBS)</p>	Fire retardant foams, metal coating, mist suppressant	[19]
3	 <p>Perfluorooctanoic acid (PFOA)</p>	Fire retardant foams, ceramic glaze	[20]
4	 <p>Perfluorooctanesulfonic acid (PFOS)</p>	Water purification, non- stick utensils, textile industry	[21]
5		Cleansing products, metal coating, textile	[22]

	6:2 Fluorotelomer sulfonamide alkyl betaine (FTAB)	industries	
6	 6:2 Fluorotelomer sulfonate (FTSA)	Chemically resilient components, rug coverings	[23]
7	 Chlorinated polyfluorinated ether sulfonate (F53-B)	General waterproofing lubricants, gaskets power-utility stoppers	[24]
8	 Viton A/B	Fire retardant foams, metal coating, mist suppressant	[25], [26]
9	 Polytetrafluoroethylene (PTFE)	Fire retardant foams, metal coating	[27]

2.2. Unintentionally produced POPs

These are mostly undesired by-products of combustion or other chemical reactions that take place in the presence of chlorine or chlorinated substances. Polycyclic aromatic hydrocarbons (PAHs), dioxins, and furans are the three primary categories.

2.2.1. Polycyclic aromatic hydrocarbons (PAHs)

Both natural and anthropogenic sources release PAHs into the environment [28]. The phenomena including incomplete burning of fuels, garbage, some organic substances such as tobacco and plant material, forest

Table 4: Pyrogenic and petrogenic PAHs reported in the literature

Pyrogenic PAHs			
Naphthalene	Acenaphthylene	Acenaphthene	Fluorene
Anthracene	Phenanthrene	Fluoranthene	Pyrene
Chrysene	Benz[a]anthracene	Benzo[b]fluoranthene	Benzo[k]fluoranthene
Benzo[a]pyrene	Indeno[1,2,3-cd]pyrene	Benzo[g,h,i]perylene	Dibenz[a,h]anthracene
Pyrogenic PAHs			

1-Methylphenanthrene	9-Ethylphenanthrene	1-Methyl-7-isopropyl phenanthrene	5-Methylchrysene
6-Ethylchrysene	Acenaphthenequinone	1,4-Anthraquinone	9,10- Phenanthrenequinone

fires, and volcanic eruptions lead to the exposure of PAHs to the environment. This is a large class of chemical compounds that include two or more fused aromatic rings, ranging from two-ring naphthalene and its derivatives to complex structures with up to 10 rings. PAHs with up to six rings are known as 'small' PAHs, while those having more than six aromatic rings are called 'large' PAHs [29]. Some of the PAHs are posing a serious threat to human health as these can be carcinogens, mutagens, and teratogens as well. Because of their physicochemical properties, PAHs are highly mobile in the environment, allowing them to spread through air, soil, and water bodies, where their presence is ubiquitous. Some of the commonly found PAHs and their chemical structures are shown in Table 4.

The US Environmental Protection Agency (US EPA) has designated 16 PAHs as priority pollutants. Benzo[a]pyrene is a well-known human carcinogen among these key PAHs and is frequently employed as an indication of PAH exposure [30]. Furthermore, these are categorized based on their origins as pyrogenic PAHs, which are produced as a result of fossil fuel combustion, and petrogenic PAHs, which are peculiar to crude oil and pollute water following an oil spill [31]. Petrogenic PAHs differ from pyrogenic PAHs in that they are either heavily alkylated or oxygenated to produce PAH quinones.

2.2.2. Dioxins and dibenzofurans

Chlorinated dibenzofurans (furans) and chlorinated dibenzo-p-dioxins (dioxins) have similar chemical properties and toxic effects, thereby being a major concern for decades. 2,3,7,8-TCDD; 2,3,7,8-TCDF and PCB-126 are examples of this kind of substance. Nowadays, there has been a new group called dioxin-like PCBs (DLPCBs) which includes a specific subgroup of PCBs. Polychlorinated dioxins (PCDDs) and dibenzofurans (PCDFs) are released from anthropogenic sources and activities rather than natural processes [32].

3. Extraction methods of POPs in the environmental samples

Extraction of POPs from solid environmental samples has frequently been done using various organic solvents with or without addition of heat. The basic extraction techniques are liquid/liquid extraction and soxhlet extraction. A number of extraction techniques along with the analytical methods used for various

POPs have been summarized in Table 5. Different methods have different recoveries depending on the type of analyte and the sample type. For instance, some reports have suggested that sediments and soils be freeze-dried [33]; while others have reported lower recoveries of PCBs from freeze-dried sediments [34].

Soxhlet extraction involves extractors in which a high amount of sample is mixed with the organic solvents and extracts the component without filtering. The major limitations of this method were large amounts of solvents (about 500 mL), and high extraction time (up to 2 days). Hence, some amendments have been made to the traditional soxhlet extraction like automated, focused microwave-assisted, and high-pressure soxhlet extraction. For example, the analysis of PCBs in fish and sediment samples achieved the best recoveries [35,36]. From sediment samples of the baltic sea, PCBs were extracted using accelerated solvent extraction (ASE) with solvent and toluene for soxhlet extraction [37].

Liquid-liquid extraction (LLE) was found to be very tedious and labor intensive, hence solid-phase extraction (SPE) was its possible replacement. It offers less solvent exposure, quick processing, and easy handling as compared to the LLE [38,39]. Solid-phase extraction was used for the extraction of POPs from water and wastewater samples, even with high loadings of analytes which resulted in the reduction of the solvent usage and the analysis time [40]. OCPs and PCBs were successfully extracted from human serum samples with this method by using n-hexane-DCM mixture with recoveries of 99 to 120% for PCBs and 88 to 115% for OCPs [41]. Some samples require solid phase microextraction (SPME) techniques and headspace sampling.

Supercritical fluid extraction (SFE) is another efficient method for removing PCBs from the soil, sediments, and other solid samples. The primary determinants of recovery rates in this approach are optimizing temperature, pressure, and flow rate. For example, PCBs were removed from sediment samples using this process at 200°C and 150-650 atm pressure with a 95% recovery rate [42]. PCBs were extracted from fatty acid samples using this approach with a basic alumina combination as an alternative to silica/silver nitrate (1:10, w/w) [43]. But poor repeatability is the major drawback of this method which has not been improved yet.

Passive sampling is also an effective method for assessing water samples utilizing semi-permeable membrane devices (SPMDs). SPMD sampling is a type of passive sampling for water, air, or sediments. Some other passive sampling techniques have been recently reviewed in the literature which is mainly based on the free flow of analytes from the sampling medium to the collecting medium [44]. Furthermore, some samples necessitate cleanup processes, which include the removal of co-extracted lipids from such samples prior to analysis. This will be determined primarily by the type of analyte. Because PCBs are highly stable in acidic conditions, it will be an effective strategy to conduct out PCB cleanup processes employing sulfuric acid or acid-impregnated silica columns for the elimination of lipids. Following that, sulfur must be removed from the samples in order to reduce co-extractive

interferences and lengthen the detector's lifetime. Sulfur can thus be eliminated by introducing copper granules during the extraction process [45].

4. Analytical techniques for POPs

Different types of chromatographic techniques have been employed for POPs analysis, such as gas chromatography (GC), ultra-high-performance liquid chromatography/or ultraperformance liquid chromatography (UHPLC/UPLC), high-performance liquid chromatography (HPLC), liquid chromatography (LC), and capillary liquid chromatography (CLC). A number of analytical methods used for the detection and quantification of various POPs have been summarized in this study.

The traditional HPLC is regularly used for the analysis of ionic PFAS, specially PFCA and PFSA, whereas GC is used for the analysis of volatile and semi-volatile PFAS. Particles collected on C18 extraction disks can be extracted quantitatively without Soxhlet or PLE extraction. PFAS have been extracted from sediments collected from different lakes by sonicating them in methanol (recoveries: 88-102%) and analyzed by UPLC/ESI-MS/MS and obtained the total concentrations ranging from 0.61 ppb to 26 ppb [46]. LC-MS/MS method was also developed for the analysis of hydroxylated PCBs with a LOD value of 0.1 ng/mL after applying suitable cleanup to them [47,48,49].

One of the most important analytical methods for the analysis of POPs is gas chromatography (GC). Pressure or temperature-programmed injection techniques, combined with increased injection volume, have been proven to be quite effective without having any detrimental effects (Tranchida and Mondello, 2012). Detection of PCBs and PBDEs in serum samples of livestock in the US, the higher concentration of PCBs and PBDEs were detected in the livestock's serum, but their origin is still unclear [50]. GC-MS-based methods have been used to confirm the presence of PCBs in the transformer oils and detected in air samples too [51]. Several GC-ECD methods have also been reported for the analysis of PCBs from sediments, serum, fish tissue, and indoor air samples in addition to the GC-MS methods [52].

Table 5: Determination of POPs in different sample matrices using various extraction methods and analytical techniques

Sr No	Analytes	Matrix	Extraction Method	Analytical Techniques	LOD values	References
1	PCBs	Urine	SPE	LC/MS/MS	0.01 ng/mL	[49]
2	PCBs	Plasma	SPE	LC/MS/MS	0.01 ng/mL	[48]

3	EPPs, PCBs	Sediment	LLE	LC/MS/MS	0.01-0.4 ng/g	[53]
4	PCBs	Water	LLE	RP-HPLC	0.25 mg/mL	[54]
5	PCBs	Mammals	LLE	RP-HPLC	2.5 mg/kg	[55]
6	PCBs, PCDFs	Fish	LLE	RP-HPLC	2-100 ng/g	[56]
7	PCBs	Herring oil	LLE	RP-HPLC	9-50 ng/g	[57]
9	PFBS, PFOS	Ground, river, and tap water	SPE	LC/(-)ESI- MS/MS	25,000 ng/L (LOQ)	[58]
10	PFOA, PFOS	Surface water	SPE	LC/(-)ESI-MS	0.2-13 ng/L (LOQ)	[59]
11	PFOS, PFOA, PFNA, PFDA,	Rain water	SPE	LC/(-)ESI-MS	0.04-7.2 ng/L	[60]
12	HCB, HCH, PCBs, OCPs	Fish	SE, LLE	GC/ECD, HRGC/ECD	-	[61]
13	PBDEs, PCBs, OCPs	Dolphin	GPC	GC/MS	-	[62]
14	OCPs, PCBs, toxaphene	Bird	LLE, SE	GC/ECD, GC- ECNI-MS	-	[63],[64]
15	PCDD/F, HCB	Air	Swipe/biofilm s, passive sample	HRGC/HRMS	-	[65]
16	PCBs, HCH, OCPs, HCB, DDT	Human	LLE, SPE, SE, LSE, hot SE	SPE-HPLC	-	[66]

5. Toxicity behavior of POPs

The ecotoxicological effects of POPs on the environment, biota, and human health have raised a big concern in the past few years which has led to their minimal usage or complete ban in many countries. These are resistant to almost all biodegradation processes, hence are highly persistent and toxic. It has been recently addressed that in children, POPs can cause cancer and tumours at many places, immune system illnesses, reproductive issues, decreased ability to fight disease, slowed growth, and irreversible cognitive function damage. As a result, POPs are a suspected carcinogen [67].

Considering the contamination of the aquatic environment, PCBs and OCPs have emerged as the most toxic among the wide range of POPs. Moderate to high exposure to PCBs (0.25-2 mg/L) has induced abnormalities in developing zebrafish retinas, resulting in visual impairment [68]. PCB 126 exposure has also resulted in heart malformations in the embryos of *Danio rerio* [69]. Many species of *Daphnia* are used as models or indicator organisms for the toxicity determination of PCBs mainly in freshwater [70]. PFASs have also been found to be toxic to the soil, wastewater, animals, and humans [71,72,73,74].

Certain PAHs, when exposed to animals through food for a prolonged time were found to cause stomach cancer from ingestion, lung cancer from inhalation, and skin cancer from skin contact. For instance, benzo[a]pyrene, benzo[a]anthracene, and chrysene were responsible for significant chromosome aberrations in rodents [75]. Among PAHs, the most common PAH to cause cancer in animals and was the first chemical carcinogen to be discovered is Benzo(a)pyrene [76].

6. Conclusions and way forward

The present compilation of various POPs and their toxicities concludes that new candidate POPs are adding to the list day by day. The 'dirty dozens' are no more dozen in number. Thus, novel research must be focused in developing new analytical approaches for trace estimation of POPs. Moreover, their highly toxic nature clearly indicates that they should be excluded from the environment as soon as possible. Various removal strategies have been proposed for this purpose such as photocatalysis, bio nanocomposites, MOF-based adsorbents or other porous absorbents etc. More work should be done for the development of environmentally benign moieties for the adsorptive removal of these POPs for the well-being of the ecosystem. Future research must also focus on estimation of toxicity/health hazards of POPs to various non-target organisms and associated environment or agro-ecosystem.

References

1. Fang H, Zhang H, Han L, Mei J, Ge Q, Long Z, Yu Y. Exploring bacterial communities and biodegradation genes in activated sludge from pesticide wastewater treatment plants via metagenomic analysis. *Environment Pollution*. 2018;243:1206–1216.

2. Fu Q, Fedrizzi D, Kosfeld V, Schlechtriem C, Ganz V, Derrer S, Rentsch D, Hollender J. Biotransformation changes bioaccumulation and toxicity of diclofenac in aquatic organisms. *Environment Science and Technology*. 2020;54:4400-4408.
3. Alharbi OM, Khattab RA, Ali I. Health and environmental effects of persistent organic pollutants. *Journal of Molecular Liquids*. 2018;263:442-453.
4. Jacob J, Cherian J. Review of environmental and human exposure to persistent organic pollutants. *Asian Social Science*. 2013;9:107-120.
5. Annex C. The Stockholm Convention on Persistent Organic Pollutants. United Nations Environmental Programme, Geneva, 2008;29.
6. Ashraf MA. Persistent organic pollutants (POPs): a global issue, a global challenge. *Environmental Science and Pollution Research*. 2017;24(5):4223-4227.
7. Harner T, Pozo K, Gouin T, Macdonald AM, Hung H, Cainey J, Peters A. Global pilot study for persistent organic pollutants (POPs) using PUF disk passive air samplers. *Environmental Pollution*. 2006;144(2):445-452.
8. Gupta PK. Pesticide exposure-Indian scene. *Toxicology*. 2004;198:83-90.
9. Joseph L, Paulose SV, Cyril N, Santhosh SK, Varghese A, Nelson AB, Kasu S. Organochlorine pesticides in the soils of Cardamom Hill Reserve (CHR), Kerala, India: Geospatial distribution, ecological and human health risk assessment. *Environmental Chemistry and Ecotoxicology*. 2020;2:1-11.
10. Koh IO, Rotard W, Thiemann WHP. Analysis of chlorinated paraffins in cutting fluids and sealing materials by carbon skeleton reaction gas chromatography. *Chemosphere*. 2002;47(2):219-227.
11. Jing R, Fusi S, Kjellerup BV. Remediation of polychlorinated biphenyls (PCBs) in contaminated soils and sediment: state of knowledge and perspectives. *Frontiers in Environmental Science*. 2018;6:79.
12. Klocke C, Sethi S, Lein PJ. The developmental neurotoxicity of legacy vs. contemporary polychlorinated biphenyls (PCBs): similarities and differences. *Environmental Science and Pollution Research*. 2020;27(9):8885-8896.
13. Buck RC, Franklin J, Berger U, Conder JM, Cousins IT, Voogt PD, Jensen AA, Kannan K, Mabury SA, Leeuwen SPJ. Perfluoroalkyl and polyfluoroalkyl substances in the environment: terminology, classification, and origins. *Integrated Environmental Assessment and Management*. 2011;7 (4): 513-541.
14. Schultz MM, Barofsky DF, Field JA. Fluorinated alkyl surfactants. *Environmental engineering science*. 2004;20 (5): 487-501.
15. National Institute of Environmental Health Sciences, 2016. Perfluorinated Chemicals (PFCs). Technical Representative., NIH.

16. Post GB, Cohn PD, Cooper KR. Perfluorooctanoic acid (PFOA), an emerging drinking water contaminant: a critical review of recent literature. *Environmental Research*. 2014;116:93–117.
17. Austin ME, Kasturi BS, Barber M, Kannan K, Mohan Kumar PS, Mohan Kumar SMJ. Neuroendocrine effects of perfluorooctane sulfonate in rats. *Environmental Health Perspectives*. 2003;111(12):1485–1489.
18. Abraham K, El-Khatib AH, Schwerdtle T, Monien BH. Perfluorobutanoic acid (PFBA): No high-level accumulation in human lung and kidney tissue. *International Journal of Hygiene and Environmental Health*. 2021;237:113830.
19. Qi W, Clark JM, Timme-Laragy AR, Park Y. Perfluorobutanesulfonic acid (PFBS) induces fat accumulation in HepG2 human hepatoma. *Toxicological and Environmental Chemistry*. 2020;102(10):585-606.
20. D'Agostino LA, Mabury SA. Identification of Novel Fluorinated Surfactants in Aqueous Film Forming Foams and Commercial Surfactant Concentrates. *Environmental Science and Technology*. 2014;48:121–129.
21. Du Z, Deng S, Liu D, Yao X, Wang Y, Lu X, Wang B, Huang J, Wang Y, Xing B, Yu G. Efficient adsorption of PFOS and F53B from chrome plating wastewater and their subsequent degradation in the regeneration process. *Chemical Engineering Journal*. 2016;290:405-413.
22. Steenland K, Fletcher T, Savitz DA. Epidemiologic Evidence on the Health Effects of Perfluorooctanoic Acid (PFOA). *Environmental Health Perspectives*. 2010;118:1100–1108.
23. Henry BJ, Carlin JP, Hammerschmidt JA, Buck RC, Buxton LW, Fiedler H, Seed J, Hernandez O. A critical review of the application of polymer of low concern and regulatory criteria to fluoropolymers. *Integrated Environmental Assessment and Management*. 2018;14:316–334.
24. Li F, Duan J, Tian S, Ji H, Zhu Y, Wei Z, Zhao D. Short-chain per- and polyfluoroalkyl substances in aquatic systems: Occurrence, impacts and treatment. *Chemical Engineering Journal*. 2020;380:122506.
25. Shaw DMJ, Munoz G, Bottos EM, Duy SV, Sauvé S, Liu J, Van Hamme JD. Degradation and defluorination of 6:2 fluorotelomer sulfonamidoalkyl betaine and 6:2 fluorotelomer sulfonate by *Gordonia* sp. strain NB4-1Y under sulfur-limiting conditions. *Science of the Total Environment*. 2019;647:690–698.
26. Dixit F, Barbeau B, Mostafavi SG, Mohseni M. PFOA and PFOS removal by ion exchange for water reuse and drinking applications: Role of organic matter characteristics. *Environment Science and Water Research Technology*. 2019;5:1782–1795.
27. Gardiner J. Fluoropolymers: Origin, Production, and Industrial and Commercial Applications. *Australian Journal of Chemistry*. 2015;68:13.

28. WHO (World Health Organization), 2003. Polynuclear aromatic hydrocarbons in drinking water. Background document for development of WHO Guidelines for Drinking-water Quality.
29. Arey J, Atkinson R. Photochemical reactions of PAH in the atmosphere. In: Douben PET, editor. PAHs: An ecotoxicological perspective. New York: John Wiley and Sons Ltd. 2003;47–63
30. Huang M, Penning TM. Processing contaminants: polycyclic aromatic hydrocarbons (PAHs). *Encyclopedia of food safety* 2014;2:416-423.
31. Zielinska B, Samy S. Analysis of nitrated polycyclic aromatic hydrocarbons. *Analytical and Bioanalytical Chemistry*. 2006;386:883–890.
32. El-Shahawi MS, Hamza A, Bashammakh, AS, Al-Saggaf WT. An overview on the accumulation, distribution, transformations, toxicity and analytical methods for the monitoring of persistent organic pollutants. *Talanta*. 2010;80(5):1587-1597.
33. Wells DE, Hess P. Separation, clean-up and recoveries of persistent trace organic contaminants from soils, sediment and biological matrices. In: Barcelo E (ed) *Sample handling and trace analysis of pollutants, techniques, applications and quality assurance*, Elsevier, Amsterdam. 2000:73-116.
34. Gevao B, Hamilton Taylor J, Jones KC. Towards a complete mass balance and model for PCBs and PAHs in a small rural lake, Cumbria UK. *Limnology and oceanography*. 2000;45(4):881-894.
35. Hess JE, Caudill CC, Keefer ML, McIlraith BJ, Moser ML, Narum SR. Genes predict long distance migration and large body size in a migratory fish, Pacific lamprey. *Evolutionary applications*. 2014;7(10):1192-1208.
36. Cui Y, Wang Z, Cong J, Wang L, Liu Y, Wang X, Xie J.. Determination of polychlorinated biphenyls in fish tissues from shanghai seafood markets using a modified QuEChERS method. *Analytical Sciences*. 2017;33(8):973-977.
37. Bandh C, Bjorklund E, Mathiasson L, Naf C, Zebuhr Y. Comparison of accelerated solvent extraction and Soxhlet extraction for the determination of PCBs in Baltic Sea sediments. *Environmental science & technology*, 2000;34(23): 4995-5000.
38. Hartmann R. Extraction of PCDD/Fs and PCBs in fish using speed extractor E-914, *Procedia environmental sciences*. 2013;18, 875-881.
39. Zhao Y, Li W, Liu J, Huang K, Wu C, Shao H, Chen H, Liu X. Modification of garlic peel by nitric acid and its application as a novel adsorbent for solid-phase extraction of quinolone antibiotics. *Chemical Engineering Journal*. 2017;.326:745-755.
40. Kurt SK, Gursel IV, Hessel V, Nigam KD, Kockmann N. Liquid–liquid extraction system with microstructured coiled flow inverter and other capillary setups for single-stage extraction applications. *Chemical Engineering Journal*. 2016;284:764-777.

41. Conka K, Drobna B, Koca, A, Petrik J. Simple solid-phase extraction method for determination of polychlorinated biphenyls and selected organochlorine pesticides in human serum. *Journal of Chromatography A*. 2005;1084(1-2):33-38.
42. Langenfeld JJ, Burford MD, Hawthorne SB, Miller DJ. Effects of collection solvent parameters and extraction cell geometry on supercritical fluid extraction efficiencies. *Journal of Chromatography A*. 1992;594(1-2):97-307.
43. Reddy AVB, Moniruzzaman M, Aminabhavi TM. Polychlorinated biphenyls (PCBs) in the environment: Recent updates on sampling, pretreatment, cleanup technologies and their analysis. *Chemical Engineering Journal*. 2019;358:1186-1207.
44. Namiesnik J, Zabiegala B, Kot-Wasik A, Partyka M, Wasik A. Passive sampling and/or extraction techniques in environmental analysis: a review. *Analytical and bioanalytical chemistry*. 2005;381(2):279-301.
45. Reddy, A.V.B., Moniruzzaman, M., Aminabhavi, T.M., 2019. Polychlorinated biphenyls (PCBs) in the environment: Recent updates on sampling, pretreatment, cleanup technologies and their analysis. *Chemical Engineering Journal*. 358, 1186-1207.
46. Li F, Huang H, Xu Z, Ni H, Yan H, Chen R, Luo Y, Pan W, Long J, Ye X, Qian X. Investigation of perfluoroalkyl substances (PFASs) in sediments from the urban lakes of anqing city, anhui Province, China. *Bulletin of environmental contamination and toxicology*. 2017;99(6):760-764.
47. Liang R, Zhao Y, Su Y, Qi, W. Determination of hydroxylated polychlorinated biphenyls by offline solid-phase extraction-liquid chromatography-tandem mass spectrometry using a molecularly imprinted polymer as a sorbent for sample preconcentration. *Talanta*. 2015;144:115-121.
48. Quinete N, Kraus T, Belov VN, Aretz C, Esser A, Schettgen, T. Fast determination of hydroxylated polychlorinated biphenyls in human plasma by online solid phase extraction coupled to liquid chromatography-tandem mass spectrometry. *Analytica chimica acta*. 2015;888, 94-102.
49. Quinete N, Esser A, Kraus T, Schettgen T. Determination of hydroxylated polychlorinated biphenyls (OH-PCBs) in human urine in a highly occupationally exposed German cohort: New prospects for urinary biomarkers of PCB exposure. *Environment International*. 2016;97:171-179.
50. Sethi S, Chen X, Kass PH, Puschner B. Polychlorinated biphenyl and polybrominated diphenyl ether profiles in serum from cattle, sheep, and goats across California. *Chemosphere*. 2017;181:63-73.
51. Francisco AP, Nardocci AC, Tominaga MY, da Silva CR, de Assuncao JV. Spatial and seasonal trends of polychlorinated dioxins, furans and dioxin-like polychlorinated biphenyls in air using passive and active samplers and inhalation risk assessment. *Atmospheric Pollution Research*. 2017;8(5):979-987.

52. Qiu C, Cochran J, Smuts J, Walsh P, Schug KA. Gas chromatography-vacuum ultraviolet detection for classification and speciation of polychlorinated biphenyls in industrial mixtures. *Journal of Chromatography A*. 2017;1490:191-200.
53. Martin J, Zafra-Gomez A, Hidalgo F, Ibanez-Yuste AJ, Alonso E, Vilchez JL. Multi-residue analysis of 36 priority and emerging pollutants in marine echinoderms (*Holothuria tubulosa*) and marine sediments by solid-liquid extraction followed by dispersive solid phase extraction and liquid chromatography–tandem mass spectrometry analysis. *Talanta*. 2017;166:336-348.
54. Shuaia J, Yua X, Zhanga J, Xiongb A, Xionga F. Regional analysis of potential polychlorinated biphenyl degrading bacterial strains from China, Braz. *Journal of Microbiology*. 2016;47:536-541.
55. Hayteas DL, Duffield DA. 1997. The determination by HPLC of PCB and p,p'-DDE residues in marine mammals stranded on the Oregon Coast 1991-1995. *Marine Pollution Bulletin*. 1997;34:844-848.
56. Zebtihr Y, Ntif C, Bandh C, Broman D, Ishaq R, Pettersen, H. An automated HPLC separation method with two coupled columns for the analysis of PCDD/Fs, PCBs and PACs. *Chemosphere*. 1993;27:1211-1219.
57. Lundgren K, Bavel B, Tysklind M. Development of a high-performance liquid chromatography carbon column based method for the fractionation of dioxin-like polychlorinated biphenyls, *Journal of Chromatography A*. 2002;962:79-93
58. Hebert GN, Odom MA, Craig PS, Dick DL, Strauss SH. Method for the determination of sub-ppm concentrations of perfluoroalkylsulfonate anions in water. *Journal of Environmental Monitoring*. 2002;4(1):90-95.
59. Boulanger B, Vargo J, Schnoor JL, Hornbuckle KC. Detection of perfluorooctane surfactants in Great Lakes water. *Environmental Science & Technology*. 2004;38(15):4064-4070.
60. Loewen M, Halldorson T, Wang F, Tomy G. Fluorotelomer carboxylic acids and PFOS in rainwater from an urban center in Canada. *Environmental science & technology*. 2005;39(9):2944-2951.
61. Fernandez P, Carrera G, Grimalt JO. Persistent organic pollutants in remote freshwater ecosystems. *Aquatic Sciences*. 2005;67(3):263-273.
62. Menone ML, Aizpun de Moreno JE, Moreno VJ, Lanfranchi AL, Metcalfe TL, Metcalfe CD. Organochlorine pesticides and PCBs in a southern Atlantic coastal lagoon watershed, Argentina. *Archives of Environmental Contamination and Toxicology*, 40(3), 355-362.
63. Barakat AO, 2004. Assessment of persistent toxic substances in the environment of Egypt. *Environment International*. 2004;30(3):309-322.

64. Gao J, Zhou H, Pan G, Wang J, Chen B. Factors influencing the persistence of organochlorine pesticides in surface soil from the region around the Hongze Lake, China. *Science of the Total Environment*. 2013;443:7-13.
65. Sanz-Landaluze J, Bartolome L, Zuloaga O, Gonzalez L, Dietz C, Camara C. Accelerated extraction for determination of polycyclic aromatic hydrocarbons in marine biota. *Analytical and bioanalytical chemistry*. 2006;384(6):1331-1340.
66. Pang GF, Liu YM, Fan CL, Zhang JJ, Cao YZ, Li XM, Li ZY, Wu YP, Guo TT. Simultaneous determination of 405 pesticide residues in grain by accelerated solvent extraction then gas chromatography-mass spectrometry or liquid chromatography-tandem mass spectrometry. *Analytical and Bioanalytical Chemistry*. 2006;384(6):1366-1408.
67. Bolt HM, Degen GH. Comparative assessment of endocrine modulators with oestrogenic activity. II. Persistent organochlorine pollutants. *Archives of toxicology*. 2002;76(4):187-193.
68. Wang YP, Hong Q, Qin DN, Kou CZ, Zhang CM, Guo M, Guo XR, Chi X, Tong ML. Effects of embryonic exposure to polychlorinated biphenyls on zebrafish (*Danio rerio*) retinal development. *Journal of Applied Toxicology*. 2012;32(3):186-193.
69. Grimes AC, Erwin KN, Stadt HA, Hunter GL, Gefroh HA, Tsai HJ, Kirby ML. PCB126 exposure disrupts zebrafish ventricular and branchial but not early neural crest development. *Toxicological sciences*. 2008;106(1):193-205.
70. Harmon SM. The toxicity of persistent organic pollutants to aquatic organisms. In *Comprehensive Analytical Chemistry*, Elsevier. 2015;67:587-613
71. Rotander A, Toms LML, Aylward L, Kay M, Mueller JF. Elevated levels of PFOS and PFHxS in firefighters exposed to aqueous film forming foam (AFFF). *Environment International*. 2015;82:28-34.
72. Houtz EF, Sutton R, Park JS, Sedlak M. Poly-and perfluoroalkyl substances in wastewater: Significance of unknown precursors, manufacturing shifts, and likely AFFF impacts. *Water research*. 2016;95:142-149.
73. Braunig J, Baduel C, Heffernan A, Rotander A, Donaldson E, Mueller JF. Fate and redistribution of perfluoroalkyl acids through AFFF-impacted groundwater. *Science of the Total Environment*. 2017;596:360-368.
74. Hoiaeter A, Pfaff A, Breedveld, GD. Leaching and transport of PFAS from aqueous film-forming foam (AFFF) in the unsaturated soil at a firefighting training facility under cold climatic conditions. *Journal of contaminant hydrology*. 2019;222:112-122.

75. Jung KH, Kim, JK, Noh JH, Eun JW, Bae HJ, Kim MG, Chang YG, Shen Q, Kim SJ, Kwon SH, Park WS. Characteristic molecular signature for the early detection and prediction of polycyclic aromatic hydrocarbons in rat liver. *Toxicology letters*. 2013;216(1):1-8.
76. Patel AB, Shaikh S, Jain K, Desa, C, Madamwar D. Polycyclic aromatic hydrocarbons: sources, toxicity, and remediation approaches. *Frontiers in Microbiology*. 2021;11:562813.

UNDER PEER REVIEW