

Original Research Article

Macronutrient Status and Soil Chemical Properties as Influenced by Methods and Schedule of Fertilizer Application Under Widely Spaced Sugarcane Crop (*Saccharum officinarum* L.) in Alfisol of Karnal District, Haryana

Comment [K1]: Need to be revised and shorter

1. Abstract:

An experiment was conducted at the Regional Research Station, Uchani (Karnal) of CCS Haryana Agricultural University, Hisar, India to assess the effect of the method and schedule of fertilizer application on macronutrient status viz. available Nitrogen (N), Phosphorus (P) and potassium (K) and soil chemical properties i.e. soil pH, electrical conductivity (EC) and soil organic carbon (SOC) at different intervals of the crop growth period. The experiment was laid out in split plot design with two methods of fertilizer application (B1-broadcasting and B2- band placement) as main plot treatments and four subplot treatments consisting of the application of a recommended doses of N and K fertilizer (RDF) in the different numbers of splits at different no. of days after planting (DAP) i.e. T1 (five splits), T2 (six splits), T3 (seven splits) and T4 (three splits). Results elucidated that the availability of all the macronutrients was significantly affected by both sub and main-plot treatments at different stages of crop growth period. Available N content was found to be significantly higher in B2 compared to B1 at all the time intervals from 75 DAP (131.86 Kg ha⁻¹) to harvest (145.91 Kg ha⁻¹) reaching the maximum at 180 DAP (142.80 Kg ha⁻¹). Among the subplot treatments, T4 and T1 respectively reported significantly higher values of available N from 75 DAP (134.51 Kg ha⁻¹) to 90 DAP (147.08 Kg ha⁻¹) and 90 DAP (153.10 Kg ha⁻¹) to 150 DAP (162.11 Kg ha⁻¹). At 180 DAP and at the time of harvesting, significantly higher values of available N (159.21 Kg ha⁻¹ and 157.01 Kg ha⁻¹) were observed in treatments receiving fertilizer in T2 and T3 splits respectively. Available K content was significantly higher in B1 compared to B2 at all the time intervals except at 0, 150, and 180 DAP and at harvest where the difference was not significant. Higher values of available K were reported under T4 upto 75 DAP (225.30 Kg ha⁻¹) and thereafter, T1 exhibited significantly higher available K values upto 150 DAP (222.06 Kg ha⁻¹). At 180 DAP and at the time of harvesting, significantly higher values of available K (225.10 Kg ha⁻¹ and 227.48 Kg ha⁻¹) were observed in T2 and T3 respectively. Available P content was significantly higher (24.63 Kg ha⁻¹) in B1 compared to B2 at the time of harvesting. Soil pH, EC and SOC did not register any significant change under any treatment.

Comment [K2]: Please mention the conclusion and best treatment

Key words: Split application, Band Placement, Broadcasting, Fertilizer application, Nitrogen, Potassium

2. Introduction:

Chemical fertilizers are the most crucial inputs to satisfy the nutrient requirements of crops for obtaining high yield in modern crop production. Modern agriculture is extensively dependent upon fertilizers and they are regarded as vital tools for global food safety (Pahalviet *et al.*, 2021). ~~The price~~ ~~Price~~ of fertilizers which are already expensive keep on increasing due to a gap in demand and supply of fertilizer minerals (Randiveet *et al.*, 2021) which directly adds up a significant portion to the cost of cultivation. Therefore, a reduction in the cost of cultivation demands the adoption of effective nutrient management techniques. The employment of fertilizers also influences ~~soil's soil~~ physicochemical and biological properties (Singh *et al.*, 2019). The imbalanced use or inefficient management of chemical fertilizers may reduce soil fertility and can cause soil, water and air pollution along with a reduction in important nutrients of soil and minerals when applied without following standard agronomic practices (Katyal and Chaudhri, 2021). To avoid these harmful effects, innovative approaches like the 4R nutrient stewardship approach may serve as a guide for ~~the~~ effective nutrient management. The concept of this approach emphasizes the application of the right source of nutrients at the right rate and time and in the right place (Fixen, 2020).

The adoption of the proper method of fertilizer application is essential to minimize the loss of nutrients from the soil with a simultaneous rise in its availability and it is equally important at the time of application to achieve higher nutrient use efficiency. Method and time of fertilizer application are important components of an effective nutrient management program ~~me~~. Band placement of nitrogenous fertilizer in the subsurface portion of soil prevents its loss through volatilization. Nitrogen is mobile in soil whereas fixation of P and K occurs in the soil suggesting their placement in bands along the crop rows to make them easily available (Shukla *et al.*, 2018). Band Placement of potassium is recommended in soils having a low level of K or with a high K fixing capacity. In upland areas, several studies indicates ~~s~~ band placement is overwhelmingly superior ~~than to~~ broadcasting as long as efficiency is concerned. Proper placement carries the same gravity in P management under wide row spacing as split application carries for N management. P and K applied through fertilizers tend to fix into the soil, therefore practicing band placement aids in reducing fixation and increases their availability (Farmahaet *et al.*, 2013 and Fernández and White, 2012) Band placement of fertilizer near the root zone aids the roots to easily draw nutrients from the soil as it minimizes ~~s~~ the distance between root hairs and nutrients placed in soil. Less contact with the soil lessens the opportunity for nutrient loss due to leaching or fixation reactions. Placement decisions depend on the crop and soil conditions, which interact to influence nutrient uptake and availability (Roberts, 2008).

Nitrogen fertilizers are expensive and crops are able to utilize only 50% of the nitrogen provided by them. Suitable fertilizers ~~s~~ application strategies which makes ~~s~~ some amount of added fertilizer, available for early growth and leftover part in later growth stages of the crop should be encouraged (Pandey *et al.*,

2019). Precision nutrient application by splitting the recommended dose of fertilizer may be functional in improving the sustainability of available N by preventing leaching or volatilization losses of fertilizers(Allen *et al.*, 2010). Split application refers to the application of a total dose of fertilizer in fragments for synchronization of supply of applied nutrients with the need of plants as well as the ability of plants to use these nutrients. If the full dose of nutrients especially N is applied in a single dose as pre-planting or at planting, the window for the potential loss of these nutrients ~~is~~ through various pathways is always open. By postponing a portion of N treatment until the crop is better able to utilize the nutrient, plants take up the nitrogen more quickly and efficiently. Fixation of K in the soil can be minimized by its split application which facilitates enhanced access and use efficiency by plants (Romheld and Kirkby, 2010).

3. Materials and Methodology:

3.1. ~~Experimental.~~ Experimental Site and Climate

A field experiment was conducted at the Regional Research Station, CCS Haryana Agricultural University, Karnal located at the latitude of 29°43'42.19" N and longitude of 76°58'49.88" E and at an altitude of 253 meters above mean sea level. It is roughly equidistant and almost midway ~~between from~~ New Delhi and Chandigarh.

The climate is sub-tropical with a mean maximum temperature ranging between 34-39 °C in summer and a mean minimum temperature ranging between 6-7 °C in winter. Most of the rainfall is received during the months of July to September and few showers ~~from during~~ December to late spring.

3.2 Soil of Experimental Plot

The field at Regional Research Station, CCSHAU, Karnal selected for conducting the experiment was uniform in fertility gradient. ~~The initial~~Initial soil fertility status of the experimental field was determined prior to planting of crop, for which four representative soil samples were collected randomly from the entire field at a depth of 0-30 cm before implementing the final layout of the experiment. The analysis was carried out by strictly following established protocols and standard procedures. From the interpretation of results obtained after soil analysis, it was interpreted that soil exhibited clay loam texture, alkaline in reaction, medium in OC content, low in available N and medium in P and K content.

Table 1: Initial physio-chemical properties of the soil under experiment

S. No.	Parameter	Value/ Category	Analytical Method Used
1.	Soil Texture	Clay	International pipette method (Piper,1966)

		loam	
2.	pH _(1:2)	8.28	pH was determined using digital pH meter
3.	EC _(1:2) (dS m ⁻¹)	0.25	EC was determined using digital EC meter
4.	OC (%)	0.48	Wet digestion method (Walkley and Black, 1934)
5.	Av. N (kg ha ⁻¹)	111.55	Alkaline Potassium Permanganate Method (Subbiah and Asija, 1956)
6.	Av. P (kg ha ⁻¹)	18.88	Sodium Bicarbonate Extractable P method (Olsen <i>et al.</i> , 1954)
7.	Av. K (kg ha ⁻¹)	201.40	Ammonium Acetate Extractable K method (Jackson, 1973)

3.3 Treatments and Layout of the Experiment:

The experiment was arranged in Split Plot Design with three replications. The experiment was designed with two main plots (Mode of fertilizer application i.e. Broadcasting and Band Placement) and four sub-plots (No. of split application i.e. 5, 6, 7 and 3 splits). The details of the experiment are given below:

3.4 Treatment Details:

3.4.1 Main plot treatments (two): Methods of fertilizer application

B1: Broadcasting

B2: Band placement

3.4.2 ~~chart~~ Chart 1: Sub Plot treatments (four): Number of splits of the recommended dose of fertilizer:

T1	Recommended dose of N and K in five splits (Basal 10% and remaining dose at 45, 75, 90 and 120 DAP)
T2	Recommended dose of N and K in six splits (Basal 10% and remaining dose at 45, 75, 90, 120 and 150 DAP)
T3	Recommended dose of N and K in seven splits (Basal 10% and remaining dose at 45, 75, 90, 120, 150 and 180 DAP)
T4	Recommended dose and schedule of nutrient applications (Half of total N and full dose of P and K at planting and rest of the N at 45 and 90 DAP)

3.5 Soil analysis:

Soil samples were collected at 0, 15, 30, 45, 75, 90, 120, 150, and 180 days after planting (DAP) and at harvesting. These collected soil samples were analyzed for available N and K during different crop growth stages. For the determination of pH, EC, Soil organic carbon (SOC) and available P soil samples were analyzed at the time of sowing and at harvesting.

4. Results and Discussion

4.1 Effects of Different Treatments on Soil Macronutrient Status

4.1.1. Available Nitrogen

The data relating related to the effect of different main and subplot treatments on available N at different time intervals presented in Table 2 indicates an overall appraisal in available N status at harvesting under all the treatments compared to the initial. This augmentation observed in N status might be directly linked to the addition of N to soil through the applied fertilizer in a balanced manner as also reported by Dubey *et al.* (2012). Further, examination of data elucidated that upto 45 DAP no significant increase was noticed in the available N content of soil under both sub or main plot treatments as at the time of sowing similar amount of a basal dose was applied in all the treatments except T4 (3 splits) where half of the total N dose was applied at the time of sowing. This, however, did not culminated in the build-up of N in T4 during the initial stage which might be due to volatilization losses owing to high temperature and absence of shade due to negligible crop cover at the time of sowing. As reported by Bouwman and Boumans (2002), potential NH₃ emissions are greatest for urea among synthetic fertilizers, for which cumulative losses may account as high as 60% of applied N (Sommer *et al.*, 2004; Rochette *et al.*, 2009). After 45 DAP available N content in soil was remarkably influenced by both main and subplot treatments.

Among the methods of fertilizer application, significantly higher values of available N were recorded in band placement (B2) compared to broadcast (B1) at all the stages after 45 DAP upto harvest. Surface broadcasting of urea may have contributed to higher volatilization and leaching losses while emissions were reduced by placement in subsurface band placement which lead to the trapping of the majority of ammonical N in the soil. Similar findings were elucidated by Prasertsaket *et al.* (2002) and de Castro *et al.* (2016). Moreover, broadcasting has an additional disadvantage of N leaching in deeper layers as reported by Chen *et al.* (2016). The amount of available N gradually increased in both main and subplot treatments with the application of each split dose which is consistent with the findings of Everaarts and Willigen (1999).

Among the number of splits, T4 reported significantly higher values of available N from 75 DAP (134.51 kg ha⁻¹) upto 90 DAP (147.08 kg ha⁻¹) and thereafter it was significantly highest in T1 from 120 DAP (142.63 kg ha⁻¹) to 150 DAP (162.11 kg ha⁻¹) which corresponds to the higher amount of fertilizer N applied in these time intervals through the respective split applications under these treatments. These findings are in congruence with those of (Kamble and Kathmale, 2015). However, from 180 DAP (153.35 kg ha⁻¹) till the harvest (145.93 kg ha⁻¹) decreasing trend was observed in available N in the soil under T1 due to no further fertilizer application. T4 also registered a consistent fall in available N content in the soil after 90 DAP (147.08 kg ha⁻¹) upto harvesting (118.31 kg ha⁻¹). In T2 from 75 DAP (125.63 kg ha⁻¹) upto 180 DAP (159.21 kg ha⁻¹) increase in available N was observed which decreased at harvest (151.78 kg ha⁻¹) its. However, in T3 available N content in soil kept on increasing significantly from 75 DAP (124.46


kg ha⁻¹) upto harvesting (157.01 kg ha⁻¹) both of which can be attributed to retention of unutilized urea applied from later splits in the soil as sugarcane crop utilizes most of the applied nutrient upto 150 DAP (Dayo-Olagbendedet *al.*, 2019). Interaction between different treatments (main and sub-sub-treatments) and time intervals was significant in influencing available N status in soil.

Table 2: Effect of different methods of fertilizer application and numbers of splits of N and K on available nitrogen of soil

Method of fertilizer application	Available N (kg ha ⁻¹)									
	0 DAP	15 DAP	30 DAP	45 DAP	75 DAP	90 DAP	120 DAP	150 DAP	180 DAP	Harvest
B1-Broadcasting	110.44	116.70	119.54	122.80	125.73	135.11	139.21	142.01	142.80	140.60
B2-Band Placement	110.80	118.72	121.79	124.62	131.86	141.55	145.17	148.10	148.66	145.91
SEm±	1.03	0.71	0.69	0.37	0.43	0.50	0.43	0.78	0.68	0.72
CD (P=0.05)	NS	NS	NS	NS	2.82	3.301	2.85	5.14	4.49	4.72
Number of splits of N and K										
T1-5 splits	110.45	117.58	120.30	123.11	130.66	142.63	153.10	162.11	153.35	145.93
T2 -6 splits	110.48	117.50	120.50	123.61	125.63	133.76	141.02	147.26	159.21	151.78
T3 -7 splits	110.84	118.01	121.13	124.00	124.46	129.86	135.60	139.55	146.96	157.01
T4-3 splits	110.72	117.76	120.73	124.13	134.51	147.08	139.06	131.31	123.40	118.31
SEm±	0.66	0.81	0.83	0.64	0.64	0.69	0.67	1.01	0.87	0.87
CD (P=0.05)	NS	NS	NS	NS	2.02	2.15	2.08	3.15	2.71	2.72


Need to be explain NS in the below of table

Fig 1: Effect of different methods of fertilizer application and numbers of splits of N and K on available nitrogen of soil



Comment [K3]: Figures 1 and 2 are the same. Please choose one of them

Fig 2: Effect of different methods of fertilizer application and numbers of splits of N and K on available nitrogen of soil (arrows represents the timing of fertilizer application)



4.1.2 Soil Available K

The data presented in Table 3 revealed a fair appreciation of ~~in~~-available K status at the time of harvesting compared to initial status under both main and subplot treatments. The rise in available K might be attributed to the frequent addition of K through a number of split applications of RDF throughout the crop period at different time intervals which enriched the soil in available K as argued by Pandey *et al.* (2019) and Tariq *et al.* (2011). Moreover, Singhet *et al.* (2019) opined that K drawn by roots from the lower layer is deposited in the surface layer which finally leads to enhancement in available K status in Alfisols which might be the case here as sugarcane is a ~~deep-deep~~-rooted crop capable of drawing sizeable amount of K from the subsurface layer.

Among different methods of fertilizer application, significantly higher values of available K were recorded in band placement (B2) compared to broadcast (B1) at all the time intervals upto harvest. However, the difference noted was non-significant at 0, 150, 180 DAP and at harvest. This might be accredited to the slow downward movement of K arising from accelerated adsorption of K on soil colloidal complex as a result of wider soil fertilizer (Nandet *et al.*, 2019) contact in broadcasting. Moreover, K uptake might be higher in band placement leading to more exhaustion of nutrient applied in the vicinity of plant roots. Similar observations were annotated by Kraska *et al.* (2021).

At 0 DAP, available K content observed in all the ~~sub-sub~~-treatments ~~were-was~~ par with each other. Among different number of ~~applied~~ splits ~~applied~~, treatment receiving fertilizer in three splits (T4) reported the highest values of available K upto 75 DAP (225.30 kg ha⁻¹) as it received the whole amount

of RDF to be applied before 90 DAP which was higher than other treatments. Thereafter treatment T1 exhibited significantly higher available K values upto 150 DAP (222.06 kg ha⁻¹) while at 180 DAP and at the time of harvesting, significantly higher values of available K were observed in treatments T2 (225.10 kg ha⁻¹) and T3 (227.48 kg ha⁻¹) respectively which might be due to higher application of fertilizer K compared to other treatments during these time intervals through the scheduled splits in respective treatments. These findings are in compliance with those of and Pandey *et al.* (2019) who reported an increase in soil available K ~~in~~ when applied in splits. Treatment T4 reported a fall in available K content in soil 45 DAP after an initial rise at 15 DAP upto harvesting while in T1 and T2 it ~~was~~ decreased after 150 and 180 DAP respectively upto harvesting. This drop in soil might be attributed to ~~non-non-~~ application fertilizers at these intervals as per findings of Nandeti *et al.* (2019). Interaction between different treatments (main and ~~sub-sub-~~ treatments) and time intervals ~~was-were~~ significant in influencing available K status in soil.

Table 3: Effect of different methods of fertilizer application and numbers of splits of N and K on available potassium of soil

Methods of fertilizer application	Available K ₂ O (kg ha ⁻¹)									
	0 DAP	15 DAP	30 DAP	45 DAP	75 DAP	90 DAP	120 DAP	150 DAP	180 DAP	Harvest
B1-Broadcasting	201.40	216.08	218.33	221.19	222.02	216.97	213.37	217.84	220.43	220.50
B2-Band Placement	201.25	213.69	215.82	218.69	217.11	212.59	206.71	209.81	213.37	214.6
SEm±	0.53	0.09	0.39	0.04	0.47	0.60	1.00	1.55	2.44	1.16
CD (P=0.05)	NS	0.63	2.59	0.28	3.12	3.96	6.57	NS	NS	NS
Number of splits of N and K										
T1 -5 splits	201.28	206.66	208.61	210.70	220.83	218.33	215.25	222.06	220.41	218.75
T2 -6 splits	201.50	206.01	207.31	209.40	217.78	215.28	211.16	217.88	225.10	223.76
T3 -7 splits	201.55	205.10	207.11	210.40	214.40	211.73	208.23	212.78	220.50	227.48
T4-3 splits	200.00	241.76	245.26	249.26	225.30	213.78	205.53	202.59	201.59	200.21
SEm±	1.00	0.75	0.82	0.76	0.92	0.90	0.72	0.67	1.06	0.64
CD (P=0.05)	NS	2.33	2.56	2.37	2.88	2.82	2.25	2.57	3.28	1.99

Fig 3: Effect of different methods of fertilizer application and numbers of splits of N and K on available potassium of soil





Fig 4: Effect of different methods of fertilizer application and numbers of splits of N and K on available potassium of soil (arrows represents timing of fertilizer application)



4.1.3 Available Phosphorus

It is clear from the perusal of data in Table 4 that at harvesting, different splits of N and K applied had a non-significant effect on P₂O₅ content of the soil. When effect of the broadcasting and band placement were compared it was seen that soil P₂O₅ content recorded was significantly higher in broadcasting (B1) than the band placement (B2) which might be due to fixation and retention of unutilized P as a result of higher soil to fertilizer contact (Rehman *et al.*, 2006). An increment in available

P values was noticed in all the treatments at harvesting which might be associated with the application of mineral fertilizers at the time of sowing (Kraska *et al.*, 2021).

Table 4: Effect of different methods of fertilizer application and numbers of splits of N and K on available phosphorus and soil organic carbon of soil

Method of fertilizer application	P ₂ O ₅ At sowing (kg ha ⁻¹)	P ₂ O ₅ At harvest (kg ha ⁻¹)	SOC At sowing (%)	SOC At harvest (%)
B1 -Broadcasting	18.62	24.63	0.48	0.50
B2-Band placement	19.62	21.68	0.48	0.51
SEm±	0.10	0.38	0.03	0.01
CD (P=0.05)	NS	3.057	NS	NS
Number of splits of N and K				
T1-5 splits	18.71	22.36	0.47	0.48
T2 -6 splits	18.88	22.96	0.47	0.49
T3 - 7 splits	19.36	23.28	0.48	0.50
T4- 3 splits	18.80	24.01	0.48	0.48
SEm±	0.40	0.58	0.06	0.02
CD (P=0.05)	NS	NS	NS	NS

4.2 Effects of Different Treatments on Soil Chemical Properties

4.2.1 Soil Organic Carbon

It is very much clear from the data presented in Table no. 4 that no significant changes were recorded in the organic carbon content of initial soil samples and those collected at harvest. Neither method of fertilizer application and nor the number of split applied splits had any significant influence on the organic content of the soil. However, a slight increase in numerical value was observed at harvesting with respect to organic carbon content at harvesting which might be due to the suppressive effect of N fertilization on microbial mineralization of soil organic matter. Addition of ammonium molecules through urea application in splits throughout the life cycle of crop reduced oxidative enzyme activity (Mahal *et al.*, 2019) by inhibiting the synthesis of ligninolytic enzymes released by ligninolytic fungi (Khalafalla and Hamed, 2015).

4.2.2 Soil pH and EC (dS m⁻¹)

Changes undergone in pH and EC of the soil during the crop growth are presented in Table 5 which shows no remarkable variations in pH and EC of the soil either under the main plot treatments (methods of fertilizer application) or subplot treatments (application of RDF in different splits). This might be due to the strong buffering capacity of the clay soils of the experimental site. However a slight decrease in soil pH values was observed in all the treatments at

harvesting relative to initial values which might be the release of root exudates and accumulation of other decomposition products which are acidic in nature. Moreover, application of urea also had an acidic residual effect in the soil causing a fall in pH values (Dayo-Olagbende *et al.*, 2019; Kraska *et al.*, 2021).

Table 5: Effect of different methods of fertilizer application and numbers of splits of N and K on pH and EC of soil

Method of fertilizer application	Initial pH	pH At Harvest	Initial EC (dSm ⁻¹)	EC At Harvest (dSm ⁻¹)
B1-Broadcasting	8.23	8.13	0.25	0.16
B2- Band placement	8.20	8.16	0.26	0.16
SEm±	0.00	0.01	0.01	0.03
CD (P=0.05)	NS	NS	NS	NS
Number of splits of N and K				
T1- 5 splits	8.15	8.12	0.26	0.16
T2 - 6 splits	8.25	8.15	0.26	0.16
T3 - 7 splits	8.21	8.15	0.25	0.17
T4- 3 splits	8.25	8.17	0.25	0.17
SEm±	0.02	0.06	0.02	0.06
CD (P=0.05)	NS	NS	NS	NS

5. Conclusion:

Based on the observations recorded and intensive interpretation of the results it can be concluded that band application of N fertilizer is an effective strategy for enhancing the availability of N in the soil while banding of K is effective in enhancing the absorption of nutrients by plant roots. However, increasing the number of splits beyond five is not helpful to crop as most of the absorption of required nutrients occur during the vegetative stage and thus fertilizer applied at the latter stages remains unutilized and is lost through various mechanisms. Providing an excessive amount of fertilizer at sowing is also not very productive as plant roots are not developed in the early stages and thus applied N can be lost through volatilization and leaching. Thus it can be said that increasing and decreasing the number of splits before or after the peak nutrient requirement stages of a crop are not fruitful and should be adjusted accordingly. In the case of sugarcane dividing the RDF into five splits emerged out to be the best treatment.

6. References:

Allen, D. E., Kingston, G., Rennenberg, H., Dalal, R. C., and Schmidt, S. (2010). Effect of nitrogen fertilizer management and waterlogging on nitrous oxide emission from subtropical sugarcane soils. *Agriculture, Ecosystems & Environment*, **136**(3-4):209-217

- Bouwman, A. F., Boumans, L. J. M., and Batjes, N. H. (2002). Estimation of global NH₃ volatilization loss from synthetic fertilizers and animal manure applied to arable lands and grasslands. *Global biogeochemical cycles*, **16**(2): 8-1.
- Chen, Z., Wang, H., Liu, X., Liu, Y., Gao, S. and Zhou, J. (2016). Effect of N fertilizer placement on the fate of urea ¹⁵N and yield of winter wheat in Southeast China. *PlosOne* ,**11**(4): 1-13
- Dayo-Olagbende, G. O., Ewulo, B. S., and Akingbola, O. O. (2019). Combined effects of tithonia mulch and urea fertilizer on soil physico-chemical properties and maize performance. *Journal of Sustainable Technology*, *10*(1), 86-93
- de Castro, S. G. Q., Decaro, S. T., Franco, H. C. J., Magalhães, P. S. G., Garside, A., and Mutton, M. A. (2017). Best practices of nitrogen fertilization management for sugarcane under green cane trash blanket in Brazil. *Sugar Tech*, *19*(1), 51-56.
- Dubey, V., Patel, A. K., Shukla, A., Shukla, S., and Singh, S. (2012). Impact of continuous use of chemical fertilizer. *International Journal of Engineering Research and Development*, *3*(11): 13-16.
- Everaarts, A. P., and De Willigen, P. (1999). The effect of nitrogen and the method of application on yield and quality of broccoli. *Netherlands Journal of Agricultural Science*, **47**(2), 123-133.
- Farmaha, B.S., Fernández, F.G. and Nafziger, E.D.(2013) Distribution of soybean roots, soil water, phosphorus and potassium concentrations with broadcast and subsurface-band fertilization. *Soil Science Society of America Journal*, **76**:1079-1089
- Fernández, F.G. and White, C. (2012). No-till and strip-till corn production with broadcast and subsurface-band phosphorus and potassium. *Agronomy Journal*,**104**(4):996-1005
- Fixen, P. E. (2020). A brief account of the genesis of 4R nutrient stewardship. *Agronomy Journal*, **112**(5):4511-4518.
- Jackson, M. L. (1973). Soil chemical analysis, pentice hall of India Pvt. Ltd., New Delhi, India, 498, 151-154
- Kamble, B. M., and Kathmale, D. K. (2015). Effect of different levels of customized fertilizer on soil nutrient availability, yield and economics of onion. *Journal of Applied and Natural Science*, **7**(2): 817-821.
- Khalafalla, M. Y., and Hamed, M. H. (2015). Impact of nitrogen fertilization on soil organic carbon decomposition. *Alexandria Science Exchange Journal*, **36**: 381-389.
- Kraska, P., Andruszczak, S., Gierasimiuk, P., and Rusecki, H. (2021). The Effect of Subsurface Placement of Mineral Fertilizer on Some Soil Properties under Reduced Tillage Soybean Cultivation. *Agronomy*, **11**(5): 859.
- Katyal, J. C., and Chaudhari, S. K. (2021). Do Fertilizers Protect or Punish Soil Health?. *Indian Journal of Fertilisers*, **17**(12): 1242-1281.
- Mahal, N. K., Osterholz, W. R., Miguez, F. E., Poffenbarger, H. J., Sawyer, J. E., Olk, D. C., Archontoulis, S.V. and Castellano, M.J., (2019). Nitrogen fertilizer suppresses mineralization of soil organic matter in maize agroecosystems. *Frontiers in Ecology and Evolution*, *7*, 59.
- Nand, M. M., Singh, S. P., Kumar, A., and Alam, M. M. (2019). Potassium fractions affected by split application of potassium in rice (*Oryza spp.*) in calcareous soil of North Bihar. *IJCS*, **7**(2): 964-966.

- Olsen, S.R., Cole, C.V., Watanabe, F.S. and Dean, L.A. (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate. *Circular United States Departments of Agriculture*, 939
- Pahalvi, H. N., Rafiya, L., Rashid, S., Nisar, B., and Kamili, A. N. (2021). Chemical fertilizers and their impact on soil health. In *Microbiota and Biofertilizers*, 2 :1-20 Springer, Cham.
- Pandey, D., Bhatnagar, A., Chandra, S. and Tewari, S. (2019). Soil nutrient balance under influence of differential placement of fertilizer doses and potassium splitting in maize (*Zea mays* L.). *Journal of Pharmacognosy and Phytochemistry*, 18(4): 1568-1572
- Piper, C.S. (1966). Soil and water analysis. Published by University of Adelaide, Australia
- Prasertsak, P., Freney, J.R., Denmead, O.T., Saffigna, P.G., Prove, B.G. and Reghenzani, J.R. (2012). Effect of fertilizer placement on nitrogen loss from sugarcane in tropical Queensland. *Nutrient Cycling in Agrosystems*, 62: 229-239
- Randive, K., Raut, T., and Jawadand, S. (2021). An overview of the global fertilizer trends and India's position in 2020. *Mineral Economics*, 34(3): 371-384.
- Rochette, P., Angers, D. A., Chantigny, M. H., MacDonald, J. D., Gasser, M. O., and Bertrand, N. (2009). Reducing ammonia volatilization in a no-till soil by incorporating urea and pig slurry in shallow bands. *Nutrient Cycling in Agroecosystems*, 84(1): 71-80.
- Rehman, O., Zaka, M. A., Rafa, H. U., and Hassan, N. M. (2006). Effect of balanced fertilization on yield and phosphorus uptake in wheat-rice rotation. *J. Agric. Res*, 44(2): 105-115.
- Roberts, T.L. (2008). Improving nutrient use efficiency. *Turkish Journal of Agriculture and Forestry*, 32: 177-182
- Romeheld, V. and Kirkby, E.A. (2010). Research on potassium in agriculture: Needs and prospects. *Plant Soil*. 335:155-180.
- Shukla, S.K., Yadav, S.K., Lal, M. and Pathak, A.D. (2018). Low cost technologies in sugarcane agriculture. ICAR- All India Coordinated Research Project on Sugarcane, AICRP(S) Technical Bulletin- No.3: 20-23
- Singh, M., Wanjari, R. H., Kumar, U., and Chaudhari, S. K. (2019). AICRP on long-term fertilizer experiments: salient achievements and future directions. *Indian Journal of Fertilisers*, 15(4): 356-372.
- Sommer, S. G., Schjoerring, J. K., and Denmead, O. T. (2004). Ammonia emission from mineral fertilizers and fertilized crops. *Advances in agronomy*, 82:557-622
- Subbiah, B.V. and Asija, G.L. (1956). A rapid procedure for the determination of available nitrogen in soils. *Current Scienc*, 25:259-60
- Tariq, M., Saeed, A., Nisar, M., Mian, I. A., and Afzal, M. (2011). Effect of potassium rates and sources on the growth performance and on chloride accumulation of maize in two different textured soils of Haripur, Hazara division. *Sarhad Journal of Agriculture*, 27(3): 415-22..
- Walkley, A.J. and Black, C.A. (1934). Estimation of soil organic carbon by the chromic acid titration method. *Soil Science* 37:29-38.

UNDER PEER REVIEW

