

Original Research Article

Parasites Associated with Open markets Vegetables: Implication for Public Health Interventions in EthiopeEast, Southern Nigeria

Abstract

Objectives: Vegetables have been reported to be a mechanical means of transmission of human intestinal parasites. This study is aimed to ascertain the association between parasites and vegetables sold in Ethiope East, Delta State, Nigeria. **Methods:** Two hundred and seventy (270) vegetables were subjected to sedimentation and zinc sulphate floatation techniques for parasitic investigations. **Results:** Of the 270 vegetables examined, 184(68.15%) were infected. *T. occidentalis* had the highest prevalence (9.26%), closely followed by *B. oleracea* and *D. carota* with a value of 8.89% and 8.15% respectively. The highest parasite prevalence was observed in Abraka (78.89%), followed by Eku (72.22%) and Okpara water-side (53.33%). Abraka had its highest prevalence in *D. carota* and *C. sativum* with a prevalence of 11.11 and 10.00% respectively. Statistically, there was no significant difference in the contaminated vegetable ($p > 0.05$) but a considerable significant difference was noted in the market locations ($p < 0.05$). The highest percentage occurrence of parasites was *A. lumbricoides* with a percentage occurrence of 23.49. The most prevalent parasites in Abrakawere *Ascaris lumbricoides* (45.56%) and *Entamoeba histolytica* (30.00%). There was significant effect ($p < 0.05$) in the prevalence of parasites ($p = 0.0003$; $df = 6$; $F = 11.07$) and markets locations ($p = 0.0019$; $df = 2$; $F = 11.01$). The Principal Component Analysis showed that the study locations positively influence the occurrence of parasites with Abraka location accounting for 95.99% of the total variance. The analysis further showed that *A. lumbricoides*, *Trichuris trichura*, and hookworm prevalence were strongly correlated to Abraka and Okpara water-side locations. **Conclusion:** Given the public health importance of the parasites isolated, there is a need for improved environmental sanitation, personal hygiene and surveillance systems in vegetable transportation, storage and means of displays in Ethiope East and Delta State at large.

Keywords: Vegetable parasites, Prevalence, Abraka, Eku, Okpara water-side, Public health.

1.0 INTRODUCTION

The voracious consumption of vegetables has increased in recent years because of their attendant health and nutritional benefits [1]. Vegetables form a key component of a healthy diet, highly beneficial for the maintenance of health and the prevention of diseases [2]. Vegetables are low in calories, fat and sodium, and supply fiber, vitamins, minerals, and other health-promoting phytochemicals. The importance of vegetables in supplying most of the vitamins and minerals such as carotene, ascorbic acid, riboflavin, iron, iodine and calcium cannot be over-emphasized.

Consumption of a diet rich in vegetables may reduce the risk of stroke, help reduce blood cholesterol levels and lowers the risk of cardiovascular diseases, type 2 diabetes, and certain cancers of the mouth, stomach, colon and rectum [3]. Potassium in some vegetables may also reduce the risk of kidney stones, bone-calcium loss and maintain healthy blood pressure; while antioxidants in vegetables help protect the body from oxidant stress, diseases and cancers by boosting immunity. Fibers absorbed in the colon retain moisture and ease faecal excretion, and defenses against hemorrhoids, colon cancer, and protracted constipation. The foliate in vegetables helps in the formation of red blood cells and reduces the risk of neural tube defects, spin Bifida and anencephaly during foetal development in pregnancy. Consumption of vegetables generally helps to maintain healthy life and body weight [3].

However, eating faecal-contaminated vegetables contributes significantly to the spread of intestinal parasite illnesses[4]. The increased consumption of fresh vegetables has been linked to several documented cases of parasitic diseases. In most cases, the vegetables are eaten raw or lightly cooked to retain their natural taste and to preserve heat-labile nutrients. This practice, however, facilitates the transmission of parasitic infections [5,6]. These vegetables are major risk factors for the prevalence and transmission of intestinal parasites in developing nations principally due to poor personal hygiene and sanitation[7].

Vegetables have been incriminated in the transmission of amoebiasis, giardiasis, toxoplasmosis, cryptosporidiosis, taeniasis, fascioliasis, strongyloidiasis and major helminthiasis, as well as eggs and larvae of *Cyclospora cayatanensis*, *Isospora* species, *Hymenolepis nana*, *Toxocara* species, *Trichostrongylus* species, *Enterobius vermicularis*[6,8]. Universally, these human intestinal parasitic infections exhibit various clinical symptoms calling for public health interventions [9]. Infants, the elderly, and immuno-compromised persons are particularly vulnerable and at more risk of parasitic infections [10]. However, environmental epidemiological factors for parasites prevalence have not been properly elucidated but outbreaks of amoebic dysenteries, giardiasis and balantidiasis caused by *Daucus carota* (carrot), *Brassica oleracea* (cabbage), *Lactuca sativa* (lettuce), and *Telfairia occidentalis* (fluted pumpkin leaf) have been documented.

Globally, approximately 3.5 billion people are infected with intestinal parasitic infections, of which 450 million exhibit symptoms with an annual mortality of 200 thousand reported [11,12]. Nearly 300 million people in resource-limited countries suffer from severe significant morbidity[13,14]. Parasitic infections are widespread in Nigeria and other Sub-Saharan African countries where they afflict various demographic groups [15]. The second most frequent factor contributing to outpatient morbidity in Nigeria is due to intestinal parasitic infections [16]. These infections have been linked to several physical and mental health issues, including diarrhea, anemia, stunting, physical weakness, poor educational performance, and growth retardation in children[15].

Intestinal parasites are primarily transmitted by fecal-oral route, mostly via ingestion with contaminated food and water or during direct hand-to-mouth contact [16]. Vegetables usually consumed raw are potential sources of infection. Empirical data have explicitly implicated *A. lumbricoides*, *E. histolytica*, *E. vermicularis*, *Fasciola gigantica*, *Giardia lamblia*, *Hymenolepis nana*, *Cryptosporidium parvum*, *Taenia saginata/solium*, *Trichuris trichiura*, and species of hookworms, *Cyclospora* and *Toxocara* infect humans who consume contaminated vegetables without cooking or proper washing [17,18]. Delta State, Nigeria is a state where many intestinal parasites abound due to poor hygiene and sanitation practices[15,9,10,19,20,21]. There is a paucity of data on vegetable parasites in Ethiopia East, Delta State, Nigeria, despite their role in the transmission of pathogens. Specifically, this study is aimed to: ascertain the association of parasites and vegetables sold in Ethiopia East, Local Government Area; and to determine the spatial prevalence of vegetable parasites in Abraka, Eku and Okpara water-side communities.

2.0 MATERIALS AND METHODS

2.1 Brief Description of Study Area.

This study was carried out in September 2021 in Abraka, Eku and Okpara water-side situated between latitude $5^{\circ} 54'$ and longitude $5^{\circ} 40'$ (Fig. 1) Abraka, Eku and Okpara-water side are three major towns in Ethiopia East, Delta State, Nigeria. The inhabitants are mainly farmers and traders together with civil servants. Sellers from Abraka, Eku and Okpara-water side markets mostly get their vegetables supplies directly from the local farmers and traders from Northern Nigeria. The inhabitants of Delta and neighbouring State usually patronize these three (3) markets for vegetables. Toilet amenities are absent in most of the markets, hence traders resort to the use of open defecation, especially for children and disposed of within the markets. Sewage disposal is also poor and there is a heap of refuses

Comment [L1]: Folostop

where faeces are often deposited in the markets. These faecal matters are washed during a heavy downpour of rain to the surroundings.




Fig. 1: Map of Ethiope East L.G.A, Delta State, Nigeria showing study locations (Source: Ito Ed.’ E, 2023; Department of Animal and Environmental Biology, DELSU, Abraka)

3.2 Sample collection

The collection of samples was carried out with slight modification to the method described by Endaleet *al.* [14]. Two hundred and seventy (270) samples of nine different types of vegetables (*Talinum triangulare* (water leaf), *Cucumis sativum* (cucumber), *Daucus carota* (carrot), *Amaranthus spinosus* (spinach), *Brassica oleracea* (cabbage), *Lactuca sativa* (lettuce), *Corchorus olitorius* (white jute), *Telfairia occidentalis* (Fluted pumpkin leaf) and *Celosia argentea* (quail grass) were purchased from farmers and vendors in three local markets, namely; Abraka, Eku and Okpara-water-side market in Ethiope East LGA, Delta State, Nigeria. Five (5) of each sample was collected randomly per week in each market from ten (10) places randomly selected. The samples were collected separately in a properly labeled plastic bag and transported to the laboratories Centre for Advance Research Delta State University, Abraka for parasitological examination.

3.3 Parasitological examination

Standard parasitological methods (Sedimentation and zinc sulphate floatation techniques) previously described were used to examine the vegetables collected [8,22].

3.4 Statistical Analysis

Data collected were analyzed using PAST software version 4.11. One-way and Two-way Analysis of Variance (ANOVA) were used to determine the association between parasitic contamination rate and types of samples. Values of *p* less than or equal to 0.05 were considered significant.

3.0 RESULTS

Of the 270 vegetables collected, 184 were infected giving a total prevalence of 68.15%. The prevalence of parasites in the vegetable is presented in Table 1. *T. occidentalis* had the highest prevalence (9.26%), closely followed by *B. oleracea* and *D. carota* with a value of 8.89% and 8.15% respectively. However, the least infected vegetable was *C. sativum* (Table 1). Statistically, no significant difference ($p=0.271$; $F=1.56$) was observed in the prevalence of vegetable parasites.

Table 1. Prevalence and Distribution of Vegetables Contaminated with Parasites in Ethiope East, Delta State

Vegetables	N.E	N. I[%]	Overall Prevalence
<i>T. triangulare</i>	30	19(63.33)	7.04
<i>C. sativum</i>	30	16(53.33)	5.93
<i>D. carota</i>	30	22(73.33)	8.15
<i>A. spinosus</i>	30	21(70.00)	7.78
<i>B. oleracea</i>	30	24(80.00)	8.89

Comment [L2]: Write the way in which the number was made

Comment [L3]: What is mean

Comment [L4]: What is mean

<i>L. sativa</i>	30	20(66.67)	7.41
<i>C. olitorius</i>	30	19(63.33)	7.04
<i>T. occidentalis</i>	30	25(83.33)	9.26
<i>C. argentia</i>	30	18(60.00)	6.67
Total	270	184	68.15

Table 2 presents the spatial percentage of infections of vegetables in the 3 selected locations. The highest prevalence was obtained in Abraka with a value of 78.89%, followed by Eku with 72.22%. Abraka had its highest prevalence in *D. carota* and *C. sativum* with a prevalence of 11.11 and 10.00% respectively. Similarly, Eku had its prevalence peak for *T. occidentalis* at 11.11% while its least prevalence (4.44%) was observed in *C. sativum*. However, *C. sativum* was the least contaminated vegetable in the Okpara water-side location (Table 2). Statistical ANOVA revealed that there was no significant difference in the prevalence of parasites ($p > 0.05$; $p = 0.316$; $F = 1.286$) but a considerable significant difference was noted in the market locations ($p < 0.05$; $p = 0.0006$; $F = 6.90$).

Table 2. Prevalence of Parasites of vegetables sold at different markets in Ethiopia East Local Government Area

Vegetables	Markets					
	Abraka		Eku		Okpara Water-side	
	N. E.	N. I.(%)	N. E.	N. I.(%)	N. E.	N. I.(%)
<i>T. triangulare</i>	10	7(7.78)	10	7(7.78)	10	5(5.56)
<i>C. sativum</i>	10	9(10.00)	10	4(4.44)	10	3(3.33)
<i>D. carota</i>	10	10(11.11)	10	8(8.88)	10	4(4.44)
<i>A. spinosus</i>	10	7(7.78)	10	9(10.00)	10	5(5.56)
<i>B. oleracea</i>	10	8(8.89)	10	9(10.00)	10	7(7.78)
<i>L. sativa</i>	10	8(8.89)	10	6(6.67)	10	6(6.67)
<i>C. olitorius</i>	10	7(7.78)	10	7(7.78)	10	5(5.56)
<i>T. occidentalis</i>	10	8(8.89)	10	10(11.11)	10	7(7.78)
<i>C. argentia</i>	10	7(7.78)	10	5(5.56)	10	6(6.67)
Total	90	71(78.89)	90	65(72.22)	90	48(53.33)

The highest percentage occurrence of parasites was *A. lumbricoides* with a cumulative percentage occurrence of 23.49%. This was followed by *E. histolytica* with 17.46%. This study showed that there was a total of 481 individual parasites (Table 3).

Table 3. Percentage Occurrence of Parasitic ova and cyst found in all the three markets

Parasites	Number of parasites (%)
<i>E. coli</i>	27(5.61)
<i>E. histolytica</i>	84(17.46)
<i>A. lumbricoides</i>	113(23.49)
<i>G. lamblia</i>	34(7.07)
<i>T. trichuria</i>	66(13.72)
<i>F. gigantica</i>	51(10.60)
Hookworm	67(13.93)
Total	481

The highest prevalence in Abraka was observed in *A. lumbricoides* with a value of 45.56%, followed by *E. histolytica* (30.00%). The Highest value in Eku was *A. lumbricoides* with 52.22% (Table 4). Okpara had *A. lumbricoides* as its most prevalent parasite with 27.77%, lower than Abraka and Eku values. ANOVA showed that there was a significant difference ($p < 0.05$) in the prevalence of parasites ($p = 0.0003$; $df = 6$; $F = 11.07$) and market locations ($p = 0.0019$; $df = 2$; $F = 11.01$). The PCA cartesian plot showed that the study

Comment [L5]: What is mean

Comment [L6]: What is mean

locations positively influence the occurrence of parasites with Abraka location accounting for 95.99% of the total variance (Table 4 and Fig. 2).

Table 4. Prevalence of Parasites in Vegetables Sold in selected Markets in Ethiopia East

Parasites	Markets				Overall Prevalence
	N. E	Abraka N.I [%]	Ekus N.I [%]	Okpara Water-side N.I [%]	
<i>E. coli</i>	90	6(6.67)	12(13.33)	9(10.00)	10.00
<i>E. histolytica</i>	90	27(30.00)	43(47.78)	14(15.56)	31.11
<i>A. lumbricoides</i>	90	41(45.56)	47(52.22)	25(27.77)	41.85
<i>G. lamblia</i>	90	11(12.22)	13(14.44)	10(11.11)	12.59
<i>T. trichuria</i>	90	24(26.67)	27(30.00)	15(16.67)	24.44
<i>F. hepatica</i>	90	18(20.00)	21(23.33)	12(13.33)	18.89
Hook worm	90	23(25.56)	28(31.11)	16(17.78)	24.81
Eigenvalue		407.89	15.68	1.34	
Variation (%)		95.99	3.69	0.32	

[Note: The overall prevalence observed is based on 270 (90 x 3) vegetables examined]




Fig. 2: PCA cartesian plot of the study locations and parasites prevalence.

The PCA further indicated that *A. lumbricoides*, *T. trichura*, and hookworm prevalence were strongly correlated to Abraka and Okpara water-side community (Fig. 2). The vector planes explicitly revealed that the variations observed in this study were mainly contributed by the three major STH (*A. lumbricoides*, *T. trichura* and hookworm).

4.0 DISCUSSION

The study recorded the presence of seven (7) parasites associated with 270 vegetables in assent with Bekele and Shumbej, [8]. This number of parasites with contamination of 68.15% is comparatively higher than 42.0% and 42.60% previously documented [6,8]. The relatively high parasitic infections in this present study are a reflection of inadequate sanitation, poor personal hygiene and geographical locations of markets. However, the intestinal parasites in this study differ in specie composition and abundance compared to other research [22,23,24] because some of these vegetables like carrots, cabbage and lettuce were transported from the northern states into the study area for consumption. During the process of harvesting and transportation, some of these parasites might have been destroyed. Despite variation in isolated parasites, *A. lumbricoides* and hookworm had a high occurrence

in vegetables in this study. This could be due to the encystation of these parasites in adverse environmental conditions and their adherence easily to vegetable surfaces [15,16].

The results of this study are in tandem with many studies which reported high lettuce parasitic contamination[24,25]. This may be ascribed to the fact that the leaves are capable of harboring parasites in-between and in addition to its uneven surfaces on which parasites are attached more easily than other vegetables with smooth surfaces i.e., cucumber and this observation is in accordance with Damen *et al.*, [26].

The types of vegetables herein analyzed are widely produced and consumed in Ethiopia East, Delta State. Although the contamination rate found among them was statistically similar. The variation detected may be attributed to the anatomical differences of vegetable foliage. Cabbage has a high prevalence of 8.89%||this might be due to its large surface area and compact structure that provides better fixation and permanence of parasitic structures [27]. In addition, the flexibility of the leaves can facilitate contact with the soil with helminth ova probably found on the market floor [28].

According to Falavigna *et al.*[29], the remarkably high number and presence of helminth rather than protozoan structures detected in the present study are probably related to the practice of washing vegetables with the same bowl containing water in the field and, subsequently, at the point of sale before commercialization. These washing steps may be more efficient in the removal of protozoa, since the structure and size of helminths may hinder their removal from the surface of vegetables.

The presence of different parasitic species demonstrates that the population of Abraka, Eku and Okpara water-side is supposedly exposed to several debilitating diseases, such as hookworm infections, ascariasis, giardiasis and amebiasis. Indeed, a survey recently carried out in the city demonstrated that local children were infected with some of these pathogens[15,21]. The closely varied contamination could be due to similarity in parasitic life cycle, especially in relation to the elimination route of potentially contaminant structures, *i.e.* human and animal faeces. In this context, the presence of animals and humans infected with these pathogens in rural areas has been described to be associated with precarious sanitary installations, which could explain possible contamination of soil and water sources used for cultivation of the vegetables [30].

There was a high level of contamination of *E. histolytica* in this present investigation. These findings represent a public health concern since cysts are also eliminated with faeces into the environment. This elevated contamination corroborated the study conducted by Guilherme *et al.*[31] on vegetables marketed in *Maringá, Paraná*, Brazil, and is probably the result of the high viability of amoebic cysts in the environment[32].

Regarding the different markets studied, parasitic structures were detected in all of them, but the samples from the market were significantly more contaminated. Ethiopia East open markets are characterized by the sale of agricultural products grown in rural areas close to major centers, where farmers exhibit their products on the floor and consumers move between. Thus, the high percentage of contamination detected at the Eku and Abraka markets reinforces the considerable risk to which the local population is exposed as highlighted by Ito and Egwunyenga, [15].

5. CONCLUSIONS

There is a need for improved surveillance systems on vegetable parasites through appropriate farming procedures, improved transportation and storage facilities in the various market destinations considering the fact that vegetables in the Ethiopia East area were highly contaminated by intestinal parasites. This study suggested that inhabitants in the study area are at high risk of getting infections from eating vegetables daily which is of public health concern.

Comment [L7]: Delete

REFERENCES

1. Gboeloh LB, Seasonal Parasitic Contamination of Vegetables Marketed in Bori Central Market, Khana Local Government Area, Rivers State, Nigeria. *European J Biol. Biotech.* 2022; 3(4): 45-50.
2. Abubakar BM, Ahmad B, Dauda A, Gagman HA. Parasitic Contamination of Fruits and Vegetables Collected from Selected Local Markets in Katagum Region, Northeast Nigeria. *South Asian J Parasitol.* 2020;4(4): 36-43.
3. Khan W, Rafiqb N, Nawazc MA, Kabird M, Ur Z, Farooqie R, Rommanf M, (2022). Parasitic contamination of fresh vegetables sold in open markets: a public health threat. *Braz. J. Biol.* 2022; 82: e242614.
4. Yahaya O, Bishop, HG. Parasitic Contamination of Fresh Vegetables Sold in Some Major Markets and Farms in Zaria. *Sci. World J.* 2022; 17(1): 130-133.
5. Ajitha S, Vazhavandal G, Uma A, Prabhakaran N. Study on parasitic contamination of common edible fruits and vegetables sold in local markets of Tiruchirappalli, South India. *Indian J Microbiol Res.* 2020; 7(4): 362–368.
6. Fumilayo AJ, Mosunmola OJ, Kayode AI, Adedokun AR, Zagi HT, Buru AS. Intestinal Parasites Isolated in Vegetables Sold in Most Important Markets within Ilorin Metropolis. *J Trop Dis Public Health.* 2020; 8:342. doi:10.35248/2329-891X.20.8.342.
7. Tomass Z, Kidane D. Parasitological Contamination of Wastewater Irrigated and Raw Manure Fertilized Vegetables in Mekelle City and Its Suburb, Tigray, Ethiopia. *Momona Ethiop J Sci.* 2012;4(1):77. doi:10.4314/mejs.v4i1.74058.
8. Bekele F, Shumbej T. (2019). Fruit and vegetable contamination with medically important helminths and protozoans in Tarcha town, Dawuro zone, South West Ethiopia. *Res Reports Trop Med.* 10: 19-23.
9. Ito EE. Spatial distribution and prevalence of parasites vectored by *Periplaneta americana* in southern, Nigeria: Implication for intervention. *Asian J. Biol. Sci.* 2019; 12: 313-319.
10. Ito EE, Egwunyenga AO. Soil-Transmitted Helminthiasis in Aviara Community: An Observation from Primary School Children in Nigeria. *IMJ.* 2017; 24(2): 205 – 208.
11. Wegavehu T, Tsalla T, Seifu B, Teklu T. Prevalence of intestinal parasitic infections among highland and lowland dwellers in Gamo area, South Ethiopia. *BMC Public Health* 2013;13(1). doi: 10.1186/1471-2458-13-151.
12. Stofer J. *Adverse Health Effects of Intestinal Parasitic Infections in Rural Peruvian Clinic Patients.* Dayton, Ohio: Wright State University; 2014. Available from: <https://corescholar.libraries.wright.edu/mpb>. Accessed Dec. 2022.
13. Duedu K, Yarnie E, Tetteh-Ouarcoo P, Attah S, Donkor E, Aveh-Kumi P. A comparative survey of the prevalence of human parasites found in fresh vegetables sold in supermarkets and open-air markets in Accra, Ghana. *BMC Res Notes*, 2014;7(1). doi: 10.1186/1756-0500-7-836.
14. Endale A, Tafa B, Bekele D, and Tesfaye F. Detection of medically important parasites in fruits and vegetables collected from local markets in Dire Dawa, Eastern Ethiopia. *Glob J Med Res.* 2018;18(1):29–36.
15. Ito EE, Egwunyenga AO. Seasonal Prevalence, Intensity and Risk Factors of Geohelminthiasis among Children in Peri-Urban Settings in Ethiopia East, Southern, Nigeria. *Nig J Parasitol.* 2023; 43(1): www.ppsn.org.ng.
16. Ito EE, Agbure PA, Egwunyenga AO. Prevalence and Intensity of Soil-Transmitted Helminths and *Schistosoma* Co-Infections in Flood-Plain Communities, Southern, Nigeria. *J Egypt Soc Parasitol.* 2023;
17. Nyirenda SS, Bukowa KM, Banda WR, Mbewe J, Hamankolom F, Banda F. et al. Parasitic contamination of common vegetables sold in Lusaka, Zambia. *J Food Qual Hazards Control.* 2021;8:112-118.
18. Osafo R, Balali GI, Amisssah-Reynolds PK, Gyapong F, Addy R, Nyarko AA. Microbial and Parasitic Contamination of Vegetables in Developing Countries and Their Food Safety Guidelines. *J Food Qual.* 2022; <https://doi.org/10.1155/2022/4141914>
19. Eboh OJ, Okaka CE, Onuoha T. Prevalence of Gastrointestinal Parasites among School Children in Delta State, Nigeria. *Lond J Res Sci: Nat Formal.* 2022;22(8): 53- 60.
20. Omorodion AO, Nmorsi OPG, Isaac CC, Umukoro D, Ogbeneovo, and Akhile AO. Distribution of Intestinal Parasites among School-Age Children in Delta and Edo States of Nigeria. *PUJ.* 2012;5(2): 121-126.

21. Egwunyenga AO, Ataikuru PD. Soil-transmitted helminthiasis among school children in Ethiopia East Local Government Area, Delta State Nigeria. *AfrJBiotech*.2005;4(9): 938-941.
22. Istifanus WA, Panda SM. Parasitic agents in fresh fruits and vegetables sold in open markets in Bauchi, Nigeria. *JFood QualHazControl*. 2018;5: 84-88.
23. Alemu G, Nega M, Alemu M. Parasitic Contamination of Fruits and Vegetables Collected from Local Markets of Bahir Dar City, Northwest Ethiopia. *ResReports TropMed*. 2020; 11:17–25.
24. Oluwasola OO, Olufemi OA, Olufarati OF, Kayode BA, Onyiche ET. Parasitic contamination and public health risk of commonly consumed vegetables in Ibadan-Nigeria. *Pan AfrMedJ*.2020;36:126. Available at: <http://www.panafrican-med-journal.com/content/article/36/126/full/>
25. El-Bakri A, Hussein NM, Ibrahim ZA, Hasan H, AbuOdeh R. Intestinal Parasite Detection in Assorted Vegetables in the United Arab Emirates. *Oman MedJ*.2020; 35(3): e128.
26. Damen JG, Banwat EB, Egah DZ, Allanana JA. Parasitic contamination of vegetables in Jos, Nigeria. *Annals AfrMed*.2007;6(3), 115-118.
27. Adamu NB, Adamu JY, Mohammed D. Prevalence of helminth parasites found on vegetables sold in Maiduguri, Northeastern Nigeria. *Food Control*,2012;25:23-6.
28. de Silva NR, Brooker S, Hotez PJ, Montresor A, Engels D, Savioli L. Soil-transmitted helminthes infections: updating the global picture. *Trends Parasitol*. 2003; 19: 547–551.
29. Falavigna LM, Freitas CBR, Melo GC, Nishi L, Araújo SM, Falavigna G. Quality of green vegetables marketed in the northwest of Paraná, Brazil. *Parasitol Latinoam*,2005; 60(3-4):144-9.
30. Pereira WR, Kloos H, Crawford SB, Velásquez-Melendez JG, Matoso LF, Fujiwara RT. *Schistosomamansonii* infection in a rural area of the Jequitinhonha Valley, Minas Gerais, Brazil: analysis of exposure risk. *Acta Trop*. 2010; 113(1):34-41.
31. Guiherme ALF, Araújo SM, Falavigna DLM, Pupulim ART, Dias MLGG, Oliveira HS. Prevalence of intestinal parasites in horticulturists and vegetables from Feira do Produtor de Maringá, Paraná, Brazil. *Rev Soc Bras Med Trop*.1999; 32(4): 405-411.
32. Tengku SA, Norhayati M. Public health and clinical importance of amoebiasis in Malaysia: A review. *Trop Biomed*. 2011; 28(2): 194-22.