

Original Research Article

“Phytochemical and Biological Investigation of *Tridax procumbens* leaves.”

Abstract

Aims: The objective of this study is to search the biological activity such as antioxidant activity, hypoglycemic activity and antibacterial activity with initial phytochemical investigations of the leaves of *Tridax procumbens*.

Methodology: The phytochemical purposes were achieved by using two different chromatographic techniques thin layer chromatography (TLC), vacuum liquid chromatography (VLC). DPPH test, glucose uptake activity of yeast cells and agar plate diffusion technique were performed to evaluate the antioxidant capacity, hypoglycemic activity and anti-bacterial properties respectively of different fraction obtained from VLC by using different solvent systems such as dichloromethane, butanol, ethyl acetate and methanol. 31.25 µg/ml, 62.5 µg/ml, 125 µg/ml, 250 µg/ml and 500 µg/ml sample were used from each fraction to investigate antioxidant and hypoglycemic activity where vitamin C tablet and metformin were used as standard respectively. Antibacterial capacity was measured by measuring zone of inhibition resulted by the fractions where ciprofloxacin was taken as reference drug.

Results: Some visible spots were identified in the TLC plate which confirms the presence of different compounds in the fractions obtained from the methanolic extract of the leaves of *Tridax procumbens*. Butanol and ethyl acetate fractions showed the highest antidiabetic activity. In addition to, ethyl acetate, butanol and methanol fractions exerted more than 80% antioxidant capacity in DPPH assay. Furthermore, butanol fraction gives maximum zone of inhibition up to 21mm against *Staphylococcus aureus* which ensure anti-bacterial properties of the leaves of *Tridax procumbens* specially against *Staphylococcus aureus* and its strains.

Conclusion: Notwithstanding the all the research, the leaves had anti-inflammatory properties, which may have been generated by their antioxidant activity. As a result, if more therapeutic research is done, the separated compounds in those fractions may be employed as future therapeutic strategies.

Keywords: *Tridax procumbens*, Hypoglycemic, Anti-Diabetic, Anti-Oxidant, Anti-Bacterial, Phytochemistry.

1. Introduction:

Medicinal plants have been recognized and utilized throughout human history owing to their ability to produce a diverse number of chemical compounds that serve significant biological activities and have been used in health and medicinal purposes for thousands of years [1-3]. *Tridax procumbens* Linn. is commonly known as "Coatbutton" or "Tridhara Flower" or "Tridax Daisy" which is a member of the Asteraceae family [4-6]. It grows to a height of 30 to 50 cm and has hairy, short, blade-like leaves. Its thinly hairy, branched stem roots at the nodes [7, 8]. Traditionally its leaf is used as a drink to cure treat bronchial catarrh, diarrhea, dysentery, and liver diseases in many countries in Africa, South and Southeast Asia [6, 9-13]. Table 1 shows the taxonomic classification of *Tridax procumbens* [14, 15]. Earlier researchers reported presence of a number chemical compounds such as alkaloids, flavonoids, carotenoids, β -sitosterol, fumeric acid, luteolin, quercetin, oxoester, lauric acid, myristic, dexamethasone, glucoluteolin, palmitic, arachidic, linoleic acid and tannin etc. were separated & purified from the plant [16-18]. Linolenic acid and a new flavonoid like compound Procumbetin (3, 6- dimethoxy-5, 7, 2', 3', 4'- pentahydroxyflavone 7- O- β -glucopyranoside) has been reported and isolated from aerial parts of plant, also, a pair of water soluble polysaccharides was purified from the leaves [17, 19]. The leaves also contain high amount of minerals for example, calcium, magnesium, potassium, sodium, selenium, etc. and it serves as a good source of plant protein and potassium supplement, carotenoids to the local generation [19-21]. Furthermore, Many bioactive compounds, such as procumbetin [6], 8,3'-dihydroxy-3,7,4'-trimethoxy-6-O- β -d-glucopyranosyl flavone, 6,8,3'-trihydroxy-3,7,4'-trimethoxyflavone; puerarin [22], centaurein, centaureidin, phenolic acids [23] and various lipid constituents [16] such as methyl 14-oxooctadecanoate, methyl 14-oxononacosanoate, 30-methyl-28-oxodotriacont-29-en-1-oic acid, β -amyrone, β -amyrin; lupeol, and fucosterol have been successfully isolated from this plant [15, 24-28].

The aim of this research project was to carry out the characterization of the functional molecules present in the methanolic extract of leaves of the *Tridax procumbens* and investigate their biological activities.

Table 1: Taxonomic Classification of *Tridax procumbens* Linn.

Kingdom	: Plantae
Subkingdom	: Tracheobionta
Division	: Spermatophyta
Subdivision	: Magnoliophyta
Class	: Magnoliopsida
Subclass	: Asteridae
Order	: Asterales
Family	: Asteraceae
Genus	: <i>Tridax</i>
Species	: <i>procumbens</i>

2. Materials and Method:

2.1 Plant materials collection, processing, and phytochemical investigations:

The plants were collected from Jhalokathi district, under Barishal division of Bangladesh by uprooting the whole plant, then the leaves were removed carefully and voucher specimens have been deposited in Bangladesh National Herbarium (BNH) for future references. After one week of shade drying, the leaves were grinded to coarse powder and successively extracted with methanol by maceration technique. After subsequent evaporation of solvents 22.5 g methanolic extract of *Tridax procumbens* leaves was obtained. Vacuum liquid chromatography (VLC) was applied collect different fractions in Dichloromethane (CHCl₂), Butanol (C₄H₉OH), Ethyl Acetate (C₄H₈O₂) and Methanol (CH₃OH) from the crude methanolic extract and marked the fractions as DCMF, BF, EAF & MF respectively. After concentrating the fractions by using rotary evaporator, they were treated with thin layer liquid chromatography. The most active fractions having the similar thin layer chromatography profile were pooled together which were conducted by three solvent systems such as non-polar (Benzene : Ethanol : Aluminum hydroxide = 9 : 1 : 0.1), intermediate polar (Chloroform : Ethyl acetate : Formic acid = 5 : 4 : 1), and polar (Ethyl acetate : Ethanol : Water = 8 : 1.2 : 0.8). After numerous trial and error, non-polar solvent was fixed as mobile phase of thin layer liquid chromatography.

2.2 Anti-oxidant activity test:

The antioxidant properties of the fractions investigated by 2,2-Diphenyl-1-picrylhydrazyl (DPPH) which was previously describe as the most popular, quickest and easiest approach for the measurement of antioxidant properties [29]. At first, 3.125 mg/ml sample solution was prepared from each fraction with methanol, and 1ml, 2ml, 4ml, 8ml and 16ml of sample solution from each fraction were transferred into five different volumetric flasks and diluted up to 10 ml with methanol. 3 ml of a 0.0004% DPPH solution was added to 1 ml of the solution, which was then diluted with methanol to make it up to 10 ml. After keeping them for 30 minutes in a dark room, the absorbance was measured at 517 nm in a UV spectrophotometer. The standard was a vitamin C tablet (250 mg), and the blanks were methanol and a DPPH mixture. Antioxidant activity was measured using the following equation:

$$\text{Antioxidant capacity (\%)} = \left(1 - \frac{\text{Abs. of sample} - \text{Abs. of blank}}{\text{Abs. of control}}\right) \times 100$$

2.3 In vitro hypoglycemic activity test:

Antidiabetic activity was investigated by the previously described method of glucose uptake in yeast cells [30-32]. A 50% (v/v) suspension of commercial baker's yeast was made in distilled water after being repeatedly centrifuged (3,000rpm; 5 min) until the supernatant fluids were clear. 1 ml of glucose solution (10 mM) and various quantities of isolated components (1 mg) were combined, and the mixture was then incubated at 37 °C for 10 minutes. The reaction was initiated by adding 100µl of yeast suspension and incubated for further 60 min at 37 °C and then they were centrifuged at 2,500rpm for 5 min to determine the amount of glucose in the

supernatant. Metformin was used as standard and a mixture of all of the chemicals, excluding the test sample were used as control. Glucose uptake was measured by using the following formula:

$$\text{Glucose Uptake (\%)} = \left(\frac{\text{Abs. of sample} - \text{Abs. of control}}{\text{Abs. of sample}} \right) \times 100$$

2.4 Antimicrobial screening test:

The agar diffusion assay is one of the established method for quantifying the ability of antibiotics to inhibit bacterial growth [33, 34]. The tested bacterium is inoculated on an agar plate first, and then an antimicrobial agent is transferred by diffusion resulting growth inhibition zones [34]. About 100µl of each fraction was added to previously incubated bacterial inoculum containing agar plates. After 30 min refrigeration, and incubation at 37°C for 18 hours, the antimicrobial activity was measured by measuring the zone of inhibition appeared after the incubation period. 10% Methanol was used as a negative control and ciprofloxacin (30µg/disk) was used as the reference standard.

3. Results:

3.1 Phytochemical investigations:

After charring of the TLC plate with 10% sulfuric acid solution and DPPH solution some visible spots were observed.

Table 2: R_f value of different fractions

Name of sample	1st spot R _f value	2nd spot R _f value	3rd spot R _f value
Crude Extract	0.31	0.50	0.75
DCMF	0.53	0.77	0.89
BF	0.55	0.75	0.96
EAF	0.60	0.87	0.93
MF	0.2	0.5	No spot

3.2 Antioxidant activity test:

The illustrations of antioxidant test of different fraction of methanolic extract of the *Tridax procumbens* leaves are given below:

Table 3: Antioxidant activity test results

Concentration (µg/ml)	Antioxidant Capacity (%)				
	DCMF	BF	EAF	MF	Metformin
31.25	12.61	19.80	62.23	57.56	58.89
62.5	13.28	42.52	72.43	70.25	60.73
125	89.72	47.86	80.78	79.28	63.40
250	40.68	51.89	83.62	81.95	66.75
500	70.25	55.89	84.62	83.29	71.90

3.3 Anti-bacterial activity test:

The result of antibiotic test of different fraction of methanolic extract of the *Tridax procumbens* leaves are given below:

Table 4: Anti-bacterial activity test results

Tested bacteria	Zone of inhibition (mm)					
	MF	EAF	DCF	BF	STD	Control
<i>S. paratyphi</i>	9	9	8	14	24	0
<i>Bacillus sereus</i>	8	9	7	7	23	0
<i>Bacillus subtilis</i>	13	11	8	9	26	0
<i>Staphylococcus aureus</i>	13	12	9	21	30	0
<i>Salmonella typhi</i>	9	8	8	11	31	0
<i>Shigella dysenteriae</i>	10	8	10	7	27	0
<i>Vibrio mimicus</i>	9	9	8	14	28	0
<i>Candida albicans</i>	9	10	9	11	27	0
<i>Aspergillus niger</i>	9	10	9	11	27	0
<i>E.coli</i>	10	11	9	11	25	0
<i>Vibrio parahemolyticus</i>	11	9	8	13	30	0
<i>Bacillus megaterium</i>	13	11	8	9	27	0
<i>Pseudomonas aureaus</i>	11	12	9	12	22	0

3.4 Hypoglycemic test:

The result of hypoglycemic test of different fraction of methanolic extract of the *Tridax procumbens* leaves are given below:

Table 5: Hypoglycemic activity test results

Concentration ($\mu\text{g/ml}$)	Glucose uptake capacity (%)				
	DCMF	BF	EAF	MF	Metformin
31.25	7.68	12.07	13.83	12.07	22.37
62.5	15.25	27.39	20.46	21.9	28.19
125	26.56	52.13	39.01	22.89	33.2
250	33.72	60.41	45.12	29.95	35.74
500	46.03	62.75	58.44	51.22	39.74

4. Discussion:

TLC plates were developed with dichloromethane (DCF), butanol (BF), ethyl acetate (EAF), methanol (MF) fractions by using non polar to most polar solvent system and the best result was found in non-polar solvent system. After charring of the sulfuric acid and DPPH solution on the TLC plate, the visible spots indicate the presence of different compounds in the methanolic extract of the leaves of *Tridax procumbens*. Further extractions and purifications from these crude drugs may lead to the possible isolation of these compounds from the crude extracts.

In the Glucose uptake in Yeast cells method the mechanism of glucose transport across the yeast cell membrane has been receiving attention as in vitro screening method for hypoglycemic effect of various compounds. It is reported that, in yeast cells (*Saccharomyces cerevisiae*) glucose is transported by a facilitated diffusion process [35]. In our result, It has shown that, butanol fraction (BF) and ethyl acetate fraction (EAF) of methanolic extract give higher antidiabetic activity 62% and 58% respectively. Whereas, methanol fraction (MF) and dichloromethane fraction (DCF) of methanolic extract of give 51% and 46% respectively.

The reduction in DPPH radical which can accept an electron of hydrogen radical was determined by the decrease of its absorbance at 517 nm (in methanol) confirmed antioxidant capacity of the samples [36]. In our result, it has shown that the fractions of dichloromethane (DCF), butanol (BF), ethyl acetate (EAF) and methanol (MF) give respectively 89%, 55%, 84%, and 83% antioxidant activity.

Various strains of gram positive and gram negative bacteria were used as the test microorganism to determine antibacterial capacity of the fractions where the zones of inhibition for the microbes were measured in millimeter by using a transparent ruler. The fraction obtained through butanol (BF) gives zone of inhibition up to 21mm against *Staphylococcus aureus* which is maximum than others and it indicates that, the leaves of *Tridax procumbens* possessing antimicrobial activity can be employed against human pathogens specially for *Staphylococcus aureus* and its strains.

5. Conclusion:

The present study on the different fraction of methanolic extract of the *Tridax procumbens* leaves showed the potentiality as an antioxidant, a hypoglycemic agent and antibacterial agent. Besides, the leaves showed anti-inflammatory activity which may be induced due to its antioxidant activity. So, the isolated compounds in those fractions may be used as future therapeutic tools if further therapeutic investigations are carried out.

6. References:

1. Samy RP, Pushparaj PN, Gopalakrishnakone P. A compilation of bioactive compounds from Ayurveda. *Bioinformation*. 2008;3(3):100.
2. Tapsell LC, Hemphill I, Cobiac L, Sullivan DR, Fenech M, Patch CS, et al. Health benefits of herbs and spices: the past, the present, the future. 2006.
3. Saha RK, Nandi T, Nahar K, Talukdar AA, Nandi PC. Phytochemical and biological investigation of VLC fraction of *Senna alata* leaves. *Journal of Pharmacognosy and Phytochemistry*. 2016;5(3):55.
4. Saxena M, Mir AH, Sharma M, Malla MY, Qureshi S, Mir MI, et al. Phytochemical screening and in-vitro antioxidant activity isolated bioactive compounds from *Tridax procumbens* Linn. *Pakistan Journal of Biological Sciences: PJBS*. 2013;16(24):1971-7.

5. Sawant RS, Godghate AG. Preliminary phytochemical analysis of leaves of *Tridax procumbens* Linn. *International Journal of Science, Environment and Technology*. 2013;2(3):388-94.
6. Ali M, Ravinder E, Ramachandram R. A new flavonoid from the aerial parts of *Tridax procumbens*. *Fitoterapia*. 2001;72(3):313-5. doi: 10.1016/s0367-326x(00)00296-3. PubMed PMID: 11295316.
7. Bhagwat DA, Killedar SG, Adnaik RS. Anti-diabetic activity of leaf extract of *Tridax procumbens*. *International Journal of Green Pharmacy (IJGP)*. 2008;2(2).
8. Chauhan BS, Johnson DE. Germination ecology of two troublesome Asteraceae species of rainfed rice: Siam weed (*Chromolaena odorata*) and coat buttons (*Tridax procumbens*). *Weed Science*. 2008;56(4):567-73.
9. Berger I, Barrientos AC, Caceres A, Hernandez M, Rastrelli L, Passreiter CM, et al. Plants used in Guatemala for the treatment of protozoal infections: II. Activity of extracts and fractions of five Guatemalan plants against *Trypanosoma cruzi*. *J Ethnopharmacol*. 1998;62(2):107-15. doi: 10.1016/s0378-8741(98)00011-7. PubMed PMID: 9741882.
10. Kuldeep G, Pathak A. Pharmacognostic and Phytochemical Evaluation of *Tridax procumbens* Linn. *Journal of Pharmacognosy and phytochemistry*. 2013;1(5).
11. Kpodar MS, Karou SD, Katawa G, Anani K, Gbekley HE, Adjrah Y, et al. An ethnobotanical study of plants used to treat liver diseases in the Maritime region of Togo. *J Ethnopharmacol*. 2016;181:263-73. Epub 20151228. doi: 10.1016/j.jep.2015.12.051. PubMed PMID: 26732630.
12. Ravikumar V, Shivashangari KS, Devaki T. Hepatoprotective activity of *Tridax procumbens* against d-galactosamine/lipopolysaccharide-induced hepatitis in rats. *J Ethnopharmacol*. 2005;101(1-3):55-60. doi: 10.1016/j.jep.2005.03.019. PubMed PMID: 15923095.
13. Safina K, Singh D. A COMPREHENSIVE REVIEW OF A HEALING HERB: TRIDAX PROCUMBENS LINN. *International Journal of Ayurveda and Pharma Research*. 2017.
14. Ghosh P, Biswas S, Biswas M, Dutta A, Sil S, Chatterjee S. Morphological, Ethnobiological and Phytopharmacological Attributes of *Tridax procumbens* Linn. (Asteraceae): A Review. *Int J Sci Res in Biological Sciences Vol*. 2019;6(2).
15. Bhagat V, Kondawar M. A comprehensive review on phytochemistry and pharmacological use of *Tridax procumbens* Linn. *Journal of Pharmacognosy and Phytochemistry*. 2019;8(4):01-10.
16. Verma RK, Gupta MM. Lipid constituents of *Tridax procumbens*. *Phytochemistry*. 1988;27(2):459-63.

17. Singh K, Ahirwar V. Acute and chronic toxicity study of *Tridax procumbens* on haemoglobin percent and blood sugar level of Sprague Dawley rats. *Journal of Pharmacology and Toxicology*. 2010;1(1):1-6.
18. Subramanian SS, Ramakrishnan S, Nair A. Isolation of luteolin and glucoluteolin from the flowers of *Tridax Procumbens*. *Current Science*. 1968;37(16):465-.
19. Raju TS, Davidson E. Structural features of water-soluble novel polysaccharide components from the leaves of *Tridax procumbens* Linn. *Carbohydrate research*. 1994;258:243-54.
20. Chen W-H, Ma X-M, Wu Q-X, Shi Y-P. Chemical-constituent diversity of *Tridax procumbens*. *Canadian Journal of Chemistry*. 2008;86(9):892-8.
21. Ikewuchi C J, Ikewuchi C C, Ngozi M. Chemical profile of *Tridax procumbens* Linn. *Pakistan journal of nutrition*. 2009;8(5):548-50.
22. Xu R, Zhang J, Yuan K. Two new flavones from *Tridax procumbens* Linn. *Molecules*. 2010;15(9):6357-64. Epub 20100909. doi: 10.3390/molecules15096357. PubMed PMID: 20877227; PubMed Central PMCID: PMC6257746.
23. Andriana Y, Xuan T, Quan N, Quy T. Allelopathic potential of *Tridax procumbens* L. on radish and identification of allelochemicals. *Allelopath J*. 2018;43:223-38.
24. Ahmed SS, Prakash KC, Tabassum S, Salma N, Devi K. Pharmacognostical and Pharmacological Review on *Tridax procumbens* Linn. *Research Journal of Pharmacology and Pharmacodynamics*. 2019;11(1):11-6.
25. Beck S, Mathison H, Todorov T, Calder E, Kopp OR. A review of medicinal uses and pharmacological activities of *Tridax procumbens* (L.). *J Plant Stud*. 2018;10.
26. Thalkari AB, Karwa PN, Shinde PS, Gawli CS, Chopane PS. Pharmacological actions of *Tridax procumbens* L.: A Scientific Review. *Research Journal of Pharmacognosy and Phytochemistry*. 2020;12(1):27-30.
27. Kumar S, Prasad A, Iyer S, Vaidya S. Pharmacognostical, phytochemical and pharmacological review on *Tridax procumbens* Linn. *International Journal of Pharmaceutical & Biological Archives*. 2012;3(4):747-51.
28. Jachak SM, Gautam R, Selvam C, Madhan H, Srivastava A, Khan T. Anti-inflammatory, cyclooxygenase inhibitory and antioxidant activities of standardized extracts of *Tridax procumbens* L. *Fitoterapia*. 2011;82(2):173-7. Epub 20100908. doi: 10.1016/j.fitote.2010.08.016. PubMed PMID: 20804828.
29. Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol*.

2007;39(1):44-84. Epub 20060804. doi: 10.1016/j.biocel.2006.07.001. PubMed PMID: 16978905.

30. Nayan M, Hossain I, Alam M, Jamil M, Hossain M, Haq I, et al. Pharmacological effect of *Heritiera fomes* on Long Evans rats against Postprandial Hyperglycemia and $\left(\text{C} \right)_{\text{H}} - \text{O} - \text{n}$ adsorption in vitro. *Journal of Diabetes & Metabolic Disorders*. 2022:1-9.

31. Kotowaroo MI, Mahomoodally MF, Gurib-Fakim A, Subratty AH. Screening of traditional antidiabetic medicinal plants of Mauritius for possible alpha-amylase inhibitory effects in vitro. *Phytother Res*. 2006;20(3):228-31. doi: 10.1002/ptr.1839. PubMed PMID: 16521114.

32. Rehman G, Hamayun M, Iqbal A, Ul Islam S, Arshad S, Zaman K, et al. In Vitro Antidiabetic Effects and Antioxidant Potential of *Cassia nemophila* Pods. *Biomed Res Int*. 2018;2018:1824790. Epub 20180123. doi: 10.1155/2018/1824790. PubMed PMID: 29607313; PubMed Central PMCID: PMC5827890.

33. Daoud A, Malika D, Bakari S, Hfaiedh N, Mnafigui K, Kadri A, et al. Assessment of polyphenol composition, antioxidant and antimicrobial properties of various extracts of Date Palm Pollen (DPP) from two Tunisian cultivars. *Arabian Journal of Chemistry*. 2019;12(8):3075-86.

34. Bonev B, Hooper J, Parisot J. Principles of assessing bacterial susceptibility to antibiotics using the agar diffusion method. *Journal of antimicrobial chemotherapy*. 2008;61(6):1295-301.

35. Lagunas R. Sugar transport in *Saccharomyces cerevisiae*. *FEMS Microbiol Rev*. 1993;10(3-4):229-42. doi: 10.1016/0378-1097(93)90598-v. PubMed PMID: 8318258.

36. Kedare SB, Singh RP. Genesis and development of DPPH method of antioxidant assay. *J Food Sci Technol*. 2011;48(4):412-22. Epub 20110225. doi: 10.1007/s13197-011-0251-1. PubMed PMID: 23572765; PubMed Central PMCID: PMC3551182.