

Original Research Article

Prevalence and Perception of Herbal Medicine Use Among Patients with Chronic Diseases in a Nigerian Teaching Hospital

ABSTRACT

Background: Man's quest to manage illness over many years has led to the development of complementary and alternative medicine (CAM). There are many types of CAM, and the most utilized is herbal medicine. The relevance of herbal medicine has increased over the years due to the surge in health demands and other factors that influence the management of diseases. Hence, this study was conducted to ascertain the frequency and pattern of herbal medicine use among patients diagnosed with chronic diseases in the University of Nigeria Teaching Hospital, its predictors, magnitude, and perception.

Main body of abstract

Methodology: We conducted a cross-sectional descriptive study among adult patients ($n=319$) diagnosed with various chronic diseases, such as Diabetes Mellitus, Hypertension, Glaucoma, cancer, etc., from October 2022 to November 2022, at the University of Nigeria Teaching Hospital, Enugu, Nigeria. Questionnaires were administered through face-to-face interviews with patients.

Result: A total of 319 patients were included in this study. The gender distribution is 130 males and 180 females (?). The rate of herbal medicine use was 65.2%. Herbal medicine use among the female gender was higher, but there was no relationship between gender and herbal medicine use ($P = 0.842$). Of participants using herbal medicine, 46.1% used it for treating their health condition, 30.4% used herbal medicine to improve well-being, and 10.3% for preventing diseases. Their perception of herbal medicine uses as compared to conventional synthetic medicine based on modifications of some issues found from literature reviews that militate against the use of herbal medicine such as dose (15% strongly agreed, 20.1% agreed, 23.2% neutral, 18.8% disagreed and 22.9% strongly disagreed) etc.

Short conclusion: The study discovered that patients in Nigeria with chronic conditions frequently (used) use herbal medicines. Age, gender, marital status, occupation, education, monthly income, religion, health insurance, the timing of sickness diagnosis, the number of medications prescribed, and multiple chronic conditions were all linked to the usage of herbal medicines.

KeyWords:Chronicdisease,herbalmedicines,perception,TeachingHospital.

UNDER PEER REVIEW

INTRODUCTION

Since man's existence, he has always sought ways to improve his health and prevent and treat diseases. This has led to the development of complementary and alternative medicine (CAM). There are many types of CAM, and the most utilized is herbal medicine. Herbal medicine is one of the most balanced and moderate approaches to achieving health [1]. Studies have shown that herbal medicine is beneficial and is usually consumed with orthodox medicine, notwithstanding the challenges [2]. The relevance of herbal medicine has increased over the years due to the surge in health demands and other factors that influence the management of diseases. Using herbal medicine has become a global trend, with Nigeria following duly. Although patients perceive herbal medicine to be safe because it is of natural origin and best suited for the long-term management of diseases [3, 4], there are issues such as inappropriate use and self-medication, obscure risk awareness, non-specialist consultation which could potentially lead to drug interactions and toxicity, especially in patients who are on many drugs [5]. A comparison of orthodox and herbal medicine in Nigeria showed that herbal medicine was better in terms of safety, accessibility, advertisement, cost, and safety [5]. Over the years, it has been shown that herbal medicines exert the following pharmacological effects: antidiabetic, anti-depressive, analgesic, anti-psoriasis, anticancer, hepatoprotective, Etc. [6]. Complementary and alternative medicine use to treat chronic disease has increased globally, even in Nigeria [7-11]. Most of the global population relies on herbal medicine, especially in developing countries (80%) [12-15]. Several studies have revealed H.M. use among health subpopulations in diverse settings and locations [16-21]. Much research has been carried out in Nigeria on herbal medicine use. However, studies of its utilization in patients diagnosed with chronic diseases in tertiary hospitals are scarce and only addressed selected chronic diseases without giving holistic knowledge [22-24]. These patients have the highest need for herbal medicine due to the perceived minimal toxicity in long-term use and the slow onset of action, and it would be of little use in acute conditions. Hence, this survey was conducted to ascertain the frequency and usage of herbal medicine among patients who have been diagnosed with chronic diseases in the University of Nigeria Teaching Hospital and to assess the factors that influence its usage and perceptions among patients to increase public health campaigns as well as inform physician and pharmacist counseling and to enlighten and counter the misconceptions associated with herbal medicine usage [24].

AIMS AND OBJECTIVES?

METHODS

Between October 2022 and November 2022, a cross-sectional descriptive study was carried out among adult patients ($n=319$) who had been given a diagnosis of various chronic diseases, including diabetes, Hypertension, glaucoma, cancer, etc., at the University of Nigeria Teaching Hospital in Enugu, Nigeria. The study only included eligible patients who provided their informed permission. All authors delivered a face-to-face questionnaire to the participants [25]. Cronbach's alpha was used to pre-test the questionnaire's reliability and validity on a sample of 20 patients who did not make up the final sample. Each participant

UNDER PEER REVIEW

provided their written informed consent, and the respondents' privacy and confidentiality were rigorously maintained. The "Committee of Research Ethics of the University of Nigeria Teaching Hospital (Ref: UNTH/HREC/2022/08/452)" accepted the study protocol. We adhered to the World Medical Association's Declaration of Helsinki concerning the moral treatment of human beings in research [26].

Study population

Sample size and recruitment strategy

The sample size was calculated according to the equation of Slovin's formula =

$$\frac{N}{1 + N \left(e^2 \right)}$$

n = sample size, N = target population, e = margin of error (0.05)

The value of N (1576), the average target population of chronic disease inpatients and outpatients in a month, was obtained from the UNTH database. Substituting target population of 1576.

$$n = \frac{1576}{1 + 1576 \left(0.05^2 \right)}$$

$$= 319 \text{ patients}$$

The sample size in this study is approximately 319 patients.

Through a systematic random sampling process, patients were chosen. The authors collected in-person data using an 18-item questionnaire that was developed and tailored from previously validated studies on herbal medicine [25-27].

The questionnaire

included information on the Patient's sociodemographic variables, disease features, diagnosis time, and the number of drugs the Patient took. The questionnaires also assess the use of herbal medicines in patients, including what it was used for, the reason for its use, the source of information of these patients, reasons for this choice as against conventional synthetic drugs, and the Patient's perception of the issues raised about herbal medicines.

Inclusion and Exclusion Criteria

All will meet the eligibility criteria for participation in this study:

1. Adult patients with chronic diseases aged 18 and above.

2. Patients with or without co-morbidities.
3. Who can read and write in the English language
4. Who agreed to participate.

Data Analysis

The (study's) data were examined using the Statistical Package for Social Sciences version 15 for Windows. Mean, standard deviation, range, and frequency (% values) are the formats used to report descriptive statistics. Categorical variables were subjected to the chi-square test or Fisher's exact test. Statistical significance is defined as $P < 0.05$.

RESULTS

A total of 319 patients participated in this study as shown in **table 1**. There were 130 male (43.6%) and 180 female (56.4%) patients (?). Whereas 38.2% of the patients are employed, 26.3% are unemployed, and 35.4% are self-employed. In **table 2**, there were 77 patients (24.1%) with hypertension, 33 patients (10.3%) with cancer, 32 patients (10.0%) with diabetes mellitus, 29 patients (9.1%) with chronic kidney disease, 13 patients (4.1%) with stroke and peptic ulcer disease each, followed by arthritis and cataract which were suffered by 12 (3.8%) and 11 (3.4%) patients respectively. There were 116 patients (53.5%) with more than one disease. Among those patients, 319 (100.0%) used conventional medicine, and 209 (65.2%) also used herbal medicine. The sociodemographic and clinical features of the study population are presented in **Table 1**. The number of herbal medicine users was 208. Herbal medicine use was found to have no significant association with the study population's social demographic and clinical features, as (seen) shown in **table 1**.

From **table 5**, Herbal medicines were commonly recommended to the users by their families or friends (60.5%), by radio/T.V. advertisement (29.2%), by medicine experts (22.6%), by internet (14.4%), by books/magazines (14.1%), and by seminar/workshop (8.5%) respectively. Of participants using herbal medicine, 46.1% used it to treat their health condition, 30.4% used it to improve well-being, and 10.3% to prevent diseases as shown in **table 4**. There was also found to be a significant association between the indications for herbal medicine use (improve health, prevent disease, and treat disease) and herbal medicine use ($p=0.000$) as (seen) shown in **table 3**.

As can be seen in **table 6**, The reasons for herbal medicine use were to be due to belief in efficacy in the past (32.0%), cost (20.4%), ease of accessibility (19.7%), it has always been done this way/people's opinion (19.4%), family tradition/culture (17.6%), decreased side effect (17.6%) and dissatisfaction with herbal medicine (14.4%).

From **Table 7**, their perception of herbal medicine uses as compared to conventional synthetic medicine based on modifications of some issues found from literature reviews that militate against the use of herbal medicine such as dose (15% strongly agreed, 20.1% agreed, 23.2% neutral, 18.8% disagreed and 22.9% strongly disagreed), taste (7.8% strongly agreed, 14.4% agreed, 29.5% neutral, 21.6% disagreed and 26.6% strongly disagreed), smell (8.2% strongly agreed, 13.8% agreed, 30.1% neutral, 21.0% disagreed and 27.0% strongly disagreed), safety (13.8% strongly agreed, 18.5% agreed, 33.2% neutral, 12.9% disagreed and 21.6% strongly disagreed), NAFDAC regulation (12.9% strongly

agreed, 12.2% agreed, 32.0% neutral, 18.5% disagreed and 24.5% strongly disagreed) and the dressing/organization of the sellers/dispensers

(3.1% strongly agreed, 10.7% agreed, 36.7% neutral, 21.6% disagreed and 27.9% strongly disagreed).

Table 1: Socio-Demographic and Clinical Characteristics of the Study Population

Patients' Characteristics	Sample	Herbal Medicine Use		P-Value
	All	Yes	No	
Participants	N(%)	N(%)	N(%)	
<i>Age (years, mean \pmSD)</i>				0.096
18-28	52(16.3)	25(7.8)	27(8.5)	
29-39	54(16.9)	33(10.3)	21(6.6)	
40-50	58(18.2)	39(12.2)	19(6.0)	
51-61	71(22.3)	51(16.0)	20(6.3)	
62 and above	84(26.3)	52(16.3)	32(10.0)	
<i>Gender</i>				0.842
Male	139(43.6)	88(27.6)	51(16.0)	
Female	180(56.4)	112(35.1)	68(21.3)	
<i>Educational level</i>				0.295
Primary	44(13.8)	30(9.4)	14(4.4)	
Secondary	93(29.2)	63(19.7)	30(9.4)	
Tertiary	174(54.5)	101(31.7)	73(22.9)	
None	8(2.5)	6(1.9)	2(0.6)	
<i>Marital Status</i>				0.164
Married	213(66.8)	142(44.5)	71(22.3)	
Single	77(24.1)	35(11.0)	42(13.2)	
Divorced	4(1.3)	3(0.9)	1(0.3)	
Widowed	25(7.8)	13(4.1)	12(3.8)	
<i>Occupational status</i>				0.407

EmployedUn	122(38.2)	71(22.3)	51(16.0)
employed	84(26.3)	54(16.9)	30(9.4)

UNDER PEER REVIEW

Selfemployed	113(35.4)	75(23.5)	38(11.9)	
<i>Monthlyincome(naira)</i>				0.382
<50,000	159(49.8)	105(32.9)	54(16.9)	
50,000-99,000	85(26.6)	50(15.7)	35(11.0)	
100,000-199,000	57(17.9)	37(11.6)	20(6.3)	
200,000-299,000	10(3.1)	4(1.3)	6(1.9)	
>300,000	8(2.5)	4(1.3)	4(1.9)	
<i>ReligionChristia</i>				0.180
nityIslamTrad	306(95.9)	189(59.2)	117(36.7)	
itionalistOther	7(2.2)	7(2.2)	0(0.0)	
s	5(1.6)	3(0.9)	2(0.6)	
	1(0.3)	1(0.3)	0(0.0)	
<i>Healthinsurance</i>				0.489
Insured	43(13.5)	29(9.1)	14(4.4)	
NotInsured	276(86.5)	171(53.6)	105(32.9)	
<i>When illness wasdiagnosed(years)</i>				0.282
1-5	223(69.9)	146(45.8)	77(24.1)	
6-10	51(16.0)	30(9.4)	21(6.6)	
11-15	20(6.3)	9(2.8)	11(3.4)	
More than15	25(7.8)	15(4.7)	10(3.1)	
<i>Number of drugsprescri bed for illness</i>				0.065
	180(56.4)	103(32.3)	77(24.1)	
1-4	99(31.0)	68(21.3)	31(9.7)	
5-8	40(12.5)	29(9.1)	11(3.4)	
More than8				

* $P < 0.05$ considered as statistically significant, SD: Standard deviation

Table 2: Associations Between Herbal Medicine Use and Chronic Diseases

Chronic Disease	Sample	Herbal Medicine Use	P-Value
	N(%)	N(%)	
Acne Vulgaris	1(0.3)	1(0.3)	0.126
Allergy	1(0.3)	1(0.3)	
Arthritis	12(3.8)	9(2.8)	
Asthma	2(0.6)	1(0.3)	
Atherosclerosis	1(0.3)	1(0.3)	
Bipolar Disorder	1(0.3)	1(0.3)	
Cancer	33(10.3)	19(6.0)	
Cardiomegaly	2(0.6)	2(0.6)	
Cataract	11(3.4)	6(1.9)	
Chronic Ankle Fracture	1(0.3)	0(0.0)	
Chronic Chest Pain	2(0.6)	1(0.3)	
Chronic Ear Problem	3(0.9)	2(0.6)	
Chronic Eye Problem	8(2.5)	3(0.9)	
Chronic fatigue syndrome	1(0.3)	1(0.3)	
Chronic Infection	3(0.9)	3(0.9)	
Chronic Kidney disease	29(9.1)	25(7.8)	
Chronic Malaria	8(2.5)	6(1.9)	
Chronic Otitis Externa	4(1.3)	2(0.6)	
Chronic Stomachache	1(0.3)	1(0.3)	
Chronic Waist Pain	1(0.3)	1(0.3)	
Cirrhosis	1(0.3)	1(0.3)	
Coronary Artery Disease	1(0.3)	1(0.3)	

DiabetesMellitus	32(10.0)	21(6.6)
DryEyes	2(0.6)	2(0.6)
Endometriosis	1(0.3)	1(0.3)
EyeTrauma	1(0.3)	1(0.3)
EyeUlcer	1(0.3)	1(0.3)
Fibroid	4(1.3)	3(0.9)
Gastroesophageal Disease	1(0.3)	0(0.0)
Reflux		
Glaucoma	5(1.5)	4(1.3)
HearingLoss	1(0.3)	0(0.0)
Heart attack	1(0.3)	0(0.0)
HepatitisB	3(0.9)	1(0.3)
Hernia	1(0.3)	1(0.3)
Hypertension	77(24.1)	37(11.6)
Incisionalhernia	1(0.3)	1(0.3)
Infertility	1(0.3)	1(0.3)
Mentaldisorder	5(1.6)	1(0.3)
Motordisorder	1(0.3)	0(0.0)
Myopia	2(0.6)	1(0.3)
Nephropathy	2(0.6)	2(0.6)
OcularHypertension	1(0.3)	0(0.0)
Otomycosis	1(0.3)	0(0.0)
Parkinson'sdisease	2(0.6)	0(0.0)
Paroxysmal Dyspnea	1(0.3)	1(0.3)
Nocturnal		
PelvicInflammatoryDisease	2(0.6)	1(0.3)
PepticUlcerDisease	13(4.1)	12(3.8)

PituitaryAdenoma	1(0.3)	0(0.0)
ProstrateHyperplasia	6(1.9)	3(0.9)
Psoriasis	1(0.3)	0(0.0)
SkinDisease	1(0.3)	0(0.0)
SpinalcordInjury	1(0.3)	1(0.3)
Stroke	13(4.1)	10(3.1)
Tuberculosis	1(0.3)	1(0.3)
Tumor	1(0.3)	1(0.3)
ValvularHeartDisease	1(0.3)	1(0.3)

Abbreviations:AOR,AdjustedOddsRatio;CI,ConfidenceInterval

Table3:IndicationsforHerbalMedicineUse

Number of reasons	N(%)	p-value
All three reasons	20(6.3)	0.000
Improve my health and prevent disease	33(10.3)	
Prevent disease and treat my disease		
Only one reason	151(47.3)	
Didn't use for any reason	115(36.1)	

Table4:PurposeofHerbalMedicineUse

Purpose	N(%)
Improve my health	97(30.4)
Prevent disease	33(10.3)
Treat my disease	147(46.1)

Table5: Sources of Information on Herbal Medicine

Source of Information	N(%)
Books/Magazine	45(14.1%)
Family/Friends	193(60.5)
Medicine Experts	72(22.6)
Radio/TV advertisement	93(29.2)
Seminar/workshop	27(8.5)
Internet	46(14.4)

Table6: Reasons for Using Herbal Medicine

Reasons for Using Herbal Medicine	N(%)
I am not satisfied with western medicine	46(14.4)
It has worked for me in the past	102(32.0)
Family tradition or culture	56(17.6)
It has always been done this way/people's opinion	62(19.4)
It is cheaper than western medicine	65(20.4)
Herbal medicine is easier to get than western medicine	63(19.7)
Herbal medicine has fewer side effects than western medicine	56(17.6)

Table7: PreferencesifSomeoftheDrawbacksofHerbalMedicinesareResolved

Would preferher balmedicine towesternmedicine, if?	StronglyAgr ee	Agree	Neutral	Disagree	StronglyDis agree
	N(%)	N(%)	N(%)	N(%)	N(%)
The dose isstandardi zed	48(15.0)	64(20.1)	74(23.2)	60(18.8)	73(22.9)
thetasteismasked	25(7.8)	46(14.4)	94(29.5)	69(21.6)	85(26.6)
theodorismasked	26(8.2)	44(13.8)	96(30.1)	67(21.0)	86(27.0)
Safety ascertained is	44(13.8)	59(18.5)	106(33.2)	41(12.9)	69(21.6)
Regulated by NAFDAC	41(12.9)	39(12.2)	102(32.0)	59(18.5)	78(24.5)
Seller is welldresse d	10(3.1)	34(10.7)	117(36.7)	69(21.6)	89(27.9)

DISCUSSION

There is an increasing prevalence of chronic diseases in the world, hence an increase in the deaths caused but these diseases ranging from cardiovascular diseases, diabetes, etc. With 17.9 million fatalities yearly, cardiovascular disease ranks first among chronic illnesses, followed by cancer (9.3 million), chronic respiratory conditions (4.1 million), and diabetes (2.0 million, including diabetic kidney disease). Over 80% of all chronic disease-related premature deaths are attributed to these four disease groups. In Nigeria, research conducted by the WHO showed that chronic disease prevalence stood at 29% in 2018, cardiovascular diseases at 11%, cancer at 4%, and diabetes at 2%. In as much as chronic diseases remain the major cause of death in Nigeria, the country is plagued with the increasing burden of chronic diseases, with premature mortality of approximately 22%.

There is a current surge in the use of herbal medicine around the world, and the WHO says that more than 75% of people utilize it as their primary form of healthcare [28]. Tulunay *et al.* [25] reported the rate of herbal medicine used to be 29% in a similar participant with chronic diseases, while according to Peltzer *et al.*, 35.9% of people utilize herbal medications [29]. However, conducted in Thailand on the same study population. This study found that herbal medicine use was 65.2%. This contrast could be due to the sociodemographic factors of the two countries (even though the study population is the same), such as the monthly income of the population, occupational status, etc.

Moreover, conventional medication use was lower in patients receiving herbal medicine [25]. (Our) The present study showed that as much as patients were on herbal medicine medication, it did not affect their use of conventional medicine. Patients are still skeptical to accept herbal medicine wholly even if some of the issues outlined with it are outlined, as can be seen in Table 5. This provided an insight into the fact that they combined both, which could lead to a possibility of interaction (ranging from inhibition to potentiation) which affects the health outcome and thus should be a major concern for our health facilities to enlighten these patients and initiate public health campaigns to tackle this rather disturbing trend. Peltzer *et al.* reported that the prevalence of herbal medicine used to be 35.9%, among which 53.7% use it for long-term health issues, 40% use it for improving their health, and 6.3% use it for treating acute illnesses [29]. Comparing with the outcome of our study, we find out that of the 65.2% who used herbal medicine, 30.4% used it to improve health, 10.3% used it to prevent diseases, and 46.1% used it to treat their diseases, while the rest claimed to use it without any reason.

Amongst the factors that exhibited a relationship with the use of herbal medicines were social demographic factors, the presence of multiple chronic conditions, and the type of illness [26]. In contrast, none of these exhibited any relationship from our study except the purpose of use and the reasons for use ($p=0.00$). Okwuonu *et al.* reported on the belief of rural populations on the utilization of herbal medications for the treatment of kidney diseases in the same geopolitical zone [7]. This study is the only study similar to our study conducted in Nigeria. However, its limitation is that it did not focus on all chronic disease patients and only on rural dwellers. The majority (83.2%) believed in alternative therapy in the management of kidney disease, which is higher than the rate from our study could be due to the fact that it was conducted in rural areas that are predominantly farmers and who do not earn enough to afford conventional medicines in that management of their illness.

Additionally, there was no connection between this assumption and the participants' educational backgrounds [27], which is in line with the findings of our study.

Li *et al.* also reported on using herbal medications among women in the same country selected from an adult population of those with breast cancer. The percentage of people who used traditional medicine was 81.6% [23]. In addition, it was found that there was a positive correlation between ethnicity, educational background, and the rate of traditional medicine use. Women from the Yoruba ethnic extraction used herbal medicine more than Igbo and Hausa women. In contrast, educated women had a lower probability of using herbal medicine than non-educated women, which proved true as most of our study population comes from the Igbo ethnic group. This could be the reason for a reduction in herbal medicine used compared to the study by Li *et al.* [23]. The highest usage is seen in women whose highest education level is secondary education. This did not play any role in our study as educational level had no association with herbal medicine use.

The study also showed the source of information for herbal medicine use ranging from family/friends, which is the major source of information, radio/T.V. advertisements, medical experts, the internet, books/magazine, and seminars/workshop. This source can also be exploited to counter misinformation concerning herbal medicine use and help tackle this issue of herbal medicine use.

Previous studies have shown that the reason for herbal medicine use is lack of regulation, ease of reaching the products, and excessive media advertisement [30], which was in tandem with our study's findings. However, additional reasons were also found, which include cost, family tradition, positive history of health outcomes with herbal medicine use, dissatisfaction with conventional medicine, cost, belief that it has reduced side effects since it's from nature (it is important to note that, we didn't access the claims of this patients clinically as it didn't form part of our scope) and opinion of neighbors on its use to patients, etc.

SUMMARY?

CONCLUSION

This study found that people with chronic diseases in Nigeria frequently use herbal medications.

There are several considerations (usage motivations, information sources, Etc.) that are connected to using herbal remedies. Healthcare professionals and policymakers will be helped by this information when making decisions about using herbal medicine. Additionally, doctors should be aware of their patients' use of herbal medications and educate them on their efficacy and potential side effects.

STUDY LIMITATIONS

Since the study was cross-sectional, no inferences on a cause-and-effect link can be made. Results cannot be extrapolated to other regions of Nigeria because the study was also carried out in a teaching hospital in South-eastern Nigeria. Self-reporting was used to collect the data for the assessment, which may have resulted in either an under or over-reporting of the use of herbal medications. Because they believe traditional medical professionals disapprove of the use of herbal medicines, some patients may not have been entirely truthful when they admitted to using them. Future studies should investigate several critical factors that could improve the use of herbal medications, including patient-provider communication about herbal medicine use and the types of herbal medicines used for the illness.

ABBREVIATIONS

OR: Odds ratio; WHO: World Health Organization; CAM: Complementary and alternative medicine; UNTH: University of Nigeria Teaching Hospital; CI: Confidence interval; H.M.: Herbal Medicine.

REFERENCES

1. Ekor M. The Growing Use of Herbal Medicines: Issues Relating to Adverse Reactions and Challenges in Monitoring Safety. *Frontiers in pharmacology*, 2014);4:177. <https://doi.org/10.3389/fphar.2013.00177>
2. Adisa R, Fakeye T. Assessment of the knowledge of community pharmacists regarding common phytopharmaceuticals sold in South Western Nigeria. *Tropical Journal of Pharmaceutical Research*, 2006;5(2):619-625.
3. Nissen M. Factors Responsible for Increased Use of Herbal Medicines and Self-Medication. *Advances in Pharmacoepidemiology And Drug Safety*, 2022;11(3):10–11.
4. Zhang J, Onakpoya IJ, Posadzki P, Eddouks M. The Safety of Herbal Medicine: From Prejudice to Evidence. *Evidence-Based Complementary and Alternative Medicine*, 2015.
5. Osemene KP. A Comparative Assessment of Herbal and Orthodox Medicines in Nigeria. *Research Journal of Medical Sciences*, 2017;5(5):280–285.
6. Pathak K, Ratna JD. Herbal Medicine - A Rational Approach in Health Care System. *International Journal of Herbal Medicine*, 2013;1(3):86-89.

7. Okwuonu CG, Ezeani IU, Olorokor AB, Aniede EF. Belief in Complementary and Alternative Medicine in the Management of Kidney Diseases in a Rural Population of South-east Nigeria. *International Journal of Medicine and Biomedical Research*, 2014;3(3):168–177.
8. Olorunniyi OF, Morenikeji OA. The Extent of Use of Herbal Medicine in Malaria Management in Ido / Osi Local Government Area of Ekiti, Nigeria. *Journal of Medicinal Plants Research* 2013;7(42):3171–3178.
9. Okoronkwo I, Okpala P, Agbo M, Ndu A. Patterns of Complementary and Alternative Medicine Use, Perceived Benefits, and Adverse Effects Among Adult Users in Enugu Urban, Southeast Nigeria. *Evidence-based complementary and alternative medicine: eCAM*, 2014;239372. <https://doi.org/10.1155/2014/239372>
10. Duru CB, Uwakwe KA, Chinomso NC, Mbachi II, Diwe KC, Agunwa CC, Iwu AC, Merenu IA. Socio-Demographic Determinants of Herbal Medicine Use in Pregnancy among Nigerian Women Attending Clinics in a Tertiary Hospital in Imo State, South-East, Nigeria. *American Journal of Medicine Studies*, 2016;4(1):1-10.
11. Oreagba IA, Oshikoya KA, Amachree M. Herbal Medicine Use among Urban Residents in Lagos, Nigeria. *BMC Complementary and Alternative Medicine*, 2011;11(117):1–8.
12. Bodeker G, Ong CK, Grundy C. *WHO Global Atlas of Traditional, Complementary and Alternative Medicine (Vol.1)*. World Health Organization. 2005
13. Bandaranayake WM. Quality Control, Screening, Toxicity, and Regulation of Herbal Drugs. *Modern phytomedicine: turning medicinal plants into drugs*, 2006:25-57.
14. Nissen N. Complementary Therapies in Clinical Practice: Practitioners of Western Herbal Medicine and Their Practice in the U.S.: Beginning to Sketch the Profession. *Complementary Therapies in Clinical Practice*, 2010;16(4):181–186.
15. Calapai G. European Legislation on Herbal Medicines: A Look into the Future. *Drugs safety*, 2008;31(5):428–431. <https://doi.org/10.2165/00002018-200831050-00009>
16. Olusanya BO, Inem VA, Abosede OA. Infants Delivered in Maternity Homes Run by Traditional Birth Attendants in Urban Nigeria: A Community-

Based on *Health Care for Women International*, 2011;32(6):474-491.

17. Sarmiento I, Zuluaga G, Andersson N. Traditional medicine used in childbirth and for childhood diarrhea in Nigeria's Cross River State: Interviews with traditional practitioners and a statewide cross-sectional study. *BMJ Open*, 2016;6(4):e010417.
18. Girma E, Tesfaye M. Patterns of treatment seeking behavior for mental illnesses in South west Ethiopia: a hospital-based study. *BMC Psychiatry* 2011;11:138 <https://doi.org/10.1186/1471-244X-11-138>
19. Awodele O. Doctor's Attitudes Towards The Use Of Herbal Medicine In Lagos, Nigeria. *Journal of Herbal Medicine*, 2012;2(1):16–22.
20. Mbutho NP, Gqaleni N, Korporaal CM, Africa S, Medicine T. Traditional Complementary and Alternative Medicine: Knowledge, Attitudes and Practices of Health Care Workers in HIV and AIDS Clinics in Durban Hospitals. *African journal of traditional, complementary, and alternative medicines: AJTCAM*, 2012;9(3):64–72. <https://doi.org/10.4314/ajtcam.v9i3s.8>
21. Van Staden AM, Joubert Georgina BA. Interest in and Willingness to Use Complementary, Alternative and Traditional Medicine among Academic and Administrative University Staff in Bloemfontein, South Africa. *African Journal of Traditional, Complementary and Alternative Medicine*, 2014;11(5):61-66.
22. Elechi-Amadi K, Briggs ON, Konne F, Giami L. Perception and Acceptance Of Herbal Medicines among Residents of Port Harcourt , Nigeria. *Journal of Complementary and Alternative Medical Research*, 2021;12(3):24–34.
23. Li S, Odedina S, Agwai I, Ojengbede O, Huo D, Olopade OI. Traditional Medicine Usage among Adult Women in Ibadan, Nigeria: A Cross-Sectional Study. *BMCComplementary Medicine and Therapies*, 2020;20(93):1–7.
24. Amaeze OU, Aderemi-Williams RI, Ayo-Vaughan MA, Ogundemuren DA, Ogunmola DS, Anyika EN. Herbal medicine Use among type 2 diabetes mellitus patients in Nigeria: understanding the magnitude and predictors of use. *International Journal of Clinical Pharmacy*, 2018;40(3):580-588.
25. Tulunay M, Aypak C, Yikilkan H, Gorpelioglu S. Herbal Medicine Use Among Patients with Chronic Diseases. *Journal of Intercultural Ethnopharmacology*, 2015;4(3):217–220. <https://doi.org/10.5455/Jice.20150623090040>

26. Pengpid S. The Use of Herbal Medicines Among Chronic Disease Patients in Thailand: A Cross-Sectional Survey. *Journal of Multidisciplinary Healthcare*, 2019;12:573-582.
27. Al-Azzawi AM, Mehdi N, Al-Juboori AG, Ejaz A, Ali H, Sarheed O. Herbal Medicines Questionnaire and Evaluation Of Attitude, Perceptions and Self-Use Among Health Care Professionals In Rak , UAE : Pilot Study. *International Journal of Pharmacy and Pharmaceutical Sciences*, 2019;11(3):86–91. doi:10.22159/ijpps.2019v11i3.29945.

UNDER PEER REVIEW

UNDER PEER REVIEW