

Review Article

Unravelling the Genetic Enigma: Exploring the Molecular Basis of Heterosis

ABSTRACT

Heterosis, the phenomenon where hybrid offspring exhibit superior traits compared to their inbred parents, has captivated scientists for decades due to its immense potential for crop improvement. Unravelling the molecular basis of heterosis has remained a challenge, but recent advancements in genomic and molecular techniques have shed new light on this intriguing phenomenon. This review paper provides a comprehensive overview of the current understanding of the molecular mechanisms underlying heterosis. We delve into the intricate interplay of genetic factors and regulatory networks that contribute to the manifestation of heterosis. Genomic studies have revealed structural variations, such as rearrangements and copy number variations, which may play a crucial role in hybrid vigour. Furthermore, we explore the dynamic nature of gene expression patterns during hybridization, highlighting the activation of novel gene networks and the modulation of epigenetic mechanisms. Transcriptomic analyses have identified key genes and pathways associated with heterosis, shedding light on the intricate molecular interactions that drive the superior performance of hybrids. Additionally, we discuss the emerging role of small RNAs, transcription factors, and regulatory elements in orchestrating heterosis. These molecular players act as conductors in the genetic symphony of hybrid vigour, fine-tuning gene expression and facilitating the coordinated functioning of diverse biological processes. Understanding the molecular basis of heterosis holds immense promise for crop improvement and sustainable agriculture. By leveraging this knowledge, breeders can harness the full potential of hybrid breeding to develop high-yielding and resilient crop varieties. This review aims to inspire further research and collaboration in unravelling the mysteries of heterosis, paving the way for innovative strategies to enhance global food security and agricultural productivity.

Keywords: *Heterosis, Transcription Factors, Hybrid Vigour, Crop Improvement, Resilient Crop, Hybrid Breeding.*

1. INTRODUCTION

Heterosis, also known as hybrid vigour, is a phenomenon that has fascinated scientists and breeders for over a century [1]. It refers to the superior performance of hybrid offspring compared to their inbred parental lines in various traits, such as yield, growth rate, disease resistance, and stress tolerance. The exploitation of heterosis has revolutionized agriculture by significantly improving crop productivity and quality. Shull [2] discovered and initially introduced the idea of heterosis by observing its occurrence in hybrid offspring of maize. He later officially coined the term "heterosis" to describe this phenomenon. The concept of heterosis was first implemented in genetic breeding of maize, leading to the production of numerous high-quality maize hybrids starting from the 1930s [3]. Notably, since 2011, the cultivation of hybrids has contributed significantly to an eightfold increase in maize yield in America [4]. Despite the practical importance of heterosis, its underlying genetic and molecular mechanisms have remained elusive. Understanding the molecular basis of heterosis is crucial for unravelling the secrets behind this phenomenon and harnessing its

potential for crop improvement. Koelreuter [5] presented the initial empirical evidence demonstrating that the growth of hybrid tobacco surpasses that of its parental plants.

Over the years, numerous studies have attempted to decipher the intricate genetic and molecular interactions that contribute to heterosis. Recent advancements in genomics, transcriptomic and molecular techniques have provided unprecedented insights into the molecular underpinnings of this phenomenon. F₁-hybrid plants exhibit various advantageous characteristics compared to their homozygous parental inbred lines, such as increased biomass, size, yield, development speed, fertility, disease resistance, resistance to insect pests, and tolerance to challenging climates [6, 7]. However, if hybrids are self-pollinated over multiple generations, the level of heterozygosity and vigour gradually decreases, leading to a phenomenon known as inbreeding depression (refer to Glossary). Therefore, heterosis and inbreeding depression are two distinct aspects of the same phenomenon [3]. Charles Darwin first described heterosis in 1876 when he noticed that cross-pollinated maize (*Zea mays*) progeny were 25% taller than inbred maize progeny [8].

At the genomic level, researchers have discovered structural variations, such as chromosomal rearrangements, copy number variations, and transposable element mobilizations, which are associated with heterosis. These alterations can lead to changes in gene dosage, gene expression, and regulatory networks, ultimately influencing the phenotypic traits observed in hybrid offspring.

The concepts of midparent heterosis (MPH) and best parent heterosis (BPH) quantify the extent of phenotypic difference exhibited by a trait in a hybrid (F₁) compared to its parental inbred lines (P₁, P₂) [9]. MPH denotes a trait that demonstrates a hybrid performance significantly superior to the average value (midparent) calculated from the two parental inbred lines ($MPH = F_1 * [(P_1 + P_2)/2]$). On the other hand, BPH indicates a hybrid trait that performs significantly better than the superior does (BP) of the two homozygous parental inbred lines ($BPH = F_1 * BP$) [10]. In addition, studies on gene expression patterns have revealed the presence of unique transcriptomic profiles in hybrid plants compared to their parental lines. Complex regulatory networks, involving transcription factors, small RNAs, epigenetic modifications, and hormone signaling pathways, have been implicated in orchestrating the differential gene expression patterns that contribute to hybrid vigour [11]. The molecular basis of heterosis is a multifaceted puzzle that requires an integrative approach, combining genetics, genomics, epigenetics, and systems biology. By unravelling the genetic and molecular factors underlying heterosis, scientists aim to develop strategies for predicting hybrid performance, optimizing parental combinations, and manipulating key genes and pathways to enhance desired traits. In this review, we aim to provide a comprehensive overview of the current understanding of the molecular basis of heterosis. We will explore the latest research findings, discuss the key genetic and molecular players involved, and highlight the challenges and future directions in this exciting field of study. By gaining a deeper understanding of the molecular mechanisms driving heterosis, we can unlock its full potential for sustainable agriculture and meet the challenges of global food security [12].

Recently, several laboratories have commenced examining various molecular aspects of heterosis utilizing newly available molecular tools. In the past, the analysis of quantitative trait loci (QTLs) served as an initial step in comprehending the molecular underpinnings of heterosis [21]. While these studies demonstrated that heterosis is determined by a limited number of individual genes inherited in a complex manner, none of the QTLs associated with heterotic traits have been successfully cloned thus far. However, with the advent of innovative molecular tools enabling comprehensive phenotyping based on QTL analysis, followed by map-based cloning, there is a potential for identifying the loci responsible for controlling heterotic traits in the foreseeable future [27].

2. STRATEGIES FOR HETEROSIS BREEDING

Conducting extensive hybridization tests to obtain hybrid F₁ lines exhibiting heterosis is not recommended due to the significant resources, time, and unreliable results it entails [13]. Melchinger and Gumber [14] proposed an alternative approach of utilizing heterotic groups as a foundation for crossbreeding. Heterotic groups are populations classified based on their breeding requirements, possessing abundant genetic variation and high combining ability. In a genome-wide association study (GWAS) conducted by Chen *et al.* [15] on yield traits, general combining ability (GCA), and specific combining ability (SCA) in rice, the use of combining ability as a basis for classifying heterotic groups was strongly supported, providing valuable insights for studying combining ability in vegetables (Figure. 1). Other studies have also demonstrated that combining ability, genetic distance, and molecular markers can serve as the foundation for evaluating parental inbred lines and predicting heterosis in F₁ hybrids of vegetables [16-17].

General combining ability (GCA) represents the average performance of a specific set of hybrid combinations and is primarily influenced by additive gene effects and additive × additive interactions. On the other hand, specific combining ability (SCA) assesses the average performance of particular hybrid combinations in relation to the parental lines and arises from dominance, epistatic deviations, and genotype × environmental interactions [18]. Parents with a high GCA effect exhibit greater adaptability and are less influenced by environmental factors [19]. It is important to note that superior traits in parents do not always guarantee their inheritance in offspring [20]. Hence, evaluating combining ability provides more reliable information than solely assessing the performance of individual lines. Various types of combining ability tests can be employed to identify superior parental lines for developing heterotic hybrids, such as line × tester analysis, topcross tests, single-cross tests, poly-cross tests, and diallel mating [21]. Singh *et al.* [22] conducted a complete diallel cross test on seven diverse bitter melon lines and observed that combinations with high × high GCA often produced substantial SCA effects, making them potential candidates for developing superior variants through the pedigree method. Combinations involving high/low × low GCA can also exhibit high but less stable SCA effects, which are suitable for heterosis breeding. This finding aligns with the results reported by Kenga *et al.* [23] in sweet sorghum and Franco *et al.* [24] in common bean.




Figure 1. Factors involved in applying heterosis-breeding strategies: obtaining heterotic lines and maintaining heterosis in the elite lines in the offspring [26].

HETEROSIS-ASSOCIATED GENE EXPRESSION IN RICE & MAIZE

In maize, the presence of small gene families contributes to the complexity of understanding the quantitative effects of gene deletions in inbred lines. It is likely that these genes are

functionally compensated by duplicate copies located elsewhere in the genome, resulting in only minor effects on plant performance [58]. In recent years, several studies have focused on analyzing gene expression associated with heterosis in maize and rice. These studies have employed different approaches, such as comparing gene expression patterns between inbred lines and hybrids using selected genes or conducting high-throughput gene expression analyses through microarray profiling or Gene Calling [31]. Various stages of plant development have been investigated, including embryos, endosperm, seedlings, shoot apical meristems, ear tissue, leaves, and rice panicles, using different genetic backgrounds [31,33] (Table 1).

However, it is important to note that these studies exhibited significant variations in their experimental designs, statistical procedures, and tissues and genotypes analyzed. Consequently, no consistent global expression patterns were observed across all the gene expression studies. Some studies highlighted non-additive gene expression as prevalent between inbred lines and hybrids, while others observed predominantly additive gene expression for most genes. Additionally, one study reported an equal number of genes exhibiting additive and non-additive expression [31, 33] (Table 1). The discrepancies in these findings may stem from the differences in genotypes, plant materials, experimental designs, and statistical procedures employed in the respective studies.

In summary, the analysis of gene expression associated with heterosis has yielded diverse and sometimes conflicting results. The variations in experimental approaches and genetic backgrounds utilized by different studies contribute to the lack of uniformity in global expression patterns. Future research should aim to address these discrepancies by employing standardized experimental designs and statistical procedures, as well as conducting comparative analyses across multiple genetic backgrounds, to gain a more comprehensive understanding of heterosis-related gene expression.

Table 1. Heterosis-associated gene expression in Rice and maize [28]


Plant organ	Developmental stage	Approach	Genetic background	Global expression trend	Refs
Rice					
Panicle	Stage III, IV, V	9K cDNA microarrays	Zhenshan97 Minghui63	Additivity	[29]
Maize					
Embryo	6 DAP	12K cDNA microarrays SSH qRT-PCR	UH005 UH301	Additivity	[30]
Endosperm	10, 14, 21 DAP	Gene Calling	7 Pioneer inbred lines	Non-additivity	[31]
Endosperm	18 DAP	RT-PCR	B73 BSSS53	Non-additivity	[32]
Embryo	19 DAP	13.5K microarrays	Mo17	Additivity	[33]
Seedling	11 DAG		B73		
Immature ear					
Seedling	14 DAG	14K cDNA microarrays qRT-PCR	Mo17 B73	Additivity	[34]

Shoot apical meristem	21–23 DAP	12K cDNA microarrays qRT-PCR	UH002 UH005 UH250 UH301	Non-additivity	[35]
Adult leaves of diploids and triploids		Quantitative Northern blotting	Mo17 B73	Non-additivity	[36]
Immature ear		Gene Calling	17 Pioneer inbred lines	Additivity and Non-additivity	[37]

Abbreviations: DAP, days after pollination; DAG, days after germination; SSH, suppressive subtractive hybridization.

VARIOUS MODELS FOR UNDERSTANDING THE MOLECULAR BASIS OF HETEROSIS

Exploring heterosis at the molecular level involves various approaches, such as genome organization studies, transcriptome-wide gene expression profiling, and investigating allele-specific contributions to gene expression [38]. These molecular investigations ultimately converge on fundamental and widely accepted models of heterosis, which include the combined expression of alleles and diverse interactions between alleles in a hybrid [39]. Despite ongoing research efforts, heterosis remains a captivating and intriguing subject, prompting continuous exploration and the acquisition of additional knowledge. Recent advancements in epigenetics, genomics, proteomics, and metabolomics have played a crucial role in unravelling the mysteries of this phenomenon. In this context, we present a summary and comprehension of the current perspectives that shed light on the existence of hybrid vigour in plant species. Several models have been proposed, represented in (Figure 2) to understand the molecular basis of heterosis. Here are some of the prominent models:-



1. Dominance Model: This classical model suggests that heterosis arises from the dominance of advantageous alleles in hybrids [40]. The superior performance of hybrids is attributed to the masking of deleterious recessive alleles by dominant alleles [41-42].

2. Over dominance/Epistasis Model: This model proposes that heterosis is driven by the interaction between different alleles at multiple loci [43]. The combined effect of these alleles in hybrids leads to a superior phenotype that exceeds the performance of both parents [44-46].

3. Additive Model: According to this model, heterosis results from the accumulation of additive genetic effects across multiple loci. The interaction between additive alleles leads to enhanced performance in hybrids [47].

4. Combinatorial/Network Model: This model emphasizes the importance of gene regulatory networks and interactions between genes. Heterosis is thought to arise from the combinatorial effects of genes and their interactions within these networks [48].

5. Metabolic Model: This model focuses on the metabolic pathways and biochemical processes involved in heterosis. It suggests that the increased metabolic activity and efficiency in hybrids contribute to their superior performance [49].

6. Epigenetic Model: Epigenetic modifications, such as DNA methylation and histone modifications, have been proposed to play a role in heterosis. This model suggests that changes in the epigenetic landscape contribute to the altered gene expression patterns and phenotypic superiority observed in hybrids [50].

It is important to note that these models are not mutually exclusive, and the molecular basis of heterosis is likely a complex interplay of multiple factors. Advances in genomics, transcriptomics, proteomics, and other molecular techniques continue to shed light on the underlying mechanisms of heterosis and may lead to the development of new models and hypotheses.

FUTURE SCOPE OF MOLECULAR HETEROSIS

The future of molecular heterosis holds promising possibilities for advancing our understanding and utilization of hybrid vigour. Here are some potential aspects to consider:

1. Genomic Selection: Genomic selection techniques can facilitate the identification of genomic regions associated with heterotic traits. By analyzing large-scale genomic data, researchers can predict hybrid performance and select parental lines with higher combining ability more efficiently [51].

2. Gene Editing and Manipulation: Advanced gene editing technologies like CRISPR-Cas9 offer the potential to precisely manipulate specific genes and regulatory elements involved in heterosis [52]. This could enable targeted enhancements of heterotic traits and the creation of custom-designed hybrids with desired characteristics.

3. Transcriptomics and Epigenetics: Investigating gene expression patterns and epigenetic modifications during hybridization can provide insights into the molecular mechanisms underlying heterosis. Transcriptomic studies can identify key gene networks and pathways associated with hybrid vigour, while epigenetic analysis can uncover heritable changes in gene expression patterns that contribute to heterosis [53].

4. Multi-Omics Integration: Integrating data from multiple omics levels, such as genomics, transcriptomics, proteomics, and metabolomics, can provide a comprehensive understanding of the molecular basis of heterosis. This holistic approach allows for a more detailed characterization of the complex interactions and regulatory networks involved in hybrid vigour [54].

5. Machine Learning and Artificial Intelligence: The application of machine learning algorithms and artificial intelligence techniques can help analyse vast amounts of molecular data and extract meaningful patterns and predictive models related to heterosis. These methods can assist in identifying genetic factors, biomarkers, and molecular signatures associated with hybrid performance [55].

6. Hybrid Breeding Strategies: Incorporating molecular markers and genomic information into traditional breeding programs can optimize hybrid-breeding strategies. Utilizing marker-assisted selection and genomic prediction can enhance the efficiency and precision of hybrid development, leading to the production of superior hybrids with improved performance [56].

7. Systems Biology Approaches: Adopting systems biology approaches can provide a holistic understanding of the complex interactions among genes, proteins, and

metabolites that contribute to heterosis. Integrating data from different biological levels can unravel emergent properties and regulatory networks that drive hybrid vigour [57].

By integrating these future aspects of molecular heterosis, researchers can deepen our understanding of the underlying molecular mechanisms, accelerate the development of improved hybrids, and ultimately enhance agricultural productivity and sustainability.

CONCLUSION

In conclusion, the study of heterosis at the molecular level has provided valuable insights into the mechanisms underlying this complex phenomenon. Various models, including the dominance model, over dominance/epistasis model, additive model, combinatorial/network model, metabolic model, and epigenetic model, have been proposed to explain the molecular basis of heterosis. These models highlight the importance of gene expression, allelic interactions, metabolic processes, and epigenetic modifications in driving hybrid vigour. While classical genetic models based on dominance and additive effects remain popular among plant breeders, it is becoming increasingly evident that the understanding of heterosis requires a broader perspective. The context of genome dynamics, particularly in polyploid species, adds complexity to the interpretation of hybrid vigour. The use of molecular markers, allozymes, and sequencing plant genomes has been suggested as valuable tools for unravelling the genetic basis of hybrid vigour and understanding the impact of self-pollination on this phenomenon. Furthermore, the application of modern molecular techniques, such as genomics, transcriptomics, proteomics, and metabolomics, holds great potential for advancing our knowledge of heterosis. These techniques enable the exploration of genome-wide changes in gene and protein expression, as well as the investigation of epigenomic and metabolic alterations in hybrids. In summary, the study of heterosis has evolved beyond classical genetic models, with increasing emphasis on molecular approaches. The integration of various models and the utilization of advanced molecular techniques have the potential to provide a more comprehensive understanding of heterosis and its underlying molecular mechanisms. This knowledge can contribute to the development of new breeding strategies and the enhancement of crop productivity by harnessing the power of hybrid vigour.

REFERENCES

1. Shull GH. What is "heterosis"? *Genetics*. 1948 Sep;33(5):439.
2. Shull GH. The composition of a field of maize. *Journal of Heredity*. 1908 Jan 1(1):296-301.
3. Shull GH. Duplicate genes for capsule-form in *Bursa bursa-pastoris*. *Z Indukt Abstamm Vererbungsl*. 1914 Dec;12(1):97-149.
4. Tian F, Bradbury PJ, Brown PJ, Hung H, Sun Q, Flint-Garcia S, Rocheford TR, McMullen MD, Holland JB, Buckler ES. Genome-wide association study of leaf architecture in the maize nested association mapping population. *Nature genetics*. 2011 Feb;43(2):159-62.
5. Koelreuter J. In *Methods of Plant Breeding* (eds. Hayes HK, Immer FR, Smith BC.) McGraw Hill Book Co. Inc., 1763.
6. Falconer DS, Mackay TFC. *Introduction to quantitative genetics* (4th edn), Longman, eds 1996.
7. Zirkle C. Early ideas on inbreeding and crossbreeding. *Heterosis*. 1952:1-3.

8. Darwin CR. The effects of cross- and self-fertilization in the vegetable kingdom (1st edn), John Murray, ed. 1876.
9. Li Z, Sun J, Hirsch CN. Understanding Environmental Modulation of Heterosis. *Plant Breeding Reviews*. 2022 Nov 18;46:219-37.
10. Feys K, Demuynck K, De Block J, Bisht A, De Vliegheer A, Inzé D, Nelissen H. Growth rate rather than growth duration drives growth heterosis in maize B104 hybrids. *Plant, Cell & Environment*. 2018 Feb;41(2):374-82.
11. Chen ZJ. Genomic and epigenetic insights into the molecular bases of heterosis. *Nature Reviews Genetics*. 2013 Jul;14(7):471-82.
12. Purugganan MD, Jackson SA. Advancing crop genomics from lab to field. *Nature Genetics*. 2021 May;53(5):595-601.
13. Singh PK, Dasgupta SK, Tripathi SK. Hybrid vegetable development. CRC Press; 2005 Jun 14.
14. Melchinger AE, Gumber RK. Overview of heterosis and heterotic groups in agronomic crops. Concepts and breeding of heterosis in crop plants. 1998 Jan 1;25:29-44.
15. Chen J, Zhou H, Xie W, Xia D, Gao G, Zhang Q, Wang G, Lian X, Xiao J, He Y. Genome-wide association analyses reveal the genetic basis of combining ability in rice. *Plant biotechnology journal*. 2019 Nov;17(11):2211-22.
16. Mather K. The balance of polygenic combinations. *Journal of Genetics*. 1942 Apr;43:309-36.
17. Griffing BR. Concept of general and specific combining ability in relation to diallel crossing systems. *Australian journal of biological sciences*. 1956;9(4):463-93.
18. Geleta LF, Labuschagne MT, Viljoen CD. Relationship between heterosis and genetic distance based on morphological traits and AFLP markers in pepper. *Plant Breeding*. 2004 Oct;123(5):467-73.
19. Jagosz B. The relationship between heterosis and genetic distances based on RAPD and AFLP markers in carrot. *Plant breeding*. 2011 Oct;130(5):574-9.
20. Sharma BB, Sharma VK, Dhakar MK, Punetha S. Combining ability and gene action studies for horticultural traits in garden pea: A review. *African Journal of Agricultural Research*. 2013;8(38):4718.
21. Fasahat P, Rajabi A, Rad JM, Derera JJ. Principles and utilization of combining ability in plant breeding. *Biometrics & Biostatistics International Journal*. 2016;4(1):1-24.
22. Rajendrakumar P, Hariprasanna K, Seetharama N. Prediction of heterosis in crop plants—status and prospects. *J. Exp. Agric*. 2015 Aug 4;9:1-6.
23. Singh AK, Pan RS, Bhavana P. Heterosis and combining ability analysis in bittergourd (*Momordica charantia* L.).
24. Kenga R, Alabi SO, Gupta SC. Combining ability studies in tropical sorghum (*Sorghum bicolor* (L.) Moench). *Field crops research*. 2004 Aug 10;88(2-3):251-60.
25. Caixeta Franco M, Tulio Cassini S, Rodrigues Oliveira V, Vieira C, Mui Tsai S, Damiao Cruz C. Combining ability for nodulation in common bean (*Phaseolus vulgaris* L.) genotypes from Andean and Middle American gene pools. *Euphytica*. 2001 Apr;118:265-70.
26. Yu D, Gu X, Zhang S, Dong S, Miao H, Gebretsadik K, Bo K. Molecular basis of heterosis and related breeding strategies reveal its importance in vegetable breeding. *Horticulture research*. 2021 Dec 1;8.
27. Lippman ZB, Zamir D. Heterosis: revisiting the magic. *Trends in genetics*. 2007 Feb 1;23(2):60-6.
28. Hochholdinger F, Hoecker N. Towards the molecular basis of heterosis. *Trends in plant science*. 2007 Sep 1;12(9):427-32.
29. Huang Y, Zhang L, Zhang J, Yuan D, Xu C, Li X, Zhou D, Wang S, Zhang Q. Heterosis and polymorphisms of gene expression in an elite rice hybrid as revealed

- by a microarray analysis of 9198 unique ESTs. *Plant molecular biology*. 2006 Nov;62:579-91.
30. Meyer S, Pospisil H, Scholten S. Heterosis associated gene expression in maize embryos 6 days after fertilization exhibits additive, dominant and overdominant pattern. *Plant molecular biology*. 2007 Feb;63:381-91.
 31. Guo M, Rupe MA, Danilevskaya ON, Yang X, Hu Z. Genome-wide mRNA profiling reveals heterochronic allelic variation and a new imprinted gene in hybrid maize endosperm. *The Plant Journal*. 2003 Oct;36(1):30-44.
 32. Song R, Messing J. Gene expression of a gene family in maize based on noncollinear haplotypes. *Proceedings of the National Academy of Sciences*. 2003 Jul 22;100(15):9055-60.
 33. Stupar RM, Springer NM. Cis-transcriptional variation in maize inbred lines B73 and Mo17 leads to additive expression patterns in the F1 hybrid. *Genetics*. 2006 Aug 1;173(4):2199-210.
 34. Swanson-Wagner RA, Jia Y, DeCook R, Borsuk LA, Nettleton D, Schnable PS. All possible modes of gene action are observed in a global comparison of gene expression in a maize F1 hybrid and its inbred parents. *Proceedings of the National Academy of Sciences*. 2006 May 2;103(18):6805-10.
 35. Użarowska A, Keller B, Piepho HP, Schwarz G, Ingvaridsen C, Wenzel G, Lübberstedt T. Comparative expression profiling in meristems of inbred-hybrid triplets of maize based on morphological investigations of heterosis for plant height. *Plant molecular biology*. 2007 Jan;63:21-34.
 36. Auger DL, Gray AD, Ream TS, Kato A, Coe Jr EH, Birchler JA. Nonadditive gene expression in diploid and triploid hybrids of maize. *Genetics*. 2005 Jan 1;169(1):389-97.
 37. Guo M, Rupe MA, Yang X, Crasta O, Zinselmeier C, Smith OS, Bowen B. Genome-wide transcript analysis of maize hybrids: allelic additive gene expression and yield heterosis. *Theoretical and Applied Genetics*. 2006 Sep;113:831-45.
 38. Cheng SH, Zhuang JY, Fan YY, Du JH, Cao LY. Progress in research and development on hybrid rice: a super-domesticated in China. *Annals of botany*. 2007 Oct 1;100(5):959-66.
 39. Birchler JA, Auger DL, Riddle NC. In search of the molecular basis of heterosis. *The Plant Cell*. 2003 Oct;15(10):2236-9.
 40. Schnable PS, Springer NM. Progress toward understanding heterosis in crop plants. *Annual review of plant biology*. 2013 Apr 29;64:71-88.
 41. Ryder P, McKeown PC, Fort A, Spillane C. Epigenetics and heterosis in crop plants. *Epigenetics in Plants of Agronomic Importance: Fundamentals and Applications: Transcriptional Regulation and Chromatin Remodelling in Plants*. 2019:129-47.
 42. Yao H, Dogra Gray A, Auger DL, Birchler JA. Genomic dosage effects on heterosis in triploid maize. *Proceedings of the National Academy of Sciences*. 2013 Feb 12;110(7):2665-9.
 43. Birchler JA, Yao H, Chudalayandi S, Vaiman D, Veitia RA. Heterosis. *The Plant Cell*. 2010 Jul;22(7):2105-12.
 44. Birchler JA, Yao H, Chudalayandi S. Unraveling the genetic basis of hybrid vigor. *Proceedings of the National Academy of Sciences*. 2006 Aug 29;103(35):12957-8.
 45. Powers L. An expansion of Jones's theory for the explanation of heterosis. *The American Naturalist*. 1944 May 1;78(776):275-80.
 46. Liang Q, Shang L, Wang Y, Hua J. Partial dominance, overdominance and epistasis as the genetic basis of heterosis in upland cotton (*Gossypium hirsutum* L.). *PLoS One*. 2015 Nov 30;10(11):e0143548.
 47. Jizhong S. A review on the research and utilization of hybrid vigor of cotton. *Acta Gossypii Sinica (China)*. 1994.

48. Omholt SW, Plahte E, Øyehaug L, Xiang K. Gene regulatory networks generating the phenomena of additivity, dominance and epistasis. *Genetics*. 2000 Jun 1;155(2):969-80.
49. Song GS, Zhai HL, Peng YG, Zhang L, Wei G, Chen XY, Xiao YG, Wang L, Chen YJ, Wu B, Chen B. Comparative transcriptional profiling and preliminary study on heterosis mechanism of super-hybrid rice. *Molecular plant*. 2010 Nov 1;3(6):1012-25.
50. Kapazoglou A, Ganopoulos I, Tani E, Tsaftaris A. Epigenetics, epigenomics and crop improvement. In *Advances in botanical research 2018* Jan 1 (Vol. 86, pp. 287-324). Academic Press.
51. Stuber CW, Polacco M, Lynn M. Synergy of empirical breeding, marker-assisted selection, and genomics to increase crop yield potential. *Crop Science*. 1999 Nov;39(6):1571-83.
52. Ansari WA, Chandanshive SU, Bhatt V, Nadaf AB, Vats S, Katara JL, Sonah H, Deshmukh R. Genome editing in cereals: approaches, applications and challenges. *International Journal of Molecular Sciences*. 2020 Jun 5;21(11):4040.
53. Liu W, Zhang Y, He H, He G, Deng XW. From hybrid genomes to heterotic trait output: Challenges and opportunities. *Current Opinion in Plant Biology*. 2022 Apr 1;66:102193.
54. Xu Y, Zhang X, Li H, Zheng H, Zhang J, Olsen MS, Varshney RK, Prasanna BM, Qian Q. Smart breeding driven by big data, artificial intelligence and integrated genomic-enviromic prediction. *Molecular Plant*. 2022 Sep 7.
55. Babu R, Nair SK, Prasanna BM, Gupta HS. Integrating marker-assisted selection in crop breeding—prospects and challenges. *Current Science*. 2004 Sep 10:607-19.
56. Schaefer RJ, Michno JM, Myers CL. Unraveling gene function in agricultural species using gene co-expression networks. *Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms*. 2017 Jan 1;1860(1):53-63.
57. Fu D, Xiao M, Hayward A, Jiang G, Zhu L, Zhou Q, Li J, Zhang M. What is crop heterosis: new insights into an old topic. *Journal of applied genetics*. 2015 Feb;56:1-3.
58. Fu H, Dooner HK. Intraspecific violation of genetic colinearity and its implications in maize. *Proceedings of the National Academy of Sciences*. 2002 Jul 9;99(14):9573-8.
59. Rehman A ur, Dang T, Qamar S, Ilyas A, Fatema R, Kafle M, et al. Revisiting Plant Heterosis—From Field Scale to Molecules. *Genes* [Internet]. 2021 Oct 24;12(11):1688.