

## Review Article

# Application of auxins in haploid embryo induction in hexaploidy wheat

### **Abstract:**

Doubled haploid (DH) plant production plays a crucial role in modern plant breeding programs, offering an efficient means to generate homozygous lines from heterozygous parents within a single generation. Different types of auxins have been ~~employed~~-utilized in wheat cross with maize DH production, with 2,4-D being the most widely used and effective hormone, followed by dicamba. Other auxins, including picloram, ~~.....~~(IAA), ~~.....~~(PAA), silver nitrate, ~~.....~~(NAA), kinetin, ~~....~~(BA), and zearalenone, have also been tested for their potential role in haploid embryo induction. Various methods have been explored for the application of 2,4-D, such as spray, tiller injection, dipping, and spikelet culture methods.

**Keywords:** Double haploid (DH), auxin, 2,4-dichlorophenoxyacetic acid (2,4-D), embryo recovery, haploid embryo induction, concentration, efficiency,

### Comments

The abstract lacks conclusion. Moreover, certain abbreviations or acronyms should be fully spelled, while the abbreviations or acronyms are placed in brackets.

### **Introduction:**

Doubled haploid (DH) plant production has emerged as a crucial component of modern plant breeding initiatives (Forster *et al.*, 2007; Weyen, 2009). Through traditional pedigree selection methods, it typically requires around a decade to develop a commercially viable wheat variety. ~~Doubled haploid~~DH technologies serve as effective techniques to expedite the generation of homozygous lines from heterozygous donor genotypes, eliminating the need for time-consuming selfing processes.

Doubled haploids (~~DH~~), which are presently utilized in various crop species such as wheat and triticale, allow breeders to obtain fully homozygous genotypes from heterozygous parents within a single generation (~~Ref???~~). This advancement results in significant time savings of approximately 4 to 6 years in the breeding process. First report of wheat haploid production was reported by crossing wheat with maize (~~Ref???~~).

The initial discovery of wheat haploid production was achieved through the crossing of wheat with maize (Barclay, 1975). After fertilization, the process of zygote induction occurs, leading to the elimination of the male parent's chromosomes from the cells of the developing embryo (Laurie and Bennett, 1989). The viability of these zygotes is limited, and the majority of them undergo abortion during the early stages of development (Laurie and Bennett, 1988). The use of auxins, on the other hand, stimulates the enlargement of the ovary and promotes the development of haploid embryos to a stage where they can be cultured on nutrient media (Laurie and Reymondie, 1991).

Nevertheless, there exist numerous variations in the hormone treatments administered, encompassing factors such as the type and concentration of auxin, as well as the method and timing of hormone application (Ref??). The hormone 2,4-dichlorophenoxyacetic acid (2,4-D) is extensively utilized and is considered the most prevalent, closely followed by dicamba (3,6-dichloro-o-anisic acid) (Ref??). Studies conducted on bread wheat have demonstrated that both hormones exhibit comparable efficacy in inducing desired effects (Matzkand Mahn, 1994).

#### Comments

The objective of the work has not been stated.

The justification of the work needs to be improved.

A number of statements need relevant references.

Most of the references utilized are old.

A nonconsistent pattern of referencing style has been utilized across the document.

#### **Auxins:**

The notion of auxin as a mobile growth regulator was notably deduced by Charles and Frances Darwin, as documented in their renowned 1880 publication, *The Power of Movement in Plants*, where he observed the impact of a hypothetical substance that influenced the elongation of plant shoots, enabling them to exhibit tropic growth towards light (Darwin and Darwin, 1880). Auxin, being a crucial plant hormone, regulates a wide range of processes, including tropic responses to light and gravity, overall root and shoot architecture, organ patterning, vascular development, and growth in tissue culture (Davies, 1995). Human intervention in auxin physiology has facilitated plant propagation and driven by artificial selection, has contributed to the development of current crop varieties (Multani *et al.*, 2003; Salamini, 2003).

#### **Types of auxins used:**

The 2,4-D (hormone-2,4-dichlorophenoxyacetic acid) hormone have has been used

most and shown optimum seed setting and embryo formation followed by close results with dicamba (3,6-dichloro-o-anisic acid) (Laurie and Bennett, 1988; Suenaga and Nakajima, 1989; Riera-Lizarazu and Mujeeb-Kazi, 1990; Kisana *et al.*, 1993; Matzk and Mahn 1994).

Other auxins like picloram (4-amino-3,5,6-trichloropicolinic acid), IAA (indole-3-acetic acid), PAA (phenylacetic acid) (Warcholet *et al.*, 2016), silver nitrate (O'Donoghue and Bennett, 1994), NAA (xxxxxx), kinetin, BA (6-benzyladenine) (Ushiyama *et al.*, 2007) and zearalenone (Biesaga-Kościelniak *et al.*, 2003).

#### **Method of application:**

For application of 2,4-D several methods have been tried by Kaushik *et al.* (2004) such as spray method, tiller injection method, dipping method, and spikelet culture method. with 240 florets pollinated in each method.

**1. Spray method:** The spikes were subjected to spraying with a concentration of 100 ppm of 2,4-dichlorophenoxyacetic acid (2,4-D) at three different time intervals: one, two, and three days after pollination. A total of 8 embryos recovered by this method.

**2. Tiller injection method:** The uppermost internodes of the wheat spikes were injected with 1 ml of 2,4-dichlorophenoxyacetic acid (2,4-D) at a concentration of 100 ppm, starting one day after pollination. This injection process was repeated for three consecutive days. A total of 9 embryos recovered.

**3. Dipping method:** The spikes that underwent pollination were immersed in an aqueous solution of 2,4-dichlorophenoxyacetic acid (2,4-D) for three consecutive days. A total of 14 embryos were recovered.

**4. Spikelet culture method:** On the second day after pollination, the spikes were extracted and subjected to surface sterilization before being dried on filter paper. The rachis (central axis) was then divided into individual spikelets and positioned upright on a growth medium called Murashige-Skoog (MS) media, which contained  $30\text{mg l}^{-1}$  sucrose and  $0.2\text{mg l}^{-1}$  2,4-dichlorophenoxyacetic acid (2,4-D). The cultures were subsequently incubated for a duration of three weeks under continuous light at a temperature of  $20^{\circ}\text{C}$ , with a daily exposure of 16 hours of light. A total of 27 embryos were reported to be recovered by this method.

Applying 2,4-dichlorophenoxyacetic acid (2,4-D) through spray, dipping, and tiller injection methods resulted in low embryo recovery rates. In contrast, when 2,4-D was administered using the spikelet culture method, significantly higher embryo recovery rates were achieved (Kaushik *et al.*, 2004).

**Table 1. Concentration of auxin used with efficiency**

S.no	Auxin	Concentration	Embryo Formation Efficiency	References
1	2,4 D	100 mg l <sup>-1</sup>	48.00%	Ushiyama, T <i>et al.</i> , 2007
2	2,4 D + ZEN	6 μM·dm <sup>-3</sup>	23.30%	Biesaga-Kościelniak, J., <i>et al.</i> , 2003
3	NAA	1000 mg l <sup>-1</sup>	3.40%	Ushiyama, T <i>et al.</i> , 2007
4	IAA	1 mg l <sup>-1</sup>	1.10%	Ushiyama, T <i>et al.</i> , 2007
5	Kinetin	1-10 mg l <sup>-1</sup>	0.90%	Ushiyama, T <i>et al.</i> , 2007
6	6-benzyladenine (BA)	10 mg l <sup>-1</sup>	1.50%	Ushiyama, T <i>et al.</i> , 2007

Formatted: Superscript

Formatted: Superscript

### Conclusion:

Auxins, including 2,4-dichlorophenoxyacetic acid (2,4-D) and dicamba, play a crucial role in inducing haploid embryo development in wheat. The application of these auxins has shown effective results, with 2,4-D being the most widely used hormone. The method of application varies, with techniques such as spray, tiller injection, dipping, and spikelet culture being employed. Among these methods, the spikelet culture method has shown the highest embryo recovery rates.

Different concentrations of auxins have been tested, and their efficiency in inducing embryo formation varies. For instance, 100 mg l<sup>-1</sup> of 2,4-D has been reported to have a 48% embryo formation efficiency. Other auxins such as NAA, IAA, kinetin, and BA have also been studied, each with varying levels of effectiveness in promoting embryo formation. This advancement has significantly contributed to the acceleration of plant breeding programs and the development of improved crop varieties.

### Comments

Auxins are plant hormones that play a critical role in controlling plant growth and development. Even at low concentrations, auxins can regulate gene expression through specific transcription factors and proteins that are modulated to environmental responses in the signalling cascade. Thus, a study on the application of auxins in haploid embryo induction in hexaploidy wheat is vitally important to foster better understanding of what exists in literature on the subject. However, the present research is shallow as there are lots more that exist in literature on the subject

that the authors underutilized. Below are some of the resources they might consider to improve on their work.

Gomes, G. L. B., &Scortecci, K. C. (2021). Auxin and its role in plant development: structure, signalling, regulation and response mechanisms. *Plant biology (Stuttgart, Germany)*, 23(6), 894–904. <https://doi.org/10.1111/plb.13303>

Mahato, A., Chaudhary, H.K. Auxin Induced Haploid Induction in Wide Crosses of Durum Wheat. *CEREAL RESEARCH COMMUNICATIONS* 47, 552–565 (2019). <https://doi.org/10.1556/0806.47.2019.31>

The authors might also consider including write ups on the following subheadings:

Potential of Haploid embryo induction in hexaploidy wheat Breeding Programs

Current Application of Haploid embryo induction in hexaploidy wheat

Generic Framework of Haploid embryo induction in hexaploidy wheat

## References:

1. Barclay, I. R. (1975). High frequencies of haploid production in wheat (*Triticum aestivum*) by chromosome elimination. *Nature*, 256(5516), 410-411.
2. Biesaga-Kościelniak, J., Marcińska, I., Wędzony, M., & Kościelniak, J. (2003). Effect of zearalenone treatment on the production of wheat haploids via the maize pollination system. *Plant cell—Cell reports Reports*, 21, 1035-1039.
3. Darwin, C. (2010). The Works of Charles Darwin, Volume 27: The Power of Movement in Plants. NYU Press.
4. Davies, P. J. (1995). Plant Hormones Dordrecht. *The Netherlands: Kluwer*.
5. Forster, B. P., Heberle-Bors, E., Kasha, K. J., & Touraev, A. (2007). The resurgence of haploids in higher plants. *Trends in plant—Plant science Science*, 12(8), 368-375.
6. Kaushik, N., Sirohi, M., & Khanna, V. K. (2004, September). Influence of age of the embryo and method of hormone application on haploid embryo formation in wheat x maize crosses. In *Proceedings of the 4th International Crop Science Congress. Brisbane, Australia* (Vol. 26).
7. Kisana, N. S., Nkongolo, K. K., Quick, J. S., & Johnson, D. L. (1993). Production of doubled haploids by anther culture and wheat x maize method in a wheat breeding programme. *Plant breeding Breeding*, 110(2), 96-102.
8. Laurie, D. A., & Bennett, M. D. (1988). The production of haploid wheat plants from wheat x maize crosses. *Theoretical and applied—Applied genetics Genetics*, 76, 393-397.
9. Laurie, D. A., & Bennett, M. D. (1988). The production of haploid wheat plants from wheat x maize crosses. *Theoretical and applied—Applied*

Formatted: Indent: Left: 0"

Comment [p1]: The authors have not followed the correct format of referencing for this journal. Please refer to the guidelines.

Formatted: Font: Italic

*genetics* *Genetics*, 76, 393-397.

10. Laurie, D. A., &Reymondie, S. (1991). High frequencies of fertilization and haploid seedling production in crosses between commercial hexaploid wheat varieties and maize. *Plant Breeding*, 106(3), 182-189.
11. Laurie, DA, & Bennett, MD (1989). The timing of chromosome elimination in hexaploid wheat × maize crosses. *Genome*, 32-(6), 953-961.
12. Matzk, F. (1991). A novel approach to differentiated embryos in the absence of endosperm. *Sexual Plant Reproduction*, 4, 88-94.
13. Matzk, F., &Mahn, A. (1994). Improved techniques for haploid production in wheat using chromosome elimination. *Plant Breeding*, 113(2), 125-129.
14. Multani, D. S., Briggs, S. P., Chamberlin, M. A., Blakeslee, J. J., Murphy, A. S., & Johal, G. S. (2003). Loss of an MDR transporter in compact stalks of maize br2 and sorghum dw3 mutants. *Science*, 302(5642), 81-84.
15. O'donoghue, L. S., & Bennett, M. D. (1994). Durum wheat haploid production using maize wide-crossing. *Theoretical and Applied Genetics*, 89, 559-566.
16. Riera-Lizarazu, O., & Mujeeb-Kazi, A. (1990). Maize (*Zea mays* L.) mediated wheat (*Triticum aestivum* L.) polyhaploid production using various crossing methods. *Cereal Research Communications*, 339-345.
17. Salamini, F. (2003). Hormones and the green revolution. *Science*, 302(5642), 71-72.
18. Suenaga, K., & Nakajima, K. (1989). Efficient production of haploid wheat (*Triticum aestivum*) through crosses between Japanese wheat and maize (*Zea mays*). *Plant Cell Reports*, 8, 263-266.
19. Ushiyama, T., Kuwabara, T., & Yoshida, T. (2007). Effects of various phytohormones on haploid wheat production in wheat x maize crosses. *Plant production-Production science* *Science*, 10(1), 36-41.
20. Warchoń, M., Skrzypek, E., Nowakowska, A., Marcińska, I., Czyczyło-Mysza, I., Dziurka, K., ... &Cyganek, K. (2016). The effect of auxin and genotype on the production of *Avena sativa* L. doubled haploid lines. *Plant growth-Growth regulation* *Regulation*, 78, 155-165.
21. Weyen, J. E. N. S. (2009). Barley and wheat doubled haploids in breeding. *Advances in haploid production in higher plants*, 179-187.

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Comment [p2]: Incomplete reference. Include volume and issue.

Formatted: Font: Italic

Formatted: Font: Italic

Comment [p3]: This book should be well referenced with editors included.