

### Original Research Article

## **Effect of yeast and different levels of sugar on Indian olive (*Elaeocarpus serratus* L.) cider**

### **ABSTRACT**

The study was conducted in Completely Randomized Design (CRD) with 8 treatments replicated three times. The treatments were T1 (Indian olive 500g +250ml water +150g sugar), T2 (Indian olive 500g +250ml water +200g sugar), T3 (Indian olive 500g +250ml water +250g sugar), T4 (Indian olive 500g +250ml water +300g sugar), T5 (Indian olive 500g +250ml water +150g sugar +1.5g yeast), T6 (Indian olive 500g +250ml water +200g sugar +1.5g yeast), T7 (Indian olive 500g +250ml water +250g sugar +1.5g yeast) and T8 (Indian olive 500g +250ml water +300g sugar +1.5g yeast). Total soluble solids, pH, and specific gravity decreased while alcohol content, acidity, and sensory qualities increased with increasing length of fermentation. From the above treatments, it is concluded that treatment T5 was found superior in respect of parameters like total soluble solids, pH, acidity, alcohol content, and specific gravity. Whereas color and appearance, aroma, taste, and overall acceptability were recorded in T6. In terms of gross return, net return, and cost-benefit ratio, treatment T6 was recorded highest. Since Indian olive contains numerous nutritional and medicinal properties and is still underutilized, the production of cider from this fruit can help to reduce post-harvest losses. This study showed that acceptable cider can be produced from Indian olive using yeast (*Saccharomyces cerevisiae*).

**Keywords:** Cider, Indian olive, *Saccharomyces cerevisiae*, Sugar, Fermentation

### **INTRODUCTION**

*Elaeocarpus serratus* (L.) commonly known as Indian olive or wild olive or Ceylon olive is a subtropical fruit tree belonging to the family Elaeocarpaceae. It is indigenous to Sri Lanka and the fruit tree is found in some parts of North Eastern states of India. It requires warmer conditions and is sensitive to frost and large fluctuations in temperatures. It is a large, evergreen, drought-tolerant, perennial broad-leaved tree with a large spreading crown. It attains a height of about 50-200 ft., 5-10 m in canopy spread, blooms during September-October, and fruiting occurs during January. Fruits are ovoid or globose berries of

4-5 cm long, weighing  $3.58 \pm 0.09$  g and a pulpy pericarp encloses a stony seed. Depending on the size of the fruit, the pulp weighs  $2.10 \pm 0.04$  g whereas the seed weighs  $1.47 \pm 0.07$  g.

*Elaeocarpus E. serratus* is one of the underutilized fruit species in India and is found and used extensively in West Bengal as Jalpai, Assam as Zolphai, and Manipur as Chorphon. The consumption of ~~this~~ these fruits has become increasingly important due to their potential health effects. Although, it is underutilized fruit species, they have the potential ~~for~~ ~~to~~ ~~contribution~~ ~~contribute~~ to food security, health and nutrition, income generation, and environmental services (Pushpakumara *et al.*, 2007). The ripened fruits stand out not only for ~~its~~ ~~their~~ unique sour-bitter taste but also for ~~its~~ ~~their~~ amazing nutritional and medicinal properties. In folk medicine, various parts of this plant have been extensively used in the treatment of stress, anxiety, depression, nerve pain, epilepsy, migraine, lack of concentration, asthma, hypertension, arthritis, and liver diseases. Apart from the raw consumption of fruit, the fruit pulp is used for the preparation of value-added products such as jam, jelly, candy, pickles, and cider/wine.

Being an underutilized fruit, many fruits are wasted as there is little knowledge about postharvest management of this fruit. Since it yields several useful items including quality fodder, firewood, and edible fruits, proper study on this fruit could minimize its wastage and maximize income generation.

Food processing is an important method to ~~minimise~~ ~~minimize~~ postharvest loss and improve linkages between industry and agriculture. The loss can be ~~minimised~~ ~~minimized~~ by converting the surplus food into various value-added products, like fermented and unfermented beverages. Fermentation is a potential tool in the development of new products from fruits with modified physio-chemical and sensory qualities, especially flavour and nutritional components (Dubey, 2002).

In most parts of the world, cider is an alcoholic beverage that is the result of fermentation made from pressed apple juice, effectively following the same process as wine. Most cider production processes rely on various strains of *Saccharomyces cerevisiae* that allow rapid and reliable fermentation, reduce the risk of sluggish or stuck fermentation, and prevent microbial contamination (Romano *et al.*, 2003). Yeasts are the prominent organisms involved in cider production and determine several characteristics of the cider, including the flavour, through a range of mechanisms and activities (Fleet, 2003).

Formatted: Highlight

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States), Highlight

Formatted: Font: (Default) Times New Roman, 12 pt, Italic, English (United States)

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States), Highlight

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States), Highlight

~~Fermentation~~ The fermentation of cider and the fermentation of wine are similar mechanisms. Alcoholic fermentation is ~~characterised~~ characterized by the conversion of sugar into ethanol by yeast, especially the ~~Saccharomyces~~ *S. cerevisiae* strain of fungi used for alcoholic fermentation. The yeast eats all the sugar inside the vat of cider and turns it into alcohol and carbon dioxide. Once the whole sugar inside the cider is converted, the yeast dies due to a lack of food, and fermentation is stopped. Temperature is the most important factor for the fermentation process; temperatures ranging from 4 to 16°C are beneficial for cider making. After the completion of fermentation, the fermented cider is made, which contains about 4 to 6 percent alcohol. Before a second fermentation, the addition of sugar or extra fruit juice increases the ethanol content of the beverage and improves the quality of the cider.

Formatted: Font: (Default) Times New Roman, 12 pt, Italic, English (United States)

Cider is a pleasant, refreshing, thirst-quenching, and hygienic beverage. It is also the most nutritive and health-giving of all fermented drinks. Besides, it is said to possess remarkable therapeutic properties, like the prevention of stone formation in the bladder on account of its diuretic properties (Chaudhary, 1969).

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States),

*Elaeocarpus serratus* L. cider, being a fruit-based fermented and undistilled product, contains most of the original nutrients present in it. Apart from the original nutrients, the nutritive value of Indian olive cider is increased due to the release of amino acids and other nutrients from the yeast during fermentation. Depending on the alcohol content, 1-5 percent of cider is considered soft cider, and 7-12 percent is considered hard cider. Research shows that Indian olive cider is nutraceutical in nature, curing various ailments naturally. It is rich in antioxidants, which delay the ageing of cells, cure diabetes and insomnia, act as anti-inflammatory agents, and reduce the risk of cardiovascular diseases, cancer, Parkinson's, and Alzheimer's diseases. Therefore, an experiment has been carried out with the aim of studying the effect of yeast and different levels of sugar on the physio-chemical properties and quality of Indian olive cider and estimating the economics of various treatments.

Formatted: Font: (Default) Times New Roman, 12 pt, Italic, English (United States)

## MATERIALS AND METHODS

The study was conducted in a Completely Randomized Design (CRD) with 8 treatments replicated thrice. The treatments were T1 (Indian olive 500g +250ml water +150g sugar), T2 (Indian olive 500g +250ml water +200g sugar), T3 (Indian olive 500g +250ml water +250g sugar), T4 (Indian olive 500g +250ml water +300g sugar), T5 (Indian olive 500g +250ml water +150g sugar +1.5g yeast), T6 (Indian olive 500g +250ml water +200g sugar

+1.5g yeast), T7 (Indian olive 500g +250ml water +250g sugar +1.5g yeast) and T8 (Indian olive 500g +250ml water +300g sugar +1.5g yeast).

### Raw Materials and Preparation of Cider


The fresh and ripened fruit of Indian olive was procured from Nungshong village of Ukhrul district, Manipur. The fruits that are free from diseases and pests were selected. Completely rotten fruits were discarded and rotten parts of the fruits were removed. The selected fruits were washed thoroughly in clean tap water to remove dust, dirt, and other undesired materials adhering to the fruits. The selected fruits were cut into pieces with a stainless knife and 250 ml of water was added in all the treatments. The “must” was sieved using muslin cloth of pore size 0.8 mm.


### Yeast and Inoculum Preparation

*Saccharomyces cerevisiae* was obtained from the local market. The inoculum was prepared by inoculating 1.5g brewer’s yeast was added in 10 ml of lukewarm water in a beaker and stirred gently. The activated yeast was added to the pulp for further fermentation.

### Preparation of Cider

Cider was prepared with different concentrations of sugar. The steps of preparation are given below.





**Fig.1. Flowchart of Indian olive cider preparation**

**Table 1. Meteorological data during the experimental period (January 2023- March 2023)**

Months	Time span	Temperature (°C)		Relative humidity (%)		Rainfall (mm)
		Max	Min	Max	Min	
January	1 <sup>st</sup> week	25	5.4	97	57	0
	2 <sup>nd</sup> week	24.6	5.2	97	47	0
	3 <sup>rd</sup> week	23.6	6	92	56	0
	4 <sup>th</sup> week	28.4	7.6	95	49	8.2
February	1 <sup>st</sup> week	25.2	10	88	38	0
	2 <sup>nd</sup> week	30.2	13.2	84	37	0
	3 <sup>rd</sup> week	31	13.6	83	42	0
	4 <sup>th</sup> week	34.2	14.2	78	39	0
March	1 <sup>st</sup> week	35.2	13.2	78	37	0
	2 <sup>nd</sup> week	35.6	13.6	80	38	0
	3 <sup>rd</sup> week	33.2	13	85	39	18.2
	4 <sup>th</sup> week	35.2	16.2	84	37	0

### Determination of Physio-Chemical Parameters

The physio-chemical changes that the cider underwent during production and storage were examined. The  $pH$  of the must  $pH$  was determined using AOAC, (2004) procedure where a

Formatted: Highlight

digital pH meter was used to determine the product's pH, while a hand-held refractometer was used to measure TSS and the results were expressed as degree brix (°B) (AOAC, 2000), titratable acidity was measured by using phenolphthalein as an indicator (AOAC, 2000) to titrate 10 ml of an aliquot against a standard solution of 0.1 N NaOH. ~~Appearance-The appearance~~ of light pink color was taken as an endpoint, and a hydrometer was used to measure alcohol content and specific gravity (Triple scale Hydrometer). The product was also examined for color and appearance, taste, aroma, and overall acceptability using a 9-point Hedonic scale with a panel of 5 experts.

## RESULTS AND DISCUSSIONS

The experiment entitled “Effect of yeast and different levels of sugar on Indian olive (*Elaeocarpus serratus* L.) cider” was undertaken in the Post-Harvest Laboratory, Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during the year 2022-2023. The results of the investigation regarding ~~the~~ production of cider from Indian olive ~~have been are~~ presented here (Tables 2 to 4).

~~Completely A completely~~ randomized block design ~~(CRBD)~~ by Panse and Sukhtme's analysis of variance (ANOVA) approach was used for the statistical analysis. Using the critical difference (C. D. at 5%) threshold of significance, the overall significance of differences between treatments was examined. A window-based computing tool called OPSTAT was used to statistically ~~analyse-analyze~~ the results (Sheoran, 2004).

Formatted: Highlight

### Total Soluble Solids (°Brix)

Changes in ~~the~~ TSS of Indian olive cider during fermentation ~~is-are~~ presented in Table 2. ~~The~~ total soluble solids of cider, at the end of ~~the~~ fermentation period, is an important quality parameter and an ~~indicative-indicator~~ of the stability and completeness of fermentation. The statistical analysis ~~has showed-shown~~ that there were significant differences in the TSS of all the treatments during 90 days ~~of~~ storage. The total soluble solids content ~~on-in~~ Indian olive cider showed decreasing trend in all the treatments. It is possible that the fermentation of sugars into alcohol by the action of yeast during storage is what caused the decrease in ~~the~~ total soluble solids content of Indian olive cider. The general decrease in TSS was a function of time and was undoubtedly caused by the yeast fermenting the sugar. This is typical cider fermentation behavior for any alcoholic fruit juice fermentation. The decrease in ~~the~~ TSS content of cider indicates the conversion of sugar into alcohol by yeast during fermentation.

The results found are similar ~~with to~~ the findings of (~~Idise et al., 2010~~) in pomegranate wine and ~~in banana wine (Isitua et al., 2010) in banana wine~~. The maximum score (29.86) of TSS (°brix) was observed in treatment T8 (500g Indian olive + 250ml water +300g sugar +1.5g yeast) and ~~the~~ lowest score (5.5) was observed in treatment T5 (500g Indian olive + 250ml water +150g sugar +1.5g yeast).

Formatted: Highlight

Formatted: Highlight

## pH

The statistical analysis has ~~showed show~~ that there were significant differences in pH during 90 days ~~of~~ storage. As fermentation time increased, the pH steadily decreased. The changes in pH are due to the effect of yeast strain and fermentation time. (~~Dhakane et al., 2016~~) encoded that low pH contributes to the stability of alcoholic beverages over time. The decrease in pH with ~~the~~ increase in acidity of cider may be due to the formation of hydrogen ions by the action of yeast. The results ~~shown obtained~~ are similar ~~with to~~ the findings of (~~Won et al., 1993~~), (~~Sigler et al., 1981~~) and (~~Castrillo et al., 1995~~). The lowest score of pH (3.21) was found in treatment T5 (500g Indian olive + 250ml water +150g sugar +1.5g yeast) whereas the maximum score (4.69) was found in treatment T8 (500g Indian olive + 250ml water +300g sugar +1.5g yeast).

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

## Titrateable Acidity (%)

~~when~~ ~~When~~ the pH dropped, it was discovered that the titrateable acidity increased, indicating that more organic acids were being generated as the microbes used the substrates. Acidity plays a vital role in determining cider quality by aiding the fermentation process and enhancing the overall characteristics and balance of the cider. ~~Lack~~ ~~A lack~~ of acidity will mean ~~a~~ poor fermentation (~~Berry, 2000~~). Titrateable acidity increased as fermentation progressed. During the storage period of 90 days, as shown in Table 2, the ~~increasing increase~~ in acidity of Indian olive cider may possibly be due to ~~the~~ production of certain organic acids such as citric, mucic, and tartaric acids by yeast. The results shown are similar ~~with to~~ the findings of (~~Ngaihling and Topno, 2023~~) in pomegranate cider. The lowest score (0.35%) of acidity (%) was observed in treatment T7 (500g Indian olive + 250ml water +250g sugar +1.5g yeast) and ~~the~~ maximum score (0.83%) was observed in treatment T5 (500g Indian olive + 250ml water +150g sugar +1.5g yeast).

Formatted: Highlight

## Alcohol content (%)

With the increase in fermentation time, alcohol concentration also increases. During 90 days of storage as shown in Table 2, the increase in alcohol content of Indian olive cider with yeast and different levels of sugar during storage may be ~~possibly-possible~~ due to the variation in performance of the sugar affecting the fermenting ability, hence the varied alcohol production. The results ~~shown of the present investigation~~ are similar ~~with to~~ the findings of ~~(Wanapu et al., 2004)~~ in rose apple cider, ~~and in banana wine~~ (Beera et al., 2013) ~~in banana wine~~. The highest score (~~10.72%~~) of alcohol content (~~(%)~~) was found in treatment T5 (500g Indian olive + 250ml water +150g sugar +1.5g yeast) and the lowest score (4.57%) was found in treatment T8 (500g Indian olive + 250ml water +300g sugar +1.5g yeast).

Formatted: Highlight

### Specific Gravity

As shown in Table 2, the specific gravity decreases as the fermentation time of cider increases. The decrease in ~~the~~ specific gravity of Indian olive cider during storage may be possibly due to ~~the~~ concentration of yeast used and different levels of sugar in the cider production. It is also reported that *Saccharomyces cerevisiae* reduces ~~the~~ specific gravity of fruit juices during fermentation. The ~~present~~ results ~~found~~ are similar ~~with to~~ the findings of ~~(Jarvis, 2014)~~ in apple cider, and ~~in Mahua wine~~ (Yadav et al., 2009) ~~in Mahua wine~~. The lowest score (~~1.021~~) of specific gravity (~~1.021~~) was observed in treatment T5 (500g Indian olive + 250ml water +150g sugar +1.5g yeast) and the maximum score (1.128) was observed in treatment T8 (500g Indian olive + 250ml water +300g sugar +1.5g yeast).

Formatted: Highlight

### Organoleptic Evaluation

As shown in Table 3, the highest sensory scores for organoleptic evaluation such as color and appearance, aroma, taste, and overall acceptability ~~was were~~ observed in treatment T6 (500g Indian olive + 250ml water +200g sugar +1.5g yeast) with 8.14, 8.20, 8.69 and 8.43 which indicates ~~that~~ the judges approved ~~of~~ it.

**Table 2. Physio-chemical parameters of Indian olive cider during storage**

Treatment symbols	Treatment details	Total Soluble Solids (°Brix)				pH				Acidity (%)				Alcohol (%)			Specific Gravity			
		Initial	30DAS	60DAS	90DAS	Initial	30DAS	60DAS	90DAS	Initial	30DAS	60DAS	90DAS	30DAS	60DAS	90DAS	Initial	30DAS	60DAS	90DAS
T1	Indian olive + water + 150 g Sugar	19.2	12.33	8.3	6.23	4.39	4.31	4.03	3.36	0.51	0.62	0.7	0.74	6.83	9.01	10.57	1.071	1.051	1.032	1.024
T2	Indian olive + water + 200 g Sugar	20.36	13.33	9.21	6.86	4.43	4.4	4.13	3.57	0.47	0.54	0.62	0.69	6.3	8.74	10.45	1.084	1.054	1.037	1.027
T3	Indian olive + water + 250 g Sugar	23.13	14.47	9.76	7.46	4.64	4.45	3.95	3.37	0.44	0.5	0.53	0.61	4.78	7.74	8.71	1.097	1.059	1.038	1.029
T4	Indian olive + water + 300 g Sugar	26.96	17.26	10.86	7.73	4.67	4.47	4.12	3.66	0.36	0.46	0.51	0.55	4.55	6.73	7.53	1.116	1.071	1.044	1.031
T5	Indian olive + water + 150 g Sugar +1.5 g yeast	15.53	10.3	7.8	5.5	4.35	4.21	3.79	3.21	0.56	0.63	0.73	0.83	7.49	9.56	10.72	1.063	1.04	1.031	1.021
T6	Indian olive + water + 200 g Sugar +1.5 g yeast	50.6	12.56	8.43	6.36	4.61	4.42	4.08	3.43	0.48	0.56	0.61	0.64	6.43	8.62	9.69	1.084	1.051	1.032	1.024
T7	Indian olive + water + 250 g Sugar +1.5 g yeast	28.93	19.1	13.7	10.56	4.62	4.41	4.1	3.45	0.35	0.42	0.49	0.56	4.47	6.56	7.68	1.124	1.071	1.055	1.042
T8	Indian olive + water + 300 g Sugar+1.5 g yeast	29.86	20.5	14.56	12.36	4.69	4.44	4.11	3.81	0.43	0.45	0.48	0.53	4.57	6.42	7.35	1.128	1.081	1.058	1.05
F-test		S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
S.E.(d)		0.289	0.172	0.149	0.139	0.067	0.61	0.54	0.52	0.01	0.009	0.009	0.009	0.14	0.09	0.069	0.002	0.001	0.001	0.001
C <sub>1</sub> D <sub>2</sub> at 0.5%		0.617	0.367	0.319	0.298	0.144	0.13	0.115	0.11	0.021	0.019	0.02	0.02	0.3	0.192	0.147	0.004	0.002	0.002	0.001
C <sub>2</sub> V <sub>1</sub> (%)		1.532	1.218	1.653	2.665	1.442	1.811	1.848	1.888	2.713	2.089	1.981	1.736	3.024	1.385	0.928	1.19	0.097	0.069	0.101

\*In all treatments 500g Indian olive fruit and 250ml of water was reused uniformly

Table 3. ~~Organoleptic~~ The organoleptic score of Indian olive cider during storage

Treatment symbols	Treatment details	Colour and appearance			Aroma			Taste			Overall acceptability		
		30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS
T1	Indian olive + water + 150 g Sugar	5.11	5.84	6.73	5.73	6.13	6.69	6.08	6.38	6.97	6	6.45	7.13
T2	Indian olive + water + 200 g Sugar	5.4	6.65	7.6	6.22	6.72	7.63	6.52	7.25	7.84	6.02	6.5	7.66
T3	Indian olive + water + 250 g Sugar	4.78	5.69	6.15	5.27	5.75	6.61	5.42	5.81	6.28	5.76	5.94	6.82
T4	Indian olive + water + 300 g Sugar	4.28	5.61	5.98	5.14	5.47	6.25	4.77	5.12	5.64	4.83	5.08	6.58
T5	Indian olive + water + 150 g Sugar +1.5 g yeast	6.28	7.81	8.19	6.79	7.14	7.61	6.28	7.62	8.14	6.64	6.92	7.93
T6	Indian olive + water + 200 g Sugar +1.5 g yeast	6.38	7.85	8.14	7.13	7.61	8.2	6.59	8.18	8.69	7.22	7.64	8.43
T7	Indian olive + water + 250 g Sugar +1.5 g yeast	5.71	6.72	7.68	5.95	6.2	6.75	6.11	6.9	7.21	6.58	7.11	7.72
T8	Indian olive + water + 300 g Sugar+1.5 g yeast	5.51	6.28	6.62	5.4	5.89	6.49	5.56	6.2	6.71	5.54	6.19	6.72
F-Test		S	S	S	S	S	S	S	S	S	S	S	S
S.E.(d)		0.044	0.060	0.032	0.041	0.026	0.031	0.066	0.031	0.025	0.078	0.030	0.025
C <sub>2</sub> D <sub>2</sub> at 0.5%		0.094	0.129	0.032	0.088	0.055	0.067	0.141	0.066	0.054	0.166	0.165	0.053
C <sub>2</sub> V <sub>2</sub> (%)		0.995	1.125	0.545	0.851	0.495	0.548	1.361	0.570	0.427	1.565	0.576	0.412

\*In all treatments 500g Indian olive fruit and 250ml of water ~~was~~ were used uniformly

**Table 4. Economics of different treatments and benefit ratio of Indian olive cider**

Treatment no.	Treatment details	Total cost (Rs)	Cider output	Selling rate (Rs/bottle)	Gross return (Rs)	Net return (Rs)	Benefit-cost ratio
T1	Indian olive + water + 150 g Sugar	189.17	3	260	780	212.5	1.44
T2	Indian olive + water + 200 g Sugar	191.67	3	270	810	234.9	1.41
T3	Indian olive + water + 250 g Sugar	194.17	3	250	750	167.5	1.29
T4	Indian olive + water + 300 g Sugar	196.67	3	220	660	69.9	1.11
T5	Indian olive + water + 150 g Sugar +1.5 g yeast	195.92	3	290	870	282.2	1.48
T6	Indian olive + water + 200 g Sugar +1.5 g yeast	198.42	3	300	900	304.7	1.51
T7	Indian olive + water + 250 g Sugar +1.5 g yeast	200.92	3	280	840	237.2	1.39
T8	Indian olive + water + 300 g Sugar+1.5 g yeast	203.42	3	230	690	79.7	1.13

\*In all treatments 500g Indian olive fruit and 250ml of water ~~was~~ were used uniformly

## CONCLUSION

Based on [the](#) findings of the present investigation it is concluded that treatment T5 (Indian olive 500g + 250ml water + 150g sugar + yeast 1.5g) was found superior in terms of Total Soluble Solids (5.50 °Brix), pH (3.21), Acidity (0.83%), Alcohol (10.72%) and Specific gravity (1.021). Whereas, in terms of Colour and appearance (8.19), Aroma (8.20) Taste (8.69), and Overall acceptability (8.43), treatment T6 (Indian olive 500g +250ml water +200g sugar +1.5g yeast) was found superior. Similarly, treatment T6 showed the highest BC ratio (1.51) [followed by T5 \(1.48\)](#).

## REFERENCES

[Akin](#), H., Brandam, C., Meyer, X.M. and Strehaiano, P.A. (2008). Model for pH determination during alcoholic fermentation of a grape must by *Saccharomyces cerevisiae*. *Chemical Engineering Processing Process Intensification*, 47:1986-1993.

Formatted: Highlight

[Anderson](#), C. and Badrie, N. (2005). Physio-chemical quality and consumer acceptance of guava wines. *Journal of Food Science Technology*. 42(3):223-226.

Formatted: Highlight

[Attri](#), B. L. (2009). Effect of initial concentration on the physio- chemical characteristics and sensory qualities of cashew apple wine. *Indian Journal of Natural Products and Resources*, 8:374-379.

Formatted: Highlight

[Banjare](#), A. and Patel, D. (2020). Preparation of apple cider: Effect of apple Juice concentration on physio-chemical composition and sensory qualities of apple cider. *Journal of Pharmacognosy and Phytochemistry* 9(3):75-77.

Formatted: Highlight

[Beech](#), F.W. (1972). Cider making and Cider research: A review. *Journal of the Institute of Brewing*, 78(6):477-491.

Formatted: Highlight

Beera, V., Mangam, R.B. and Enthoti, K. (2013). Mango wine: Standardisation and screening of cultivars using different strains of *Saccharomyces cerevisiae*. *Asian Journal of Multidisciplinary Studies*, 1(4):35-42.

[Chowdhury](#), J.F. (2019). Physicochemical, microbial, and sensory properties analysis of olive jelly prepared from different maturity stages of olive. *Chattogram Veterinary and Animal Sciences University Chattogram*.

Formatted: Highlight

Jawahar, A., Kotecha, P.M. and Kadam, S.S. (2001). Wine preparation from guava juice. *Beverage and Food World*. 28(1):37-38. Formatted: Highlight

Jayashree, I., Geetha, D.H. and Rajeswari, M. (2014). Evaluation of antimicrobial potential of *Elaeocarpus serratus* L. *International Journal of Pharmaceutical Sciences and Research*, 51:67-72. Formatted: Highlight

Khoisnam, N., Thokchom, R., Maisnam, R. and Sapna, J. (2020). Ethnobotanical uses of *Elaeocarpus serratus*. *European Journal of Molecular and Clinical Medicine*, 7(7):79-88. Formatted: Highlight

Kumar, Y.S., Varakumar, S. and Reddy, O.V.S. (2012). Evaluation of antioxidant and sensory properties of mango (*Mangifera indica* L.) wine. *Journal of Food*, 10(1):12-20. Formatted: Highlight

Malleshwari, P., Faseela, V.A. and Shanmugasundaram, P. (2022). Traditional Uses, Phytochemical and Pharmacological Activity of *Elaeocarpus Serratus*. *Journal of Pharmaceutical Negative Results*, 13(8):1837-40. Formatted: Highlight

Ngaihling, C. and Topno, S.E. (2023). Effect of different levels of local yeast in pomegranate (*Punicagranatum* L.) cider. *International Journal of Plants and Soil Science*, 35(14):80-88.

Ningli, Q., Lina, M., Liuji, L., Xiao, G. and Jianzhi, Y. (2017). Production and Quality Evaluation of Pineapple Fruit Wine. *International Global on Renewable Energy and Development*. Formatted: Highlight

Pooja, N., and Kocher, G.S. (2015). Fermentative Production of Guava wine (*Psidium guajava* L.) Using *S. cerevisiae*. *Current Nutrition & Food Science*, 11(1):21-30. Formatted: Highlight

Sahu, U.C., Panda, S.K., Uma, B., Mohapatra, U.B. and Ray, R.C. (2012). Preparation and evaluation of wine from Tendu (*Diospyros melanoxylon* L.) fruits with antioxidants. *International Journal of Food Fermentation Technology*, 2(2):167-178. Formatted: Highlight

Song, J., Yuchen, S., Xi, C., Tian, W. and Jianbo, L. (2015). The olive fruit wine research on the fermentation process and aromatic ingredients. *Atlantis press Journal*:603-608. Formatted: Highlight

Sriti, K., Manjari, S.K. and Kanti, C.C. (2011). Correlation between phytochemical screening, antibacterial and aromatic activities of *Elaeocarpus serratus* L. *International Journal of Pharmacy*. 2(9)133-136. Formatted: Highlight

Walker, G.M., and Stewart, G.G. (2016). *Saccharomyces cerevisiae* in the Production of Fermented Beverages. *Beverages*, 2(4):30. Formatted: Highlight

Yadav, P., Garg, N. and Diwedi, D. H. (2009). Effect of location of cultivar, fermentation temperature and additives on the physio-chemical and sensory qualities on Mahua (*Madhuca indica* J. F. Gmel.) wine preparation. *Indian Journal of Natural Products and Resources*, 8(4): 406-418. Formatted: Font: Not Italic  
Formatted: Font: Not Italic

Yao, Q., Gang, H., Yibing, H., Shen, Y. and Gou, X. (2016). Antioxidant activity of olive wine, a by-product of olive mill wastewater. *National Library of Medicine*, 54(10):2276-81. Formatted: Highlight

UNDER PEER REVIEW