


[Original Research Article]

Influence of Phosphorus and Iron on Yield and Economics of Lentil (*Lensculinaris* L.)



ABSTRACT

A field experiment was conducted during *Rabi* 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P.) on the topic “Influence of Phosphorus and Iron on Yield and Economics of Lentil (*Lens culinaris* L.)”, to study treatments consisting of three levels of Phosphorus viz. 20 kg, 30 kg and 50 kg/ha and three levels of Iron viz. 5, 10 and 20 kg/ha. The soil of experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.28%), available N (22.5 kg/ha), available P (19.50 kg/ha) and available K (92 kg/ha). There were 10 treatments each being replicated thrice and laid out in Randomized Block Design. The results revealed that treatment 9 (Phosphorus at 50 kg/ha + FeSO₄ at 20 kg/ha) recorded significantly higher number of pods/plant (113.07), number of seeds/pod (1.93), Test weight (31.20 g), seed yield (1.25 kg/ha), Stover yield (2.07 kg/ha), harvest index (37.67%), gross returns (1,00,266.67 INR/ha), net returns (66,866.71 INR/ha) and B:C ratio (2.00) as compared to other treatments.

Keywords: *Phosphorus, Iron, Yield, Economics.*

Comment [mm1]: Add background, problems, aims, and novelty

Introduction


The family Leguminosae includes lentils (*Lens culinaris* Medic). Legumes like lentils are a healthy food source. It is one of the oldest annual grains and legumes that is grown and consumed the most in the world, and is primarily eaten as dhal. As early as 6000 B.C., lentil was grown in South Western Asia. In addition to having a high concentration of the important amino acids isoleucine and lysine, lentils also include other nutrients such as nutritive fibre, folate, vitamin B₁, and minerals (Rozalet al., 2001). Because it contains a high amount of protein, lentil is also referred to as "poor man's meat." Lentils are also enjoyed equally by all socioeconomic strata in South East Asia (Bhattya, 1988). As a thick soup made from whole grains or split pulses known as "dhal," lentils are frequently consumed. Its seeds can be fried and eaten, while lentil flour is excellent for soups, stew purées, bread and pastries made with cereal, as well as for additions to softness to cuisine (Williams and Singh, 1988).

The Indus subcontinent, which includes Pakistan, the Middle East, and many other nations, uses lentil as a main course, a side dish, or in salads. It is used as a substitute for meat in vegetarian diets in some places. South East Asia is where lentils are most frequently cultivated. The most well-known nations for the production and cultivation of lentil include India, Canada, Turkey, Bangladesh, Iran, China, Nepal, and Syria, among many more countries. On an area of 19.6 thousand hectares, lentils are grown in Pakistan during the winter as the second-most important pulse crop, after chickpea, with an annual production of 9.7 thousand tonnes and an average grain yield of 541 kg ha⁻¹ (GOP, 2013), which is a very low yield and cannot satisfy the demands of the growing population. Due to the spread of various illnesses and weed infestations, the area used to cultivate lentils is gradually decreasing.

Due to its role in root development, stalk and stem strength, flower and seed formation, crop maturity and production, N-fixation, crop quality, and resistance to plant diseases, phosphorus is a crucial component for the successful production of pulses. It is vital to the stimulation

of biological processes such as Legume yield is increased via nodulation, nitrogen fixation, and nutrient uptake in the rhizosphere environment. Application of phosphorus reduces the negative effects of drought on physiological parameters and can increase yield in water situations of stress (Singh et al., 2005). The majority of soils used for producing lentils have low to medium levels of phosphorus that are readily available, therefore they respond well to the suggested quantity of phosphorus fertilizers. Pulses' increased yield is a result of phosphorus application, which also improves the soil's nitrogen content for subsequent non-legume crops that require less nitrogen. However, high rates of phosphorus application cause P fixation because they chelate with iron and aluminium in acidic soils and calcium in alkaline soils, making them unavailable to plants. So, to get the ideal lentil yield, phosphorus needs to be applied in the proper amount. Therefore, phosphorus must be applied in the proper amount to produce the highest possible yield of lentils. In several crops, including lentil, iron (Fe) insufficiency is a prevalent nutritional problem (Erskine et al., 1993). The yield losses of the vulnerable genotypes ranged from 18 to 25%. According to Sakalet al. (1984), the Fe²⁺ concentration of the leaf tissue, as compared to the total iron content, was closely correlated with the symptoms of Fe lack and was determined to be a good index to identify soil where response to Fe attitudes can be expected. Leghemoglobin, ferredoxin, and nitrogenase are all made up of iron. During the process of fixing nitrogen, bacteria have used this element. Some legumes have low nitrogen concentrations in their shoots as a result

of iron deficiency, which typically reduces nodule formation, leghaemoglobin production, and



Nitrogenase activity. The nitrogen fixation processes have been shown to benefit from iron and molybdenum fertilisation, increasing lentil yield and nitrogen status. The ability to actively fix nitrogen depends on the crop's health and an appropriate supply of nutrients, as is well established (**Bra r and Sidhu, 1992**). According to **Lindsay and Norwell (1969)**, the essential level of iron is 4.5 ppm.

Keeping

in view of the above facts, the present experiment was undertaken to find out “**Influence of Phosphorus and Iron on Growth and Yield of lentil (*Lens culinaris* L.)**”

MATERIALS AND METHODS:

The experiment was conducted during *rabi* season of 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). The soil of the field constituting a part of central gangetic alluvium is neutral and deep. The soil of experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.28%), available N (225 kg/ha), available P (19.50 kg/ha) and available K (92 kg/ha). The treatment consists of Phosphorus @ 20 kg, 30 kg and 50 kg/ha and Iron consists 5 kg, 10 kg and 20 kg/ha. The experiment was laid out in RBD with 10 treatments each replicated thrice. The treatment combinations are T1- Phosphorus at 20 kg/ha + Iron at 5 kg/ha, T2- Phosphorus at 20 kg/ha + Iron at 10 kg/ha, T3 - Phosphorus at 20 kg/ha + Iron at 20 kg/ha, T4 - Phosphorus at 30 kg/ha + Iron at 5 kg/ha, T5 - Phosphorus at 30 kg/ha + Iron at 10 kg/ha, T6- Phosphorus at 30 kg/ha + Iron at 20 kg/ha, T7 - Phosphorus at 50 kg/ha + Iron at 5 kg/ha, T8 - Phosphorus at 50 kg/ha + Iron at 10 kg/ha, T9- Phosphorus at 50 kg/ha + Iron at 20 kg/ha, T10- Control N:P:K (20:40:20 Kg/ha). The growth parameters and yield production was recorded at harvest from randomly selected plants in each plot. The data was computed and analysed by following statistical method of **Gomez and Gomez (1984)**.

RESULT AND DISCUSSION

YIELD PARAMETERS

Number of pods/plant

Treatment 9 with (Phosphorus at 50 kg/ha + FeSO₄ at 20 kg/ha) recorded significantly highest Number of pods per plant (113.07). However, treatments with (Phosphorus at 50 kg/ha + FeSO₄ at 10 kg/ha) and (Phosphorus at 50 kg/ha + FeSO₄ at 20 kg/ha) were statistically at par with the (Phosphorus at 50 kg/ha + FeSO₄ at 20 kg/ha).

It might be the reason of moderate plant nutrients availability due to which the plant produce more pods per plant¹ as compare to other treatments and also phosphorus strongly increases the reproduction of the plants i.e. flowering and fruiting (**Ali et al. 2017**). These results were similar with that of (**Saleh, 1976**) and (**Jayapaul et al. 1990**).

Number of seeds/pod

Treatment 9 with (Phosphorus at 50 kg/ha + FeSO₄ at 20 kg/ha) recorded significantly highest Number of seeds per pods (1.93). However, treatments with (Phosphorus at 50 kg/ha + FeSO₄ at 10 kg/ha), (Phosphorus at 50 kg/ha + FeSO₄ at 5 kg/ha), (Phosphorus at 30 kg/ha + FeSO₄ at 20 kg/ha), (Phosphorus at 30 kg/ha + FeSO₄ at 5 kg/ha), (Phosphorus at 30 kg/ha + FeSO₄ at 10 kg/ha), (Phosphorus at 30 kg/ha + FeSO₄ at 20 kg/ha), and (Phosphorus at 20 kg/ha + FeSO₄ at 20 kg/ha) were statistically at par with the (Phosphorus at 50 kg/ha + FeSO₄ at 20 kg/ha).

The increase in number of seeds/pod by the application of molybdenum along with iron may be due to the fact that molybdenum and iron may fix that much amount of nitrogen which was required by the plant to show better performance as molybdenum is related directly to nitrogen fixation by legumes. Result also showed that the molybdenum and iron nutrition had similar effect on lentil. Similar observations were found by **Landgeet et al. (2002)** and **Tahir et al. (2011)** in chickpea.

Comment [mm2]: Add discussion. Compare with previous research

Test weight (g):

Highest test weight (31.20 g) was recorded in Treatment 9 with application of (Phosphorus at 50kg/ha+FeSO₄at20kg/ha),thoughtherewassignificantdifferenceamong thetreatments.

The possible reason for thousand grain weight could be the effect of cell division, phosphoruscontentspresentinsidetheseed aswellastheformationofalbumin andfats(**Aliet al., 2017**).

Seed Yield(kg/ha):

Treatment 9 with (Phosphorus at 50 kg/ha + FeSO₄ at 20 kg/ha) recorded the highest seed yield(1253.33 kg/ha). However, treatment with Phosphorus at 50 kg/ha + FeSO₄at 10kg/ha wasstatisticallyatparwiththe (Phosphorusat50kg/ha + FeSO₄at20kg/ha).

The reason for increasing grain yield is the balance nutrient supply and timely application of thenutrients which enhances the plant growth and the grain yield will be increased. Similar resultssupporthatincreasingphosphoruslevel,thegrainyieldincreasedtoo(**Chaubeyetal.,1999**)and(**Singh et al., 1999**). The increase in seed yield by application of 60 kg P₂O₅ /fed might beassociated with high number of pods /plant, 1000 seed weight and seed yield /plant (**Zeidan,2007**). Similar results were reported by (**Chaubeyet al., 1999**) and (**Singh et al., 1999**) and

theyfoundthatphosphorusfertilizationat50kg/P₂O₅/haincreasedseedyieldascomparedwithcontrol.It is widely known that Fe helps improve the chlorophyll content necessary for photosyntheticactivities. Moreover, Fe and Mo are integral components of nitrogenase enzyme which areessential for symbiotic N₂ fixation (**Nasar et al. 2017**). These findings are supported by

ShariefandSaid(1998)foundthatfoliarapplicationofmicronutrientseitherseparatelyorincombination significantlyimprovedthe grainyieldoflentil.

Stover yield (kg/ha)

Treatment 9 with (Phosphorus at 50 kg/ha + FeSO₄ at 20 kg/ha) recorded the highest stover yield(2073.33 kg/ha). However, treatments with Phosphorus at 50 kg/ha + FeSO₄at 10kg/ha andPhosphorus at 50 kg/ha + FeSO₄ at 5kg/ha were statistically at par with the (Phosphorus at 50kg/ha +FeSO₄at20kg/ha).

The higher stover yield with suitable dose of phosphorus might be contributed by better growthof the plant as expressed in terms of plant height, number of branches per plant, fresh and dryweight of the plant as a result of improved nutrient uptake. These findings were supported by(**Choubeyetal.2013**)and(**Zeidan,2007**)and(**RasoolandSingh(2016)**)inlentil.This mightbedue to increased availability of physiologically active iron (Fe²⁺) in the plant system which inturns affects various physiological functions of plants favourably. Translocation of the same toreproductive structures. Since uptake is the function of seed and straw yield and their nutrientconcentration,therewassignificantimprovementinconcentrationofthesenutrientscoupledwithseedandstrawyield(**Meenaetal.2013**).

Harvest Index(%)

Treatment9with(Phosphorusat50kg/ha+FeSO₄at20kg/ha)recordedthehighestharvestindex(37.67%).

Each increment of P from (25 to 75 kg ha⁻¹) gave superior HI value of lentil (**Fatima et al., 2013**), however, the low HI at low level of P might be due to poor development of plant.

ECONOMICS

Cost of cultivation varied due to Phosphorus and different levels of Iron. Highest cost of cultivation was seen in treatment 9 with Phosphorus at 50 kg/ha + FeSO₄ at 20 kg/ha (31,649.96 INR/ha) and lowest was seen in the control (28,549.96 INR/ha).

Gross returns varied due to Phosphorus and different levels of Iron on Yield and Yield components of Lentil. Highest Gross returns were seen in treatment 9 with Phosphorus at 50 kg/ha + FeSO₄ at 20 kg/ha (1,00,266.67 INR/ha) and lowest Gross returns were seen in the treatment control (70,400.00 INR/ha).

Net Returns varied due to Phosphorus and different levels of Iron on Yield and Yield components of Lentil. Highest Net returns were seen in the treatment 9 with Phosphorus at 50 kg/ha + FeSO₄ at 20 kg/ha (66,618.71 INR/ha) and lowest Gross returns were seen in the treatment control (41,850.04 INR/ha).

Highest B: C Ratio was recorded with the treatment 9, Phosphorus at 50 kg/ha + FeSO₄ at 20 kg/ha (2.17) and lowest B: C Ratio was seen in the treatment Phosphorus at 20 kg/ha + FeSO₄ at 20 kg/ha (1.39).

Table 1: Influence of Phosphorus and Iron on Yield attributes and Yield on Lentil.

Sl No.	Treatments	Number of pods/plant	Number of Seeds/Pod	Test weight (g)	Seed yield (kg/ha)	Stover Yield (kg/ha)	Harvest Index (%)
1	Phosphorus at 20 kg/ha + FeSO ₄ at 5 kg/ha	94.80	1.13	27.67	900.00	1533.33	37.02
2	Phosphorus at 20 kg/ha + FeSO ₄ at 10 kg/ha	97.47	1.33	27.73	910.00	1596.67	36.34
3	Phosphorus at 20 kg/ha + FeSO ₄ at 20 kg/ha	97.80	1.47	27.73	936.67	1706.67	35.46
4	Phosphorus at 30 kg/ha + FeSO ₄ at 5 kg/ha	98.67	1.53	27.80	996.67	1793.33	35.73
5	Phosphorus at 30 kg/ha + FeSO ₄ at 10 kg/ha	99.67	1.53	29.40	1033.33	1853.33	35.79
6	Phosphorus at 30 kg/ha + FeSO ₄ at 20 kg/ha	104.47	1.53	29.73	1036.67	1856.67	35.81
7	Phosphorus at 50 kg/ha + FeSO ₄ at 5 kg/ha	109.33	1.53	30.40	1110.00	1930.00	36.50
8	Phosphorus at 50 kg/ha + FeSO ₄ at 10 kg/ha	110.00	1.80	30.47	1153.33	1973.33	36.87
9	Phosphorus at 50 kg/ha + FeSO ₄ at 20 kg/ha	113.07	1.93	31.20	1253.33	2073.33	37.67
10	Control (RDF 20-40-20 NPK/ha)	91.47	1.07	27.60	880.00	1470.00	37.47
	F-Test	S	S	NS	S	S	NS
	SEm(±)	1.74	0.16	1.10	42.26	62.86	1.42
	CD (p=0.05)	5.17	0.46	-	125.55	186.76	-

Table 2: Influence of Phosphorus and Iron on Economics of Lentil.

SlNo.	Treatments	Total cost of cultivation (INR/ha)	Gross Returns (INR/ha)	Net Returns (INR/ha)	B:C ratio
1	Phosphorus at 20kg/ha + FeSO ₄ at 5kg/ha	29099.96	72000.00	42900.04	1.47
2	Phosphorus at 20kg/ha + FeSO ₄ at 10kg/ha	29849.96	72800.00	42950.04	1.44
3	Phosphorus at 20kg/ha + FeSO ₄ at 20kg/ha	31349.96	74933.33	43583.37	1.39
4	Phosphorus at 30kg/ha + FeSO ₄ at 5kg/ha	29199.96	79733.33	50533.37	1.73
5	Phosphorus at 30kg/ha + FeSO ₄ at 10kg/ha	29949.96	82666.67	52716.71	1.76
6	Phosphorus at 30kg/ha + FeSO ₄ at 20kg/ha	31449.96	82933.33	51483.37	1.64
7	Phosphorus at 50kg/ha + FeSO ₄ at 5kg/ha	29399.96	88800.00	59400.04	2.02
8	Phosphorus at 50kg/ha + FeSO ₄ at 10kg/ha	30149.96	92266.67	62116.71	2.06
9	Phosphorus at 50kg/ha + FeSO ₄ at 20kg/ha	31649.96	100266.67	68616.71	2.17
10	Control (RDF 20-40-20 NPK/ha)	28549.96	70400.00	41850.04	1.47

CONCLUSION

It is concluded that application of 50 kg/ha Phosphorus with 20 kg/ha Iron (treatment 9) recorded higher seed yield and maximum net returns and benefit cost ratio in lentil crop.

REFERENCES

1. Brar, J. S. and Sidhu A S. (1992). Effect of Rhizobium Inoculation under Different Levels of Phosphorus and Molybdenum on N, P, and Mo contents of straw and Seed of Moong (*Phaseolus aureus* Roxb.).
2. Bhatta R. S. 1988. Composition and quality of lentil (*Lens culinaris* Medik): a review. *Can. Inst. Food Sci. Technol. J.* 21:144-160.
3. Chaubey AK, Kushik MK & Singh SB (1999). Response of lentil to phosphorus and zinc sulphate nutrition. *Crop Res Hasar* 17:309-12.
4. Datta, S.K., Sarkar, M.A.R. and Uddin, F.M.J. (2013). Effect of variety and level of phosphorus on the yield and yield components of lentil. *International Journal of Agricultural Research Innovation Technology* 3:78-82.
5. Erskine, W., Saxena, N. P. and Saxena, M. C. (1993). Iron deficiency in lentil: Yield loss and geographic distribution in germplasm collection. *Plant Soil* 151:249-254.
6. Fatima, K., Hussain, N., Pir, F.A. and Mehdi, M. (2013). Effect of nitrogen and phosphorus on growth and yield of lentil (*Lens culinaris*). *Elixir Applied Botany* 57:14323-14325.
7. GOP. 2013. Economic Survey of Pakistan, 2012-13. Finance Division, Economic Advisor's Wing, Islamabad, 22-22.
8. Jayapaul P & Ganesaraja V (1990). Studies on response of soybean varieties to nitrogen and phosphorus. *Indian Journal Agriculture* 35:329-330.
9. Kuldeep, Kumawat, P.D., Bhadu, V., Sumeriya, H.K. and Kumar, V. (2018). Effect of iron and zinc nutrition on growth attributes and yield of chickpea (*Cicer arietinum* L.). *International Journal of Current Microbiology and Applied Sciences*, 7(8):2837-2841.
10. Landge, S. K., S. U. Kakade, P. D. Thakare., A. P. Karunakar and D. J. Jiotode. (2002). Response of soybean to nitrogen and phosphorus. *Pakistan Journal Genotypes Sciences* 3(3):653-655.

Comment [mm3]: Add references using limited 5 years ago

11. Lindsay, W.L. and Norwell, W.A. (1969). Development of a DTPA micronutrient soil test. *Soil Science Annual. Procedures* **35**(1):600-602.
12. Meena, K. K., Meena, R. S., Kumawat, S. M. 2013, Effect of sulphur and iron fertilization on yield attributes, yield and nutrient uptake of mungbean (*Vigna radiata*) *Indian Journal of Agricultural Sciences* **83**(4):472-6.
13. Nasar, J., Shah, Z. 2017. Effect of iron and molybdenum on yield and nodulation of lentil. *ARN Journal of Agricultural and Biological Science* **12**(11).
14. Prasad, U. K., Prasad, T. N., Mishra, R. B. 1995. Effect of irrigation and phosphorus on yield, moisture use and phosphorus uptake of late-sown lentil (*Lens culinaris* L.) in calcareous soil. *Indian Journal of Agricultural Sciences* **65**(3):178-81.
15. Rasheed, M., Jilani, G., Shah, I.A., Najeeb, U. and Iqbal, T. (2010). Improved lentil production by utilizing genetic variability in response to phosphorus fertilization. *Soil and Plant Sciences*. **60**:485-493.
16. Rozaan P., Kuo Y. H and Lambein F. 2001. Amino acids in seeds and seedlings of the genus *Lens*. *Phytochemistry*. **58**:281-289.
17. Sakal, R., Singh, B.P. and Singh, A.P. (1984) Determination of threshold value of iron in soils and plants for the response of rice and lentil to iron application in calcareous soil. *Plant and Soil* **82**:141-148.
18. Saleh SA (1976). Response of six soybean varieties to phosphorus fertilizer at wadi Jizan region, Saudi Arabia. *Annals Agricultural Sciences Moshtohor* **5**:3-10.
19. Sharief A. E and Said El. M. 1998. Response of lentil productivity to phosphorus fertilizer levels and some micronutrients. Proc. 8th Conf. Agron., Suez Canal University, Ismailia, Egypt, 28-29, November. pp.326-34.
20. Singh OB, Shama HB & Singh VK (1999). Effect of nitrogen, phosphorus and rhizobium culture on yield and yield attributes of lentil under dry land conditions. Dept Agron G.B Pant University Agri & Tech Pantnagar Distt. Udham Singh Nagar 263145, India. *Ind J Pulses Research* **12**:260-2.
21. Singh, K.K., Srinivasarao, Ch. and Ali, M. (2005). Root growth, nodulation, grain yield, and phosphorus use efficiency of lentil as influenced by phosphorus, irrigation, and inoculation. *Commun. Soil Science and Plant Anal.* **36**:1919-1929.
22. Tahir, M., A. Ali, N. Aabidin, M. Yaseen and H. Rehman. (2011). Effect of molybdenum and seed inoculation on growth, yield and quality of mungbean. *Genotypes and Environment*. **2**(2):37-40.
23. Togay, Y., Togay, N., Dogan, Y. (2008) Research on the effect of phosphorus and molybdenum applications on the yield and yield parameters in lentil (*Lens culinaris* Medic.) *African Journal of Biotechnology* **7**(9):1256-1260.

- 24. Trivedi, A. K., Hemantaranjan, A. and Pandey, S. K. 2011.** Iron application may improve growth and yield of soybean. *Indian Journal of plant Physiology*, 16(3/4): 309-313.
- 25. Williams P. Cand Singh U. 1988.** Quality screening and evaluation in pulse breeding. J. In: *World Crops: Cool Season Food Legumes*. (Ed.): R.J. Summerfield. Kluwer Academic Publishers, Dordrecht the Netherlands. 445-457.
- 26. Zeidan, M.S. 2007,** Effect of Organic manure and Phosphorus Fertilizers on Growth, Yield and Quality of Lentil Plants in Sandy Soil. *Research Journal of Agriculture and Biological Sciences*, 3(6):748-752.