

Original Research Article

“Effect of NPK and Zinc Inorganic fertilizers on Soil Health, Growth and Yield of Chickpea (*Cicer arietinum* L.)”

Abstract

The objective of study to show the effect on inorganic sources of NPK and Zn fertilizers on soil health. The experiment was laid out in Randomized Block Design with three levels of NPK (00%, 50% and 100%), and Zinc (00%, 50% and 100%) Recommended Dose of fertilizer applied in all treatments. The result showed that the soil properties, growth and yield of chickpea were significantly affected by application of NPK and Zinc. The highest growth and yield were observed in T₉(100%NPK +100% Zinc), followed by T₈ (100% NPK and +50%Zinc), whereas the lowest growth and yield were observed in T₁ (00%NPK +00% Zinc). The treatment T₉ (NPK @ 100 % +Zn @ 100%) was recorded as best treatment for major soil parameters. The treatment T₉ (NPK @ 100% + Zn @ 100 %) also shows the significantly highest vegetative growth as well as yield attributes and net profit ₹45,186.00 ha⁻¹ with cost benefit ratio is 1:2.17.

Key word: Chickpea, NPK, Zinc, Growth and Yield, Physico-chemical Properties of Soil.

Introduction

Soil plays a critical role in chickpea production by providing nutrients, water and physical support for growth. Physico-chemical properties of soil, including texture, pH, organic matter, fertility, moisture, drainage and temperature, play critical role in growth and development of chickpea. Proper management of these properties can lead to a healthy and productive chickpea crop. Soil is the loose surface material that covers most land. It consists of inorganic particles and organic matter. Soil provides the structural support to plants used in agriculture and is also their source of water and nutrients. Soils vary greatly in their chemical and physical properties. Processes such as leaching, weathering and microbial activity combine to make a whole range of different soil types. Each type has strengths and weaknesses for agricultural production (Roy *et al.*, 2006).

The three most significant nutrients, without any one of which plants could not endure, are known as the primary macronutrients: Nitrogen (N), Phosphorus (P), and Potassium (K). In

Comment [MF1]: An introduction of two to three lines should be written on the importance of the topic.

Comment [MF2]: The study area, the location of the field, the experiment in which the study was conducted, and whether it is affiliated with a university, an institution, or a company, must be mentioned.

Comment [MF3]: was applied

Comment [MF4]: Mention the statistical techniques and software, if any, in conducting the study.

Comment [MF5]: chickpeas

Comment [MF6]: the application

Comment [MF7]: The researcher's opinion on the study and its importance in society.

Comment [MF8]: Keywords

Comment [MF9]: roles in the growth

Comment [MF10]: chickpeas

Comment [MF11]: This paragraph serves as an introduction at the beginning of the abstract.

Comment [MF12]: Delete

particular, nitrogen is crucial to chlorophyll, which allows plants to carry out photosynthesis (the process by which they uptake sunlight to produce sugars from carbon dioxide and water). Nitrogen is also a major component in amino acids, the base of proteins. (Mishra *et al.*, 2011).

Nitrogen also aids in the compounds that let for storage and use of energy. The plants that are nutritious and healthy while developing have good nitrogen content in them. Phosphorus also encourages the growth of roots, promotes flowering, and is necessary for DNA. Phosphorus fertilizer extract from the phosphate rock (Dixit, G. P., 2016).

Phosphorus is a vital component of ATP, the "energy unit" of plants. ATP forms during photosynthesis, has phosphorus in its structure, and processes from the beginning of seedling growth through to the formation of grain and maturity. Thus, phosphorus is essential for the general health and vigour of all plants (Mishra *et al.*, 2011).

Potassium is often referred to as the "value element," because of its contribution to many of the features we connect with quality, such as size, figure, color, and even taste, among others. Potassium provides strength to the plants and can resist diseases (Mishra *et al.*, 2011).

Zinc is an important component of various enzymes that are responsible for driving many metabolic reactions in all crops. Growth and development would stop if specific enzymes were not present in plant tissue. Carbohydrate, protein, and chlorophyll formation is significantly reduced in zinc-deficient plants (Singh *et al.*, 2009).

Materials and Methods

A field experiment conducted at the Soil Science Research Farm, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, during the *Rabi* season 2021-22 growing chickpea *Var.* Sadabahar applied 3 levels of NPK and Zinc respectively 0 %, 50 % and 100 % including RDF for chickpea = 20:40:20kg ha⁻¹ experiment is lead to observe the physical and chemical parameters. In physical parameters like that bulk density, particle density, pore space and water holding capacity through method by 100 ml graduated measuring cylinder and process by Muthuvel *et al.*, 1992.

In chemical parameters through method by-

- Soil pH –method given by (Jackson, M. L. 1958) through using digital pH meter.
- Soil EC (dSm⁻¹) - method given by (Wilcox, 1950) through using digital EC meter.
- Organic Carbon (%) - Wet oxidation method given by (Walkley and Black, 1947)
- Available Nitrogen (kg ha⁻¹) - Kjeldhal Method (Subbiah and Asija, 1956)

Comment [MF13]: Delete

Comment [MF14]: Delete

Comment [MF15]: Delete

Comment [MF16]: Before this title, a new title was proposed under the name (description of the study area), and a shape should be placed in the study area in the form of a map with a picture of the plant under study.

Comment [MF17]: Write the same to the abstract.

Comment [MF18]: Delete

Comment [MF19]: led

Comment [MF20]: observing

Comment [MF21]: Delete

Comment [MF22]: the method

Comment [MF23]: a digital

- e) Available Phosphorus (kg ha^{-1}) - Colorimetric method by using Jasper single beam U.V. Spectrophotometer at 660 nm wavelength given by (Olsen *et al.*, 1954)
- f) Available Potassium (kg ha^{-1})- Flame photometric method by using Metzer Flame Photometer given by (Toth and Prince, 1949)
- g) Available Zinc (mg kg^{-1}) –DTPA extractant method by using AAS given by (Lindsay and Norvell, 1978)

Comment [MF24]: The flame

Result and Discussion

Physical Properties of Soil

Bulk density (Mg m^{-3})

The data presented in table and fig. 1 shows the effect of NPK and Zinc on bulk density (Mg m^{-3}) of soil after crop harvested. The response of NPK and Zinc on bulk density of soil was found non-significant. The maximum bulk density of soil 1.33 and 1.36 Mg m^{-3} was found at 0-15 and 15-30 cm in treatment T_1 (NPK@0%+Zn@0%) and minimum bulk density of soil 1.15 and 1.17 Mg m^{-3} was found at 0-15 and 15-30 cm in treatment T_9 (NPK @ 100% + Zn@100%) respectively. Similar result has been recorded by Hussain *et al.*, (2022); Chinth *et al.*, (2021).

Comment [MF25]: the crop is harvested

Comment [MF26]: the minimum

Comment [MF27]: A similar

Particle density (Mg m^{-3})

The data presented in table and fig. 1 shows the effect of NPK and Zinc on particle density (Mg m^{-3}) of soil after crop harvested. The response of NPK and Zinc on particle density of soil was found non-significant. The maximum particle density of soil 2.52 and 2.55 Mg m^{-3} was recorded at 0-15 and 15-30 cm in treatment T_1 (NPK @ 0% + Zn @ 0 %) and minimum particle density of soil 2.38 and 2.45 Mg m^{-3} was recorded at 0-15 and 15-30 cm in treatment T_9 (NPK @ 100% + Zn @ 100 %) respectively. Similar result has been recorded by Hussain *et al.*, (2022); Chinth *et al.*, (2021) and Dang *et al.*, (2020).

Comment [MF28]: harvesting

Comment [MF29]: the minimum

Comment [MF30]: A similar

Comment [MF31]: , and

Percent pore space (%)

The data presented in table and fig 1 shows the effect of NPK and Zinc on percent pore space of soil after crop harvested. The response of NPK and Zinc on percent pore space of soil was found significant. The maximum percent pore space of soil 48.7 and 47.60 % was found at 0-15 and 15-30 cm in treatment T_9 (NPK @ 100 % + Zn @ 100 %) and minimum percent pore space of soil 45.11 and 44.77 % was found at 0-15 and 15-30 cm in treatment T_1 (NPK @ 0 % + Zn @ 0 %) respectively. Similar result has been recorded by Hussain *et al.*, (2022); Chinth *et al.*, (2021) and Dang *et al.*, (2020).

Comment [MF32]: the table and Fig 1

Comment [MF33]: the percent

Comment [MF34]: the crop is harvested

Comment [MF35]: the percent

Comment [MF36]: the minimum

Comment [MF37]: A similar

Water holding capacity (%)

The data presented in table and fig. 1 shows the effect of NPK and Zinc on water holding

Comment [MF38]: the table and Fig. 1

Comment [MF39]: the water

capacity (%) of soil after crop harvested. The response of NPK and Zinc on water holding capacity of soil was found significant. The maximum water holding capacity of soil 44.58 and 42.09 % was found at 0-15 and 15-30 cm in treatment T₉ (NPK @ 100 % + Zn @ 100 %) and minimum water holding capacity of soil 39.23 and 38.05 % was found at 0-15 and 15-30 cm in treatment T₁ (NPK @ 0 % + Zn @ 0 %) respectively. Similar result has been recorded by Hussain *et al.*, (2022); Chintha *et al.*, (2021) and Dangiet *et al.*, (2020).

Comment [MF40]: the crop is harvested

Comment [MF41]: the water holding capacity of the soil

Comment [MF42]: the minimum

Comment [MF43]: A similar

Chemical Properties of Soil

Soil pH (1:2.5) w/v

The data presented in table and fig. 2 shows the effect of NPK and Zinc on pH of soil after crop harvested. The response NPK and Zinc on pH of soil was found non-significant. The maximum pH of soil 7.36 and 7.41 was found at 0-15 and 15-30 cm in treatment T₁ (NPK @ 0 % + Zn @ 0 %) and minimum pH of soil 7.27 and 7.27 was found at 15-30 cm in treatment T₉ (NPK @ 100 % + Zn @ 100 %) respectively. Similar result has been recorded by Parmar and Poonia (2020); Sahu *et al.*, (2020) and Deshlahare *et al.*, (2019).

Comment [MF44]: the pH of soil after the crop is harvested.

Comment [MF45]: the pH of the soil were

Comment [MF46]: the minimum

Comment [MF47]: A similar

Soil EC (dSm⁻¹)

The data presented in table and fig. 2 shows the effect of NPK and Zinc on EC of soil after crop harvested. The response of NPK and Zinc on EC of soil was found non-significant. The maximum EC of soil 0.476 and 0.482 dSm⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₉ (NPK @ 100 % + Zn @ 100 %) and minimum EC of soil 0.445 and 0.448 dSm⁻¹ was recorded at 15-30 cm in treatment T₁ [control (NPK @ 0 % + Zn @ 0 %)] respectively. Similar result has been recorded by Parmar and Poonia (2020); Sahu *et al.*, (2020) and Deshlahare *et al.*, (2019).

Organic carbon (%)

The data presented in table and fig. 2 shows the effect of NPK and Zinc on organic carbon of soil after crop harvested. The response of NPK and Zinc on organic carbon of soil was found non-significant. The maximum organic carbon of soil 0.43 and 0.38 % was found at 0-15 and 15-30 cm in treatment T₉ (NPK @ 100 % + Zn @ 100 %) and minimum organic carbon of soil 0.36 and 0.32 % was found at 0-15 and 15-30 cm in treatment T₁ (NPK @ 0 % + Zn @ 0 %) respectively. Similar result has been recorded by Parmar and Poonia (2020); Sahu *et al.*, (2020) and Deshlahare *et al.*, (2019).

Available nitrogen (kg ha⁻¹)

The data presented in table and fig. 2 shows the effect of NPK and Zinc on available nitrogen

of soil after crop harvested. The response of NPK and Zinc on available nitrogen of soil was found significant. The maximum available nitrogen of soil 322.75 and 316.59 kg ha⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₉(NPK @100% + Zn @ 100 %) and minimum available nitrogen of soil 290.65 and 286.32 kg ha⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₁[control(NPK@0%+Zn @ 0%)]respectively. Similar result has been recorded by Pingoliyaet al., (2014); Yadav et al., (2021) and Banjara andMajgahe (2019).

Comment [MF48]: the crop is harvested

Comment [MF49]: the available

Comment [MF50]: in

Comment [MF51]: the minimum

Comment [MF52]: A similar

Availablephosphorus (kg ha⁻¹)

The data presented in table and fig. 2 shows the effect of NPK and Zinc on available phosphorus of soil after crop harvested. The response of NPK and Zinc on available phosphorus of soil was found significant. The maximum available phosphorus of 35.07 and 33.82 kg ha⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₉(NPK @ 100% + Zn @ 100 %) and minimum available phosphorus of soil 19.40 and 16.43 kg ha⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₁(NPK @0%+Zn @ 0 %)respectively. Similar result has been recorded by Pingoliyaet al., (2014, Yadav et al., (2021); Banjara andMajgahe (2019).

Comment [MF53]: available

Comment [MF54]: the crop is harvested

Comment [MF55]: the minimum

Comment [MF56]: A similar

Availablepotassium(kg ha⁻¹)

The data presented in table and fig. 2 shows the effect of NPK and Zinc on available potassium of soil after crop harvested. The response of NPK and Zinc on available potassium of soil was found significant. The maximum available potassium of soil was recorded 208.42 and 204.67 kg ha⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₉(NPK@100%+Zn@100%) and minimum available potassium of soil 182.23 kg and 178.55 kg ha⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₁[control(NPK @0%+Zn@0 %)]respectively. Similar result has been recorded by Pingoliyaet al., (2014, Yadav et al., (2021); Banjara andMajgahe (2019).

Comment [MF57]: the crop is harvested

Comment [MF58]: the minimum

Comment [MF59]: A similar

AvailableZn(mg kg⁻¹)

The data presented in table and fig. 2 shows the effect of NPK and Zinc on available Zn of soil after crop harvested. The response of NPK and Zinc on available Zn of soil was found significant. The maximum available Zn of soil 0.345 and 0.353 mg kg⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₉(NPK @100%+Zn@100%) and minimum available Zn of soil 0.284 and 0.292 mg kg⁻¹ was recorded at 0-15 and 15-30 cm in treatment T₁[control(NPK@0%+Zn @ 0%)]respectively. Similar result has been recorded by Pingoliyaet al., (2014, Yadav et al., (2021);

Comment [MF60]: the crop is harvested

Comment [MF61]: the minimum

Comment [MF62]: A similar

Banjara andMajgahe (2019).

UNDER PEER REVIEW




Table1: Effect of NPK and Zn on bulk density (Mg m^{-3}), particle density (Mg m^{-3}), pore space (%) and water holding capacity (%) of soil after crop harvest.

Comment [MF63]: , and

Comment [MF64]: light font

Treatment		Bulk density (Mg m^{-3})		Particle density (Mg m^{-3})		Pore space(%)		Water holding capacity(%)	
		0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm
T₁	NPK @ 0 % + Zn @ 0 %	1.33	1.36	2.52	2.55	45.11	44.77	39.23	38.05
T₂	NPK @ 0 % + Zn @ 50 %	1.30	1.32	2.50	2.53	45.97	45.80	39.03	38.64
T₃	NPK @ 0 % + Zn @ 100 %	1.28	1.31	2.48	2.51	46.24	45.88	41.96	39.26
T₄	NPK @ 50 % + Zn @ 0 %	1.27	1.29	2.51	2.54	45.55	45.11	39.76	38.29
T₅	NPK @ 50 % + Zn @ 50 %	1.24	1.26	2.47	2.52	47.05	46.87	41.51	39.83
T₆	NPK @ 50 % + Zn @ 100 %	1.21	1.25	2.44	2.49	47.38	47.07	42.27	41.87
T₇	NPK @ 100 % + Zn @ 0 %	1.20	1.23	2.46	2.51	46.48	45.52	41.89	40.34
T₈	NPK @ 100 % + Zn @ 50 %	1.18	1.21	2.42	2.48	47.60	47.52	43.05	41.25
T₉	NPK @ 100 % + Zn @ 100 %	1.15	1.17	2.38	2.45	48.71	47.60	44.58	42.09
	F-Test	NS	NS	NS	NS	S	S	S	S
	S.Ed. (\pm)	-	-	-	-	0.61	0.59	0.68	0.79
	C.D. at 0.5%	-	-	-	-	1.84	1.76	2.06	2.39

Table 2: Effect of NPK and Zn on pH, electrical conductivity (dSm⁻¹), organic carbon (%), available nitrogen (kg ha⁻¹), available phosphorus (kg ha⁻¹), available potassium (kg ha⁻¹) and available zinc (mg kg⁻¹) of soil after crop harvest.

Comment [MF65]: light font

Treatment		Soil pH (1:2.5) w/v		Electrical Conductivity (dSm ⁻¹)		Organic Carbon (%)		Available Nitrogen (kg ha ⁻¹)		Available Phosphorus (kg ha ⁻¹)		Available Potassium (kg ha ⁻¹)		Available Zinc (mg kg ⁻¹)	
		0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm	0 – 15 cm	15 – 30 cm
T ₁	NPK @ 0 % + Zn @ 0 %	7.36	7.41	0.445	0.448	0.36	0.32	290.65	286.32	19.40	16.43	182.23	178.55	0.284	0.292
T ₂	NPK @ 0 % + Zn @ 50 %	7.36	7.40	0.449	0.451	0.36	0.33	291.45	287.05	20.26	17.87	184.41	181.82	0.328	0.334
T ₃	NPK @ 0 % + Zn @ 100 %	7.35	7.40	0.453	0.456	0.38	0.34	293.37	289.91	22.87	19.09	187.58	185.56	0.288	0.296
T ₄	NPK @ 50 % + Zn @ 0 %	7.32	7.38	0.456	0.459	0.40	0.36	297.72	291.65	23.50	20.60	188.08	186.72	0.321	0.329
T ₅	NPK @ 50 % + Zn @ 50 %	7.31	7.38	0.460	0.465	0.41	0.38	300.68	294.27	25.24	22.28	191.56	189.80	0.331	0.338
T ₆	NPK @ 50 % + Zn @ 100 %	7.31	7.37	0.464	0.469	0.41	0.39	304.80	298.53	28.16	26.54	194.78	192.45	0.337	0.342
T ₇	NPK @ 100 % + Zn @ 0 %	7.28	7.31	0.467	0.473	0.42	0.39	310.06	304.26	30.45	28.27	199.81	195.72	0.340	0.344
T ₈	NPK @ 100 % + Zn @ 50 %	7.28	7.30	0.472	0.477	0.42	0.39	317.35	309.38	33.71	31.20	202.95	198.65	0.342	0.348
T ₉	NPK @ 100 % + Zn @ 100 %	7.27	7.29	0.476	0.482	0.43	0.38	322.75	316.59	35.07	33.82	208.42	204.67	0.345	0.353
	F-Test	NS	NS	NS	NS	NS	NS	S	S	S	S	S	S	S	S
	S.Ed. (±)	-	-	-	-	-	-	2.27	1.84	1.08	0.78	1.57	1.35	0.07	0.08
	C.D. at 0.5%	-	-	-	-	-	-	4.62	3.71	2.19	1.61	2.25	1.78	0.16	0.19




Fig.1: Effect of NPK and Zn on bulk density (Mg m⁻³), particle density (Mg m⁻³), pore space (%) and water holding capacity (%) of soil after crop harvest.

Comment [MF66]: light font




Fig.2: Effect of NPK and Zn on pH, electrical conductivity (dSm⁻¹), organic carbon (%), available nitrogen (kg ha⁻¹), available phosphorus (kg ha⁻¹), available potassium (kg ha⁻¹) and available zinc (mg kg⁻¹) of soil after crop harvest.

Comment [MF67]: light font

Conclusion

The results of the experiment are concluded as the effect of NPK and Zinc on nitrogen (kg ha^{-1}), phosphorus (kg ha^{-1}), potassium (kg ha^{-1}), % pore space and water holding capacity (%) of soil after crop harvest was found significant except on bulk density (Mg m^{-3}), particle density (Mg m^{-3}), pH, EC (dsm^{-1}) and organic carbon (%) of soil after harvest. The treatment T₉ (NPK @ 100 % +Zn @ 100%) was recorded as best treatment for major soil parameters. Therefore, it can be recommended for farmers to obtain best combination Treatment (T₉) for higher farm income and sustainable agriculture.

Comment [MF68]: the best

Comment [MF69]: the best

Comment [MF70]: Write the values please.

Reference

Banjara, G. P. and Majgahe, S. K. (2019) Effect of Biofortification of zinc and iron on yield attributes and yields of chickpea (*Cicer arietinum* L.) through agronomic intervention. *The Pharma Innovation Journal*; **8(10)**: 45-47.

Bouyoucos, G.J., (1927) The hydrometer as a new method for the mechanical analysis of soils. *Soil Science*, **23**: 343-353.

Chintha, H. B., Swaroop, N., Thomas, T., Barthwal, A., Harit, H. and Amjad, A. (2021) Effect of nitrogen, phosphorus, potassium, sulphur and zinc on soil health parameters of chickpea (*Cicer arietinum* L.). *The Pharma Innovation Journal*; **10(11)**: 1955-1961.

Dangi, M., Thomas, T., David, A. A., Kumar, S. and Joshi, A. (2020) Response of different levels of zinc and FYM on soil health and yield attributes of chickpea (*Cicer arietinum* L.). *International Journal of Chemical Studies*; **8(4)**: 3559-3563.

Deshlahare, H. and Banjara, G. P. (2019) Impact of biofortification of zinc and iron on growth parameters and yields of chickpea (*Cicer arietinum* L.) through agronomic intervention in Chhattisgarh plains. *Journal of Pharmacognosy and Phytochemistry*; **8(6)**: 383-386.

Dixit, G. P. (2016) Project Coordinator's Report, All India Coordinated Research Project on Chickpea, *Indian Institute of Pulses Research*, Kanpur.

Fisher, R.A., (1955) Statistical methods and scientific induction. *Journal of the royal statistical society series.17*: 69-78.

- Hussain, M., Banoo, M., Sinha, B. K. and Chand, G. (2022)** Effect of foliar application of zinc and boron on growth, yield and quality attributes in chickpea (*Cicer arietinum* L.). *Journal of Pharmacognosy and Phytochemistry*; **11(3)**: 270-275.
- Jackson, M. L., (1958)** *Soil chemical analysis Prentice Hall of India Ltd.* New Delhi. **219-221.**
- Lindsay, W. L. and Norvell, W. A. (1978)** Development of DTPA soil test for zinc, iron, manganese, and copper. *Soil Science Society of America Journal*, **42**: 421-428.
- Mishra, P. K., Bisht, S. C., Pooja, R., Joshi, G. K., Singh, G., Bisht, J. K. and Bhatt. J. C. (2011)** Bio associative effect of cold tolerant *Pseudomonas* spp. and *Rhizobium leguminosarum*-PR 1 on Iron acquisition, nutrient uptake and growth of lentil (*Lens culinaris* L.). *European Journal of Soil Biology*, **47**: 35-43.
- Munsell, A. H. (1971)** Munsell's description of his colour system, from a lecture to the American Psychological Association. *American Journal of Psychology*, **23(2)**: 236-244.
- Muthuvel, P., Udayasoorian, C., Natesan, R. and Ramaswamy, P. P. (1992)** Introduction to Soil Analysis, *Tamil Nadu Agricultural University Coimbatore*-641002.
- Olsen, S. R., Cole, C. V., Watanabe, F. S. and Dean, L. A., (1954)** Estimation of available phosphorus in soils by extraction with sodium bicarbonate (NaHCO_3), *U.S.D.A. Circular*. **939**: 1-19.
- Parmar, P. M. and Poonia, T. C. (2020)** Effect of zinc biofortification on growth, yield and economics of chickpea (*Cicer arietinum* L.). *International Journal of Chemical Studies*; **8(2)**: 1782-1786.
- Pingoliya, K. K., Dotaniya, M. L. and Lata, M. (2014)** Effect of iron on yield, quality and nutrient uptake of chickpea (*Cicer arietinum* L.). *African Journal of Agricultural Research*, **9(37)**: 2841-2845.
- Roy, R. N., Finck, A., Blair, G. J. and Tandon, H. L. S. (2006)** Plant nutrition for food security. A guide for integrated nutrient management. FAO fertilizer and Plant Nutrition Bulletin 16. Rome, Italy, food and Agriculture Organization of the United Nations.

- Sahu, A., Swaroop, N., David, A. A. and Thomas, T. (2020)** Effect of Different Levels of NPK and Zinc on Soil Health Growth and Yield of Chickpea (*Cicer arietinum* L.) var. PUSA 362. *Int. J. Curr. Microbiol. App. Sci.*, **9(10)**: 591-597.
- Singh, A., Singh, V. K., Chandra, R. and Srivastava, P. C. (2009)** Effect of the integrated nutrient management on chickpea based intercropping system and soil properties in Mollisols of the tarai region. *Journal of the Indian Society of Soil Science*, **60**: 38-44.
- Subbiah, B. V. and Asija, E. C., (1956)** A rapid procedure for estimation of available nitrogen in soil. *Current Science*; **25(8)**: 259-260.
- Toth, S. J. and Prince, A. L. (1949)** Estimation of cation exchange capacity and exchangeable Ca, K and Na content of soil by flame photometer technique. *Soil Sci.*, **67**: 439-445.
- Walkley, A. and Black, I. A., (1947)** Estimation of soil organic carbon by the chromic acid titration method. *Soil Science*. **47**: 29-38.
- Wilcox, L. V., (1950)** Electrical conductivity Am. water work, Association **42**: 775-776.
- Yadav, R., Thomas, T., Swaroop, N. and Kumar, T. (2021)** Response of different levels of phosphorus and zinc on soil health after harvesting chickpea (*Cicer arietinum* L.). *The Pharma Innovation Journal*; **11(5)**: 402-405.