

# Effect of different level of NPK Fertilizers, FYM and Vermicompost on Yield attributes of Okra (*Abelmoschus Esculentus L.*) var. Devika

## ABSTRACT

The field experiment was carried out at soil science research farm of Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during *kharif* season 2022. The soil experimental area was sandy loam in texture. The design was laid out in randomized block design with three levels of NPK (0%,50%, & 100%), three levels of FYM and three levels of Vermicompost (0%, 50%, & 100%) respectively. The results obtained with treatment T<sub>9</sub>-[120:60:50 Kg NPK + 25 t FYM + 6 t Vermicompost] showed highest plant height, number of branches plant<sup>-1</sup>, number of leaves plants<sup>-1</sup>, number of fruits plant<sup>-1</sup>, total fruit yield. No significant was observed in yield and growth of okra under control. Application of Organic manures well as its mixture with full NPK considerably increase in growth and total yield attributes of okra. In contrast to all other treatments, use of 100% NPK, 100% FYM and 100% Vermicompost shows the most significant impact with effective on okra growth.

**Keywords:** NPK: Nitrogen, Phosphorus, Potassium, FYM: Farmyard manure, Vermicompost, Yield attributes.

## INTRODUCTION

“Be it deep or shallow, red or black, sand or clay, the soil is the link between the rock core of the earth and living things on its surface. It is the foot fold for the plants we grow, therein lies the main reason for our interest in soils Simonson” [1]. “India ranks first in the World with production of 6416.0 thousand metric tonnes (74% of the total world production) with an area of 523 thousand ha of okra. It is also cultivated in Nigeria, Sudan, Pakistan, Ghana, Egypt, Benin, Saudi Arabia, Mexico and Cameroon. Gujarat is the leading okra producing state which has production of around 1019.42 thousand tons from an area of 85.15 thousand hectare, with a productivity of 11.97 t ha<sup>-1</sup>. It is followed by Uttar Pradesh (335.86 thousand tonnes from 24.80 thousand hectare with 13.54 tonnes ha<sup>-1</sup> productivity) National

Horticulture Board” [2]. “Okra is an important vegetable crop which supplies higher nutrition, the green pod per 100g edible portion of okra contain moisture 89.6g, carbohydrates 6.6g, protein 1.9g, fat 0.2g, fibre 1.2g, mineral 0.7g, calcium 66mg, magnesium 43mg, phosphorus 56mg, potassium 103 mg, thiamine 0.07mg, riboflavin 0.1mg, nicotinic acid 0.6mg, vitamin-C 13mg, Oxalic acid 8mg” Choudhary *et al.*, [3]. “Okra fruit is principally consumed freshly cooked and is a major source of vitamins A, B, C, mineral, iron and iodine and important vegetable source of viscous fibre but it reportedly lows in sodium saturated and cholesterol” Singh *et al.*, [4]. “Okra is a rich source of iodine and other vital minerals and vitamins. Mucilage present in okra is polysaccharides *i.e.*, galacturonic and glucuronic acids” Singh and Ram [5]. “Nitrogen is an essential macro nutrients and determinant in growth and development of crop plants. Nitrogen plays a vital role in chlorophyll, protein, nucleic acid, hormones and vitamins synthesis. Nitrogen also helps in cell division, cell elongation and linear increase in green pod yields of okra” Das *et al.*, [6]. “Organic manure increases cation exchange capacity, water holding capacity and phosphate availability of the soil beside improving the fertilizer use efficiency and microbial population of soil, it reduces nitrogen loses due to slow release of nutrients” Tadesse *et al.*, [7]. “Phosphorus is a key element in the formation of high energy compounds, such as AMP, ADP and ATP, which play an essential role in photosynthesis and respiration. It is a vital component of nucleic acids and phospholipids. Plants take up phosphorus in the inorganic form, mainly as the orthophosphate ion. Phosphorus supports early phase of crop development, synchronizes the germination process and leading to enhance the final yield, especially in P deficient soil” Meena *et al.*, [8].

## METHODOLOGY

A field experiment was conducted in the year 2022 during the Kharif season at research farm of Department of Soil Science and Agricultural Chemistry, Sam Higginbottom University of Agriculture, Technology and Sciences Prayagraj (Allahabad) 211007 U.P., India. The soil of experimental area falls in order Inceptisol and the experimental field is alluvial in nature. The design applied for statistical analysis was carried out with 3<sup>2</sup> factorial randomized block design having three levels of NPK @ 0, 50 and 100 % ha<sup>-1</sup>, three levels of FYM @ 0, 50 and 100 % ha<sup>-1</sup> and three levels of Vermicompost @ 0, 50 and 100 % ha<sup>-1</sup> respectively. The details of the treatment combinations are given below Table No.1. and observation were recorded plant height, number of leaves plant<sup>-1</sup>, number of branches plant<sup>-1</sup>, number of fruits

and total yield of fruits.

The source of inorganic nutrients sources as Urea, SSP, MOP and organic nutrients sources as FYM and Vermicompost. Basal dose of fertilizer was applied in respective plots according to treatment allocation in furrows opened by about 5cm depth before sowing seeds in soil at the same time of sowing of seeds was shown on well-prepared beds in shallow furrows, at the depth of 5 cm , row to row distance was maintained at 30 cm and plant to plant distance was 45 cm, during the course of experiment, observations were recorded as mean values of the data.

## RESULTS AND DISCUSSION

### Plant Height

The response of plant height of okra recorded at 30 DAS, 60 DAS and 90 DAS detailed results were in shown in Table 2 as influenced by different levels of NPK FYM and Vermicompost. The plant height of okra was found to have increased significantly with the age of plants and increase in the levels of inorganic fertilizers and organic manures. The maximum plant height was recorded as 23.6 cm, 90.2 cm and 117.2 cm in T<sub>9</sub> at 30 DAS, 60 DAS and 90 DAS respectively and the minimum plant height was recorded as 20.2 cm, 83.0 cm and 107.0 cm in T<sub>1</sub>-[ Absolute control] at 30 DAS, 60 DAS and 90 DAS respectively. “Inorganic fertilizer and organic manure play an important role in increasing production, improving quality of vegetables and sustaining soil fertility. Organic manure contains all nutrients which are required for healthy growth of crops and helps to improve physical, chemical and biological properties of soil” Ola *et al.*, [9].

### Number of Leaves

The number of leaves of okra at different days after sowing (DAS) at 30, 60 and 90 detailed results were in shown in Table 2 as influenced by different levels of NPK FYM and Vermicompost were found significantly in treatment T<sub>9</sub> was 18.6, 44.2 and 45.2 respectively. While the minimum values of the result were found in treatment T<sub>1</sub>-[ Absolute control] 12.2, 36.8 and 39.4 respectively. The effect of different doses of NPK fertilizers on growth of okra regarding number of leaves and height of had significant superiority over other means Khetran *et al.*, [10].

### Number of Branches

The response of number of branches plant<sup>-1</sup> recorded at 30 DAS, 60 DAS and 90 DAS detailed results were in shown in Table 2 as influenced by different levels of NPK FYM and Vermicompost. The Number of branches plant<sup>-1</sup> of okra was found to have increased significantly with the age of plants and increase in the levels of inorganic fertilizers and organic manures. The maximum number of branches plant<sup>-1</sup> was recorded as 1.8,2.4 and 2.6 in T<sub>9</sub> at 30 DAS, 60 DAS and 90 DAS respectively and the least number of branches was recorded as 0.6, 0.8 and 1.0 in T<sub>1</sub>-[ Absolute control] at 30 DAS, 60 DAS and 90 DAS respectively.

“Concluded that the use of organic manure in combination with inorganic fertilizers in the production of vegetables like okra should be encouraged as it is beneficial for the physical growth of okra plants” Gayathri *et al.*, [11].

### **Number of Fruits**

The response of number of fruits plant<sup>-1</sup> recorded and detailed results were in shown in Table 2 as influenced by different levels of NPK FYM and Vermicompost. The number of fruits plant<sup>-1</sup> of okra was found to have increased significantly with the application of inorganic fertilizers and organic manures. The maximum number of fruits plant<sup>-1</sup> was recorded as 23.2 in T<sub>9</sub> and minimum number of fruits plant<sup>-1</sup> was recorded as 14.0 in T<sub>1</sub>-[ Absolute control].

“Potassium is one of the three major nutrient elements (N, P and K) required by plants. Potassium imparts vigour and disease resistance to the plant and plays an important role in crop productivity” Ginindza *et al.*, [12].

### **Fruit Yield**

Highest yield of fruits due to influence of NPK FYM and Vermicompost was recorded as 178.50 in T<sub>9</sub> and minimum yield was recorded as 110.81 in T<sub>1</sub>-[ Absolute control].

“The requirements of fertilizers in okra are important for the early growth and total yield of fruit. Integrated use of Organic and Inorganic fertilizers can improve crop productivity. The soil enriched with vermicompost provides additional substances that are not found in chemical fertilizers” Mal *et al.*, [13].

**Table.1: Treatment combination**

<b>Treatment</b>	<b>Treatment combination</b>
T <sub>1</sub>	Absolute Control
T <sub>2</sub>	[NPK @ 0% + FYM @ 0% + Vc @ 50%]
T <sub>3</sub>	[NPK @ 0% + FYM @ 0% + Vc @ 100%]
T <sub>4</sub>	[NPK @ 50% + FYM @ 50% + Vc @ 0%]
T <sub>5</sub>	[NPK @ 50% + FYM @ 50% + Vc @ 50%]
T <sub>6</sub>	[NPK @ 50% + FYM @ 50% + Vc @ 100%]
T <sub>7</sub>	[NPK @ 100% + FYM @ 100% + Vc @ 0%]
T <sub>8</sub>	[NPK @ 100% + FYM @ 100% + Vc @ 50%]
T <sub>9</sub>	[NPK @ 100% + FYM @ 100% + Vc @ 100%]

*Note: NPK 100% (100:60:50), FYM 100% (25 t) and Vc 100% (6 t)*

**Table 2: Effect of different level of NPK FYM and Vermicompost on Yield attributes of Okra**

Treatment	Plant height (cm)			Number of leaves			Number of branches			Number of fruits	Fruit Yield (q ha <sup>-1</sup> )
	30DAS	60DAS	90DAS	30DAS	60DAS	90DAS	30DAS	60DAS	90DAS		
T <sub>1</sub>	20.2	83.0	107.0	12.2	36.8	39.4	0.6	0.8	1.0	14.0	110.81
T <sub>2</sub>	21.2	84.0	107.6	13.0	37.4	40.0	0.8	1.0	1.2	14.2	117.84
T <sub>3</sub>	20.8	85.2	108.0	13.4	38.4	40.0	0.8	1.2	1.4	14.6	120.56
T <sub>4</sub>	21.6	86.4	109.8	14.0	39.2	40.8	1.0	1.4	1.6	15.4	122.38
T <sub>5</sub>	22.0	87.2	111.0	14.8	39.2	41.8	1.2	1.6	1.8	16.2	130.60
T <sub>6</sub>	22.0	87.4	111.4	15.4	41.0	42.4	1.4	1.8	2.0	18.0	136.70
T <sub>7</sub>	22.4	87.8	112.6	16.0	41.8	43.0	1.4	1.8	2.2	19.6	150.35
T <sub>8</sub>	23.0	89.0	115.6	17.2	43.2	44.2	1.6	2.0	2.4	20.8	174.55
T <sub>9</sub>	23.6	90.2	117.2	18.6	44.2	45.2	1.8	2.4	2.6	23.2	178.50
F-test	S	S	S	S	S	S	S	S	S	S	S
S.Em. (±)	0.35661	1.15136	1.55396	0.30157	0.54027	0.75387	0.01611	0.02102	0.01669	0.23018	1.61692
C.D. (P=0.05)	1.13091	3.46592	4.67787	0.90781	1.62638	2.26938	0.0485	0.06327	0.05024	0.69291	4.8674

## CONCLUSION

It revealed from the trial that application of NPK FYM and Vermicompost in treatment T<sub>9</sub> was found best in increasing growth and yield of okra. Since the results is based on one season experiment, further trail is needed to substantiate the result.

## REFERENCES

1. Simonson, R. (1989). Historical Highlights of soil survey and soil classification with emphasis on the United States.
2. Anonymous (2021) National Horticulture Board.
3. Choudhary, K., More, S. J. and Bhanderi, D. R. (2015). Impact of bio-fertilizers and chemical fertilizers on growth and yield of okra (*Abelmoschus esculentus* L. Moench). *International Quarterly Journal of Environmental Science*. 9(1&2): 67-70.
4. Singh, P., Chauhan, V., Tiwari, B.K., Simon, S. and Abidi, A.B. (2015). An overview on Okra (*Abelmoschus esculentus* L.) and its importance as a nutritive vegetable in the World. *International Journal of Pharmacy and Biological Sciences*. 4(2):227-233.
5. Singh, S. and Ram, R.B. (2018) Effect of organic, inorganic and bio-fertilizers on yielding and fruiting traits of okra [*Abelmoschus esculentus* (L.) Moench]. *Journal of Pharmacognosy and Phytochemistry*. 7(5): 90-93 .
6. Das, A.K., Prasad, B. and Singh, R. (2014). Response of chemical fertilizer and vermicompost on okra (*Abelmoschus esculantus* L.). *The Asian Journal of Horticulture*. 9: 372-376.
7. Tadesse, T., Dechassa, N., Bayu, W. and Gebeyehu, S. (2013). Effects of farmyard manure and Inorganic fertilizer application on soil Physico-chemical properties and nutrient balance in rain-fed lowland rice ecosystem. *American Journal of Plant Sciences*. 4:309-316.
8. Meena, N.K., Meena, R.K., Dhaka, R.S. and Meena, O.P. (2017). Response of Nitrogen, Phosphorus and Potassium Levels on Growth and Yield of Okra [*Abelmoschus esculentus* (L.) Moench]. *Int. J. Pure App. Bio-sci*. 5 (4): 1171-1177.
9. Ola, R., David, A. A., Singh, P. and Baloda, S.S. (2018). Response of different levels of N P K and FYM on growth and yield of okra (*Abelmoschus esculentus*) Var. Arka Anamika. *International Journal of Chemical Studies*. 6(5):1098-1101.
10. Khetran, R., Kasi, M. A., Agha, S. H. and Fahmid, S. (2016). Effect of different doses of

NPK fertilizers on growth of okra. *International Journal of Advanced Research in Biological Sciences*. ISSN:2348- 8069.

11. Gayathri, K. and Reddy, P.S. (2013). Effect of integrated nutrient management growth and yield of okra (*Abelmoschus esculentus* (L). Moench) cv. Arka Anamika. *Vegetable Science*. 40(2): 246-248.
12. Ginindza, T. K., Masarirambi, T. M. and Wahome, P. K. (2015) Effect of different concentrations of NPK fertilizers on growth and development of okra. *Agriculture and Biology Journal of North America*. 6(3):2151-7525.
13. Mal., M. Mahapatra., P. and Mohanty, S. (2014). Effect of Diazotrophs and Chemical Fertilizers on Production and Economics of Okra (*Abelmoschus esculentus*, L.) Cultivars. *American Journal of Plant Sciences*. 5:168-174.

UNDER PEER REVIEW