

Case report

INTENSIVE CARE MANAGEMENT OF GITELMAN SYNDROME: A RARE CASE

ABSTRACT

Gitelman syndrome (GS) is an autosomal-recessive disorder distinguished by hypokalemia, hypomagnesaemia, and hypocalciuria. Elderly people and women of childbearing age are highly affected by GC. There isn't much evidence known about its effects on maternal and fetal outcomes. GS is caused by mutations in the thiazide-sensitive Na-Cl cotransporter gene. Due to its rarity and lack of knowledge, It is susceptible to misdiagnosis or being overlooked. In our case, the patient was suffering from recurrent hypokalemia, hypomagnesemia, hypochloraemia, and hypocalciuria with hypotension. After taking proper medication patients recovered slowly and during patient counseling provided diet chart by nutritionists to avoid recurrent electrolyte imbalance.

Keywords: [Hypokalemia, hypochloraemia, hypocalciuria, hypotension]

1. INTRODUCTION

Gitelman syndrome (GS) is a rare autosomal-recessive disorder, first described in 1966 by Gitelman et al. GS also known as Gitelman's version of Bartter's Syndrome (BS), is a hereditary tubulopathy which considers salt-losing that runs in the autosomal-recessive family [1]. Estimated data from several genomic databases, GS affects around 1 in 40,000 individuals, despite the fact that Asians may be more prone to it [2]. GS is characterized by hypokalemia, hypomagnesemia, metabolic alkalosis, and hypocalciuria. [3]. Low serum potassium level is the characteristic of hypokalemia. Hypokalemia mainly represents electrolyte imbalance. When potassium levels are below 2.5 mEq/L than severe, life-threatening neurological and cardiac problems may appear. It can be due to various causes. Hypokalemia cannot be corrected just by lowering potassium levels, which may lead to recurrence and severe complications [4]. Although BS (Barter's syndrome) and GS (Gitelman's syndrome) share several phenotypic characteristics, GS can be distinguished by having hypomagnesemia and hypocalciuria. Hypomagnesemia may be present, and hypocalciuria is exceedingly variable. It occurred due to the presence of an inactivating mutation in the gene SLC12A3, on chromosome 16, which codes for the thiazide-sensitive sodium-chloride cotransporter and the magnesium transporter on the apical membrane of the Distal Convoluted Tubule (DCT). GS is a benign tubulopathy that may have no symptoms. But the symptoms that have been described in the literature can range from mild weakness, weariness, salt seeking, thirst, and nocturia to muscle weakness, paralysis, paresthesias, or the symptoms of neuromuscular excitability, such as tetany, and infrequent seizures [5].

2. CASE PRESENTATION

This is a case of 55-year-old male with recurrent hypokalemia, admitted to the general medicine department. The patient presented with complaints of muscle weakness, dizziness and low blood pressure of 108/72, systolic blood pressure was always below 110 mmHg with positive anamnesis, and no further symptoms of the urinary

tract, he only mentioned polyuria and nocturia. His medical history was not significant with negative family history. He denied using any drugs other than vitamin supplements intake. When the patient was being treated in a hospital for community-acquired pneumonia at the age of 53, this illness was first discovered. Although the last six months he had been taking oral magnesium and potassium supplements inconsistently. During the physical examination, he showed normal hydration, skin and mucosa coloration, a blood pressure of 108/68 mm Hg, and a regular pulse frequency of 80 beats/min. The cardiopulmonary assessment was normal with no evidence of peripheral edema. The remaining physical checkup was normal. A metabolic alkalosis was presented during the gas analysis (pH 7.435; HCO₃ 33.1 mmol/l; pCO₂ 38 mm Hg). In biochemical analysis, the report confirm hypokalemia (2.42 mmol/l), hypomagnesemia (0.35 mmol/l), and hypochloreaemia (89 mmol/l). urea (58mg/dl), Serum creatinine (2.8 gm/dl), serum sodium level 123.4 mmol/l and remainder ionogram was normal. Further investigation found raised plasma-active renin (2719 fmol/l/sec), with normal aldosteronaemia (16.7 ng/ml in orthostatic; NR 4–31), hypocalciuria (0.73 mmol/l; NR 2.5–7.5) and a greater amount of urine excretion of sodium (126.5 mmol/l; NR 20-110) and chloride (166 mmol/l; 55-125). The estimated glomerular filtrate rate (GFR) was 120 ml/min/1.73 m² and the potassium transtubular gradient was 11.6. And Other tests like Renal ultrasonography, and renal and adrenal CT indicate normal kidney function. A correlation between hypokalemia, hypomagnesemia, metabolic alkalosis, hypocalciuria, and low blood pressure led to the diagnosis of GS.

Table.1 During the initial presentation, biochemical parameters:

Test	Value	Normal range
Serum potassium, mmol/l	2.42	3.6–5.4
Serum magnesium, mmol/l	0.35	0.7–1.0
Serum Sodium level, mmol/l	30	135–145
Blood urea, mg/dl	58	10–50
Blood calcium level mmol/l	0.73	2.5-7.5
Serum creatinine gm/dl	2.4	0.6–1.6
Plasma renin activity, fmol/l/s	2,719	100–1,500
Serum Chloride mmol/l	89	96-108
Urine Sodium mmol/l	126.5	20-110
Urine Chloride mmol/l	166	55-125

TREATMENT

The patient took medication according to the prescribed manner, magnesium aspartate 1230 mg once a day (OD), Tolvaptan tablet 15 mg once a day (OD), potassium chloride capsule 600 mg twice daily (BD), and started taking magnesium aspartate/potassium aspartate 250/250 mg orally four times per day (qid) from admission time. During counseling, The patient was advised to keep up a high-sodium and high-potassium diet by nutritionists. 20 days later patient's serum potassium levels of 2.9 mmol/l and magnesium levels of 0.52 mmol/l. The patient recovered slowly with the proper care of the healthcare team.

3. DISCUSSION

A frequent clinical issue with potentially fatal symptoms is chronic hypokalemia. long-term symptomatic hypotension, hypomagnesaemia, and metabolic alkalosis. Our patient is affected by these conditions. The two main diagnoses in this situation—vomiting and diuretic abuse—were ruled out by measuring high urinary chloride excretion and having no history of using diuretics, respectively. The genetic syndromes Gitelman and Bartter were the only remaining differential diagnoses. Due to its more severe phenotype, earlier start, frequent rise in urinary calcium excretion, and normal or modestly hypomagnesaemia, Bartter syndrome is considered implausible. On the basis of laboratory test reports final diagnosis was GS. It is an autosomal recessive renal tubulopathy that causes salt loss. Sodium-chloride cotransporter sensitive to thiazides, which is found in the epithelial cells of the renal distal convoluted tubule (DCT), and it is encoded by the gene in the vast majority of instances [6]. Its distinguishing characteristics, secondary hyperaldosteronism, hypomagnesaemia, and hypocalciuria, cause hypokalemia and metabolic alkalosis. Hypomagnesemia can inhibit parathyroid hormone secretion [7]. The clinical signs are the same as when thiazide diuretics are used for a long time [8]. Clinical signs and biochemical abnormalities, such as hypomagnesaemia, hypokalemia, metabolic alkalosis, and hypocalciuria, are used to make the diagnosis of GS. This type of patient responds to thiazide with a blunted natriuretic response, but responds quickly to furosemide, indicating that the abnormality is at the level of the distal tubule. Analysis of the GS gene's DNA for mutations may help to confirm the diagnosis [9] According to global consensus suggestion, blood potassium and magnesium levels should be maintained at or above 3.0 mmol/L and > 0.6 mmol/L, respectively in GS patients [10-11]. Most asymptomatic patients do not receive treatment, and they only sometimes have ambulatory monitoring. Renal insufficiency can progress but it is incredibly uncommon [12].

4. CONCLUSION

This case report features a patient who diagnosed with GS symptoms of hypomagnesemia, persistent hypokalemia, and metabolic alkalosis. Inconsistently, patient was taking oral magnesium and potassium supplements over the previous six months. Frequent rise in urinary sodium and chloride excretion, and normal or mild hypomagnesaemia is considered implausible. When thiazide diuretics are used for a prolonged period of time, then the clinical symptoms are the same. During patient counseling provided diet chart by nutritionists to avoid recurrent electrolyte imbalance. After 20 days, his serum potassium and magnesium levels were 2.9 mmol/l and 0.52 mmol/l, respectively. We strengthened ion supplementation, and the patient gradually achieved a full recovery.

CONSENT

Written informed consent was obtained from the patient for the publication of this case report.

ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

REFERENCES

1. Riveira-Munoz E, Chang Q, Godefroid N, Hoenderop JG, Bindels RJ, Dahan K, Devuyst O. Transcriptional and functional analyses of SLC12A3 mutations: new clues for the pathogenesis of Gitelman syndrome. *Journal of the American Society of Nephrology*. 2007;18(4):1271-83.
2. Kondo A, Nagano C, Ishiko S, Omori T, Aoto Y, Rossanti R, Sakakibara N, Horinouchi T, Yamamura T, Nagai S, Okada E. Examination of the predicted prevalence of Gitelman syndrome by ethnicity based on genome databases. *Scientific Reports*. 2021 ;11(1):16099.
3. Cruz DN, Shaer AJ, Bia MJ, Lifton RP, Simon DB. Gitelman's syndrome revisited: an evaluation of symptoms and health-related quality of life. *Kidney international*. 2001;59(2):710-7.
4. Bettinelli A, Bianchetti MG, Girardin E, Caringella A, Cecconi M, Appiani AC, Pavanello L, Gastaldi R, Isimbaldi C, Lama G, Marchesoni C. Use of calcium excretion values to distinguish two forms of primary renal tubular hypokalemic alkalosis: Bartter and Gitelman syndromes. *The Journal of pediatrics*. 1992;120(1):38-43.
5. Blanchard A, Bockenhauer D, Bolognani D, Calo LA, Cosyns E, Devuyst O, Ellison DH, Frankl FE, Knoers NV, Konrad M, Lin SH. Gitelman syndrome: consensus and guidance from a kidney disease: improving global outcomes (KDIGO) controversies conference. *Kidney international*. 2017;91(1):24-33.
6. Simon DB, Nelson-Williams C, Johnson Bia M, Ellison D, Karet FE, Morey Molina A, Vaara I, Iwata F, Cushner HM, Koolen M, Gainza FJ. Gitelman's variant of Barter's syndrome, inherited hypokalaemic alkalosis, is caused by mutations in the thiazide-sensitive Na-Cl cotransporter. *Nature genetics*. 1996 ;12(1):24-30.
7. Pantanetti P, Arnaldi G, Balercia G, Mantero F, Giacchetti G. Severe hypomagnesaemia-induced hypocalcaemia in a patient with Gitelman's syndrome. *Clinical endocrinology*. 2002 Mar;56(3):413-8.
8. Jeck N, Schlingmann KP, Reinalter SC, Kömhoff M, Peters M, Waldegger S, Seyberth HW. Salt handling in the distal nephron: lessons learned from inherited human disorders. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2005;288(4):R782-95.
9. Knoers NV, Starremans PG, Monnens LA. Hypokalemic tubular disorders. *Oxford textbook in clinical nephrology*. 2005;3:995-1004.
10. Gitelman Syndrome Collaborative Study Group. Expert consensus for the diagnosis and treatment of patients with Gitelman syndrome. *Zhonghua nei ke za zhi*.;56(9):712-6.
11. Knoers NV. Gitelman syndrome. *Advances in chronic kidney disease*. 2006;13(2):148-54.
12. Bonfante L, Davis PA, Spinello M, Antonello A, D'Angelo A, Semplicini A, Calò L. Chronic renal failure, end-stage renal disease, and peritoneal dialysis in Gitelman's syndrome. *American journal of kidney diseases*. 2001;38(1):165-8.
- 13.