

## *Original Research Article*

### **Applicability of multispectral images to detect soil organic carbon content in land suitability assessment: A case of a sugarcane plantation**

**Abstract** - Soil organic carbon is important for sugarcane production as it plays a significant role in the development of soil aggregates and soil improvement in any agricultural soil, hence increasing soil health. Increasing soil organic matter encourages soil aggregation and slows the rate of organic matter breakdown. Soil aggregates act as nuclei for soil stabilization with time. It improves soil attributes by increasing organic carbon content in the soil and enhances the physical properties of the soil that favor water infiltration and retention. However, traditional methods of SOC determination like laboratory analyses are expensive and time-consuming. The objective of the study was to assess the applicability of multispectral images to determine SOC content in the surface soil. The research adopted two means to determine SOC of sugarcane. Under the first method, the lands were selected according to the sugarcane productivity in the study area, namely: low (35-54 t ha<sup>-1</sup>), medium (55-79 t ha<sup>-1</sup>), and high (80-100 t ha<sup>-1</sup>) productivity. Soil samples were collected up to 15cm depth. Walkley and Black Method was used for SOC determination in the laboratory. Simultaneously, multispectral images of each land were obtained using a drone platform. Multispectral images were then used to calculate Normalized Difference Vegetation Index (NDVI), Bare Soil Index (BSI), and Modified Secondary Soil Adjusted Vegetation Index (MSAVI2). These indices were used to predict SOC. The outcome was compared with SOC obtained from the laboratory analysis results. The results showed that the soil organic carbon (SOC) varied between 2.72 and 3.63% from the mean

for the the highest and lowest productivity lands respectively. The results of regression analysis observed a moderate correlation between SOC and the BSI values (0.4828). Weak correlations were observed in MSAVI2 (0.0269) and NDVI (0.0858). Future research should focus on improving these indices for SOC determination with increasing sample quantity.

**Keywords** - Soil organic carbon, land suitability assessment, multispectral images, sugarcane

## **1. Introduction**

Human beings depend on land for their survival (Schoonover and Crim, 2015). However, land is becoming less productive due to overuse (Montort, et al, 2021, Eswaran, 2019, Kath et al 2019). Human activities are the main drivers of land degradation, which leads to loss of biodiversity and vegetation cover, imbalance and depletion of soil nutrients and organic matter (SOM), and reduced soil water holding capacity (Mng'ong'o et al., 2021). Soil degradation is a specific form of land degradation that occurs when lands are poorly managed.

Organic carbon (OC) is a key component of SOM, which is essential for soil health (Ngatia et al., 2021). To improve crop productivity through soil management, it is important to measure SOC content in the soil regularly. However, laboratory methods for SOC detection are expensive and time-consuming (Koparan, 2019). Alternative methods that use Geographic Information Systems (GIS) and Remote Sensing (RS) may replace conventional SOC detection techniques in the future (Chaminé et al., 2021). Koparan (2019) states that remote sensing images can be obtained from satellites and Unmanned Aerial Vehicle (UAV) systems. Gomez et al. (2008) have used RS data and hyper-spectral proximate data from visible and near-infrared reflectance to compare the predictions of SOC and other soil properties. Vašát et al. (2014) have used visible and near-infrared diffuse reflectance spectroscopy (VNIR-DRS) to predict soil nutrients based on the peak parameters from continuum-removed spectra. Castaldi et al. (2018) have proposed a

'bottom-up' method to predict SOC from RS data without any laboratory tests. However, the accuracy of these methods needs more research. Therefore, it would be beneficial if remote sensing applications could be used as a new technology to detect SOC content in a fast and cost-defective means.

The purpose of this study was to investigate whether multispectral images can be used to detect SOC instead of laboratory testing for land suitability assessment in a sugarcane field. The study also aimed to determine the correlation between SOC measurements and different vegetation indices (Normalized Difference Vegetation Index, Bare Soil Index, and Modified Soil Adjusted Vegetation Index 2) in this context.

## 2. Material and methods

### 2.1 Study Area

Sugarcane lands of Lanka Sugar Company Ltd., Pelwatta, which is located in Monaragala District of the Uva Province in Sri Lanka ( $6.7230^{\circ}$  N and  $81.2044^{\circ}$  E) (Figure 1) and 175m above sea level, under tropical climate, were used as the study site. Nine (9) lands, each with 1 ha extent, were selected for obtaining soil samples and multi-spectral images. The field is covered with sugarcane cultivation and replanting is carried out, as needed.




Figure 1: Study area of Lanka Sugar Company Ltd., Pelwatta Unit.

These lands were chosen for the experiment based on the productivity of the sugarcane crop. Accordingly, the lands with a yield of 35-54 t ha<sup>-1</sup> were considered as lands belonging to the low productivity zone; the yield between 55-79 t ha<sup>-1</sup> was under the moderate productivity zone; and the yield between 80-100 t ha<sup>-1</sup> was categorized into the lands belonging to the high productivity zone by considering yield quantity of sugarcane lands. 3 replicates from each productivity category were used. Figure 2 depicts the research methodology related to the present study.




Figure 2: Flow chart of research methodology.

## 2.2 Soil Sampling and Analysis

Soil samples were collected from each 1 ha land, under three yield productivity classes, 2 to 4 weeks before ploughing using an augur up to a depth of 0-15 cm. Under each category of productivity, three plots were selected and from each plot, 5 representative soil samples were collected. Consequently, 45 soil samples were collected from 9 plots and SOC contents were measured. The obtained soil samples were subjected to laboratory testing to calculate the SOC concentration using Walkley-Black Wet Oxidation Method (World Bank, 2021) according to Eq. (1).

$$\%OC = [M_{Fe} \times (V_{blank} - V_{sample}) \times 0.003 \times 100 \times f \times mcf] / W \quad (1)$$

where;

$V_{blank}$  = volume of titrant in blank, ml

$V_{sample}$  = volume of titrant in sample, ml

$M_{Fe}$  = concentration of standardized  $FeSO_4$  or  $(NH_4)_2 Fe(SO_4)_2 \cdot 6H_2O$  solution,

molarity

f = correction factor, 1.3

W = weight of soil, g

mcf = moisture correction factor, 1

Besides, undisturbed ring samples were taken for bulk density determination. A hand held GPS device was used to get the location information of the each sampling point.

### 2.3 Spatial Data Collection and Processing

Acquisition of multispectral images was carried out using drone technology parallel with the collection of soil samples. The drone was flown horizontally at an altitude of 25 m and images were taken with a spatial resolution of 3 cm pixels and the ground coordinates were obtained correctly by the RTK GPS receiver. Obtained multispectral images were processed and geo-referenced using Agisoft Metashape Professional 1.7.6 software. Thereafter, multispectral images, which consist of 5 bands, were processed using ERDAS Imagine 2014 software. MATLAB 2009b software was used to perform analytics on resultant images.

### 2.4 Calculation Indices

Normalized Difference Vegetation Index (NDVI) (Koparan, 2019), Bare Soil Index (BSI) (Bhunja *et al.*, 2019), and Modified Secondary Soil Adjusted Vegetation Index (MSAVI2) (Koparan, 2019) associated with each land plot were calculated using processed images from ERDAS Imagine 2014 and MATLAB 2009b software. Following equations show the parameters associated with the calculation of these indices.

$$NDVI = \frac{NIR - Red}{NIR + Red} \quad (2)$$

$$BSI = \frac{[(Red + Green) - (Red + Blue)]}{(NIR + Green) + (Red + Blue)} \times 100 + 100 \quad (3)$$

$$MSAVI2 = \frac{[2NIR + 1 - \sqrt{(2NIR + 1)^2 - 8(NIR - Red)}]}{2} \quad (4)$$

where;

NIR – Near Infrared Radiation

Red – Red Band

Green – Green Band

Blue – Blue Band

## 2.5 Statistical Analysis

Linear regression analysis method was performed to investigate the correlation between the calculated indices and the SOC determined by the laboratory testing. Accordingly, NDVI, BSI, and MSAVI2 indices and respective SOC of the plots obtained from laboratory analysis were graphed using scatter plots. The correlations between these parameters were determined using MATLAB 2009b software.

## 3. Results & Discussion

The SOC of 9 plots was obtained by using laboratory analysis method and indices analysis method and the results are presented by dividing plots into three productivity categories of low, medium and high. Furthermore, by using the index analysis method, the relationships between each productivity category and the obtained SOC were found and represented by graphs (Figure 3, Figure 4 and Figure 5).

### 3.1 Soil Sampling and Analysis

Table 1 shows the mean SOC values of the samples collected from 9 plots according to their three different productivity categories.

Productivity Category	Mean SOC (%)
Low	2.96±0.22
Medium	3.24±0.26

High

3.38±0.22

---

**Table 1: Measured SOC content values of three productivity classes. Each value has been presented as a mean ± standard deviation of them taken from each field plot.**

In accordance with the laboratory analysis method, SOC values obtained in the low productivity zone were in the range of 2.7-3.0% whereas SOC values in the medium and high productivity zones were in the range of 3.0-3.5 and 3.2-3.6%, respectively (Table 1). This shows that there was a positive relationship between SOC and productivity and correspondingly, the lower productivity was observed for the lower SOC soils.

### 3.2 Relationship between Observed SOC and Indices

Table 2 shows the mean NDVI, BSI, and MSAVI2 values obtained for three different productivity classes.

---

<b>Productivity Category</b>	<b>Mean NDVI</b>	<b>Mean BSI</b>	<b>Mean MSAVI2</b>
Low	-0.0197±0.051	-0.031±0.016	-0.014±0.039
Moderate	-0.004±0.077	0.01±0.010	-0.008±0.060
High	-0.133±0.061	-0.006±0.076	-0.091±0.045

---

**Table 2: Descriptive statistics of indices.**

As per the Figure 3, the correlation between NDVI and observed SOC content of each plot was 0.02694. This is relatively a weak relationship and the observed  $R^2$  was 0.0007256 (Table 3). The observed SOC had 0.07% variation with the regression of the mean NDVI of each plot.

When comparing with Figure 4 the correlation between BSI and observed SOC content of each plot was 0.4828. This highlights a moderate relationship between the two variables. There was a 23.3% variation with observed SOC and the regression of the Mean BSI of each plot. The correlation value 0.08582 (Figure 5) showed a weak relationship between MSAVI2 and observed SOC content of each plot. The observed  $R^2$  was 0.007365 and also, there was 0.7% variation with observed SOC and the regression of the mean MSAVI2 of each plot.




Figure 3: Relationship between measured SOC and mean NDVI.

Regression Line Equations	Variables	$R^2$	R
Predicted SOC = 3.200 + 0.090 * NDVI	NDVI	0.0007256	0.02694

Predicted SOC = 3.222 + 3.123 * BSI	BSI	0.2331	0.4828
Predicted SOC = 3.210 + 0.408 * MSAVI2	MSAVI2	0.007365	0.08582

**Table 3: Relationship between observed SOC and indices.**




Figure 4: Relationship between measured SOC and mean BSI




Figure 5: Relationship between measured SOC and mean MSAVI2

Land can be considered as a very essential thing for man. When a land is used for agricultural purposes its soil is the most valuable resource. Therefore, in order to promote productivity in agriculture, it is necessary to assess whether the land is suitable for each crop. There, soil quality can be taken as a major criterion for determination of productivity. SOC is one of the leading indicators of agricultural productivity when it comes to soil quality. Therefore, it is important to detect SOC regularly. In many cases, soil samples are taken from the land and subjected to laboratory testing. But, this is a very expensive and time-consuming task and therefore, present study was conducted to see whether a cost-effective and efficient method like remote sensing

could be used for this purpose without destruction. Accordingly, the ability to capture multispectral images using a drone platform and thereby to detect SOC was carried out by indices analysis.

According to the laboratory analysis method, SOC values obtained in the low productivity zone were in the range of 2.7-3.0% whereas SOC values in the medium and high productivity zones were in the range of 3.0-3.5 and 3.2-3.6%, respectively (Table 1). It was clear that there was a positive relationship between SOC and productivity and accordingly, the lower productivity was observed for the lower SOC soils. It is proven that application of fertilizers has significant effect on the level of SOC and thereby on crop productivity (Studdert, 2000). Therefore, changes of SOC in the sugarcane fields is partially supported by application of fertilizers (Table 4). But, retention of SOC in the different productivity lands might be attributed to the unique features of such lands in particular soil moisture content, clay type, parent material etc.

<b>Plant Crop</b>				
<b>Application</b>	<b>Urea (Kg/ha)</b>	<b>TSP (Kg/ha)</b>	<b>MOP (Kg/ha)</b>	<b>Time Period</b>
Basal Dressing	501	100	75	
Top Dressing 01	1100			45 Days After Planting
Top Dressing 02	150		75	90 Days After Planting
<b>Ratoon Crops</b>				
Top Dressing 01	50	100	75	14 Days After Planting
Top Dressing 02	100			60 Days After Planting
Top Dressing 03	150		75	120 Days After Planting

**Table 4: Fertilizer application of Lanka Sugar Company Ltd., Pelwatte Unit.**

Corresponding to previous analysis conducted by Sugarcane Research Institute of Sri Lanka in Pelwatte Nucleus Estate, SOC percentage of soils have ranged from 0.2 to 1.0% (Crop Nutrition Division, 2022). These values are lesser than that of the current study. The reason might be they have sampled soils from different locations to what we collected and it supports the variability of

SOC in the lands of Lanka Sugar Company Ltd., Pelwatte. Thus, improving SOC content in all lands at least to the values obtained at the locations of the present study will be essential in terms of improving soil quality. On the other hand, it is linked with the productivity of sugarcane (Bot *et al.*, 2005). Minimizing the activities such as burning, tillage, overgrazing and continuous cropping can be advocated to maintain SOC levels at the desired condition in these lands.

According to Bhunia *et al.* (2019), there is a significant correlation between SOC and NDVI ( $R = 0.74$ ,  $P < 0.0075$ ). In contrast, Koparan (2019) shows a weak negative correlation between SOC and NDVI which was obtained from Landsat 8 and PlanetScope multispectral imageries:  $-0.39$  and  $-0.35$ , respectively. However, in this study, there was a very weak direct relationship between mean NDVI and observed SOC of each category.

When considering the relationship between mean BSI and observed SOC of each productivity category (Figure 4), it has given a medium satisfactory correlation *i.e.*  $0.4828$ . Hence, it can be concluded that there is a moderate direct relationship between mean BSI and observed SOC ( $P = 0.1880$ ). Bhunia *et al.* (2019) show a statistically significant relationship between SOC and BSI ( $R = -0.72$ ;  $P < 0.0000$ ). Conversely, Koparan (2019) shows a weak negative relationship as  $-0.44$  between SOC and BSI which has been obtained from both Landsat 8 and PlanetScope satellite imageries.

It is apparent that the amount of reflected and emitted spectra from soil surface is affected by soil humidity by decreasing reflectance values (Nocita *et al.*, 2013). The present study showed a moderate correlation with SOC concentration of lands where moisture content was near field capacity. Contrasting results are reported in some other studies as well (Kumar *et al.*, 2016).

The relationship between mean MSAVI2 and observed SOC of each yield category soil was quite similar to the relationship between mean NDVI and observed SOC. With regards to Figure

5, the correlation is 0.08582 and the  $P$  value was 0.8267. Thus, it is apparent that there was a weak correlation between mean MSAVI2 and observed SOC of each productivity category. According to Koparan (2019), the relationship between SOC and MSAVI2 has a moderate negative correlation ( $R = -0.54$ ,  $P < 0.0001$ ) under both Landsat 8 and PlanetScope satellite multispectral imageries.

Soil organic carbon is associated with soil color, nutrient-holding capacity, and also helps maintain soil composition (Ismail *et al.*, 2012). Under this context, soil color and image intensity relationships can be established between SOC concentration and satellite imagery data. In the present study, high productivity soils had a dark brown color and the medium and low productivity lands had a reddish brown and light brown color, respectively. This is in agreement with Castaldi *et al.* (2016) and accordingly, soils with dark hues have higher SOC contents when compared to the soils with bright hues in the case of equal moisture and the same parent materials.

BSI regression equation for SOC detection developed in the present study can be used to determine SOC in land with bare soil. The amount of soil samples required to precisely estimate SOC levels for a region will need to be determined in future research work. Further, drone images are least subject to the effects of atmospheric effects. The present findings will enhance and deliver more accurate soil data. In fact, this study will help researchers and farm managers to choose the best management strategies for their work, farm operations, and research. It will also help preserve SOC contents and improve precision agriculture farming systems.

#### 4. Conclusion

SOC content of the sugarcane lands related to the study as assessed by soil samples collection and laboratory test is in the favorable range for sugarcane cultivation. The study found that NDVI and MSAVI2 indices show a weak relationship with the measured SOC in the selected study area and a moderate relationship with BSI. Therefore, further studies should be conducted to determine how different vegetation indices would affect the estimation of SOC.

BSI is suitable for SOC detection by multispectral images of a land. But, this index should be further enhanced in future studies when predicting SOC in sugarcane lands.

It is also desirable to see how effective the contribution of Bright Index (BI) is to identify salt spot areas in multispectral images. Also, it is worthwhile to improve NDVI and MSAVI2 indices and examine their applicability for sugarcane plots.

#### References

Bhunias GS, Kumar SP, Pourghasemi HR. Soil organic carbon mapping using remote sensing techniques and multivariate regression model. *Geocarto International*. 2019;34(2):215–226. doi: 10.1080/10106049.2017.1381179.

Bot A, Benites J. *The importance of soil organic matter: Key to drought-resistant soil and sustained food production*. 2005;80. Food & Agriculture Org.

Castaldi F, Chabrillat S, Jones A, Vreys K, Bomans B, Van WB. Soil organic carbon estimation in croplands by hyperspectral remote APEX data using the LUCAS topsoil database. *Remote Sensing*. 2018;10(2). doi: 10.3390/rs10020153.

Castaldi F, Palombo A, Santini F, Pascucci S, Pignatti S, Casa R. Evaluation of the potential of the current and forthcoming multispectral and hyperspectral imagers to estimate soil texture and organic carbon. *Remote Sensing of Environment*. 2016;179:54-65. doi:10.1016/j.rse.2016.03.025.

Chaminé HI, Pereira AJ, Teodoro AC, Teixeira J. Remote sensing and GIS applications in earth and environmental systems sciences. *SN Applied Sciences*. 2021;3:1-3. doi:10.1007/s42452-021-04855-3.

Crop Nutrition Division (2020). Crop Nutrition – Sugarcane Research Institute, Sri Lanka. 2020. Accessed 31 Oct. 2022. Available at: <https://sugarres.lk/research-development/main-research-programs-divisions/crop-nutrition>.

Eswaran H, Lal R, Reich PF. Land degradation: an overview. *Response to land degradation*, 2019;20-35.

Gomez C, Viscarra RRA, McBratney AB. Soil organic carbon prediction by hyperspectral remote sensing and field vis-NIR spectroscopy: An Australian case study. *Geoderma*. 2008;146(3–4):403–411. doi: 10.1016/j.geoderma.2008.06.011.

Ismail M, Yacoub RK. Digital soil map using the capability of new technology in Sugar Beet area, Nubariya, Egypt. *The Egyptian Journal of Remote Sensing and Space Science*. 2012;15(2):113-124. doi:10.1016/j.ejrs.2012.08.001.

Kath J, Le Brocque AF, Reardon-Smith K, Apan A. Remotely sensed agricultural grassland productivity responses to land use and hydro-climatic drivers under extreme drought and rainfall. *Agricultural and Forest Meteorology*. 2019;268:11-22. doi:10.1016/j.agrformet.2019.01.007.

Kheir RB, Greve MH, Bøcher PK, Greve MB, Larsen R, McCloy K. Predictive mapping of soil organic carbon in wet cultivated lands using classification-tree based models: The case study of

Denmark. *Journal of Environmental Management*. 2010;91(5):1150-1160. doi: 10.1016/j.jenvman.2010.01.001.

Koparan MH. Estimating soil organic carbon in cultivated soils using soil test data, remote sensing imagery from satellites (Landsat 8 and PlantScope), and web soil survey data. *Electronic Theses and Dissertations*. 2019;3177. Available at: <https://openprairie.sdstate.edu/etd/3177>.

Kumar P, Pandey PC, Singh BK, Katiyar S, Mandal VP, Rani M. et al. Estimation of accumulated soil organic carbon stock in tropical forest using geospatial strategy. *The Egyptian Journal of Remote Sensing and Space Science*. 2016;19(1):109-123. doi:10.1016/j.ejrs.2015.12.003.

Mng'ong'o M, Munishi LK, Blake W, Comber S, Hutchinson TH, Ndakidemi PA. (2021). Soil fertility and land sustainability in Usangu Basin-Tanzania, *Heliyon*. 2021;7(8): p.e07745. doi: 10.1016/j.heliyon.2021.e07745.

Montfort F, Begue A, Leroux L, Blanc L, Gond V, Cambule AH, et al. From land productivity trends to land degradation assessment in Mozambique: Effects of climate, human activities and stakeholder definitions. *Land Degradation & Development*. 2021;32(1):49-65. doi:10.1002/ldr.3704.

Mugiyo H, Chimonyo VG, Sibanda M, Kunz R, Masemola CR, Modi AT. et al. Evaluation of land suitability methods with reference to neglected and underutilised crop species: A scoping review. *Land*. 221;10(2);125. doi:10.3390/land10020125.

Ngatia LW, Moriasi D, Grace III JM, Fu R, Gardner CS, Taylor RW. Land use change affects soil organic carbon: An indicator of soil health. *Environmental Health*. 2021. doi:10.5772/intechopen.95764.

Nocita M, Stevens A, Noon C, van Wesemael B. Prediction of soil organic carbon for different levels of soil moisture using Vis-NIR spectroscopy. *Geoderma*. 2013;199:37-42. doi:10.1016/j.geoderma.2012.07.020.

Schoonover JE, Crim JF. An introduction to soil concepts and the role of soils in watershed management. *Journal of Contemporary Water Research & Education*. 2015;154(1):21-47. doi:10.1111/j.1936-704X.2015.03186.x.

Studdert, GA. Crop rotations and nitrogen fertilization to manage soil organic carbon dynamics. *Soil Science Society of America Journal*. 2000;64(4):1496-1503. doi:10.2136/sssaj2000.6441496x.

Vašát R, Kodešová R, Borůvka L, Klement A, Jakšík O, Gholizadeh A. Consideration of peak parameters derived from continuum-removed spectra to predict extractable nutrients in soils with visible and near-infrared diffuse reflectance spectroscopy (VNIR-DRS). *Geoderma*. 2014; 232:208-218. doi:10.1016/j.geoderma.2014.05.012.

World Bank. Soil Organic Carbon MRV Sourcebook for Agricultural Landscapes ,World Bank, Washington, DC; 2021.