

Original Research Article
ON THE EXISTENCE OF A HOLLOW NEUTRON STAR

Abstract:

Based on the Ni's solution of Einstein Field Equations (EFE) we discuss the possibility of formation of a new type Neutron Star in which case the maximum density appears at a distance far from its center (not at the center for normal neutron star case). As the inner core of such Neutron Star is a hollow sphere consideration of vacuum energy at the core offers the possibility of two types neutron star models — one with no exotic phase at the core and another one with a significant portion of it in the new phase. Ni's solutions of EFE provide a new type neutron star model with hollow sphere in the inner core without violating the general relativity conditions. It is proposed that existence of such new kind neutron star is physically possible. It is suggested that existence of this type Neutron Star can be detected through observations of electromagnetic counterpart by LIGO and VIRGO while theoretical studies can increase our knowledge of understanding this new, unknown stream of relativistic compact objects (neutron star) in astrophysics. Some problems related to this hollow neutron star are discussed.

Key Words : Field Equations, Exact solutions, Relativistic stars, Neutron star core, Asymmetric matter

1. INTRODUCTION

In 1915 Einstein General theory of Relativity (GTR) introduces the idea that gravitation is not a force, rather it is a consequence of curvature in a four dimensional space-time [1a,b] i.e., Matter tells space how to curve and space tells matter how to move [2].

Thus solutions of Einstein's Field Equations (EFE) of general relativity provide the picture of space-time which relate the Einstein tensor to stress-energy tensor representing the distribution of energy, momentum and stress in the space-time manifold. The fact is that the Einstein's field equations alone are not enough to determine the evolution of a gravitational system in many cases where dependency on the dynamics of matter and energy are relevant i.e. on the gravitational field. Since the Einstein's field equations are a system of coupled non-linear partial differential equations, several effective techniques / approximations offer different exact solutions (Lorentz metrics) which are comfortable to express a physically realistic stress-energy tensors which are in general found in complex, non-physical stress in space. This means that such types of materials that would go into the make up of such spaces are seldom found in nature (practically impossible to create such situation on the earth), it is frequently desired to understand the geometrical properties of such space of a given mass distribution. So, exact solutions of Einstein's field equations play important role in physical problems [3].

In the year 1916 Schwarzschild [4] first gave the solution (interior and exterior) of the Einstein's field equation with a prediction on the existence of a black-hole. This creates interest on studies of stellar compact objects in the realm of relativistic astrophysics. Searching for the exact solutions and the best of these solutions provide a limit on certain physical parameters (such as mass, radius, etc.) as well as the behavior of gravitational field. For this reason the

construction of exact solutions, therefore, can be used for physical modeling and there after to predict the measurable quantities about physical systems (compact objects such as white dwarfs, neutron stars,....).

2. Ni's Solution of EFE

In 2011 Ni published the result of his solution of the Einstein's Field Equations, based on Tolman-Oppenheimer-Volkoff equations [5,6], for the structure of stable neutron stars [7].

Considering the general static metric for stars (inside neutron stars) with spherical symmetry

$$ds^2 = - e^\lambda dr^2 - r^2 d\theta^2 - r^2 \sin^2\theta d\phi^2 + e^\nu dt^2 \quad (1)$$

$$\text{and the auxiliary function } u = r (1 - e^{-\lambda}) / 2 \quad (2)$$

he obtained the field equations (in $c = G = 1$ units) as

$$(du / dr) = 4\pi \rho r^2 \quad (3)$$

$$(dp / dr) = - \{ (p + \rho) / r (r - 2u) \} (4\pi p^2 r^3 + u) \quad (4)$$

Where ρ and p are the macroscopic energy density and pressure measured in proper coordinators, respectively. Note that equ. (3) and (4) are the Tolman-Oppenheimer-Volkoff equations.

For neutrons, inside the neutron star, with rest mass m_0 obeying the Fermi Dirac statistics, he used parametric form of ρ and p with the parameter 't' related to the Fermi momentum P_F as

$$t = 4 \ln \{ P_F / m_0 c + [1 + (P_F / m_0 c)^2]^{1/2} \} \quad (5)$$

$$\text{energy density } \rho = K (\text{sinht} - t) \quad (6)$$

$$p = \frac{1}{3} K (\text{sinht} - 8 \sinh t/2 + 3 t) \quad (7)$$

where $K = \pi m_0^4 c^5 / (4h^3)$ and the density of the particle number $n = N/V = 8\pi P_F^3 / 3 h^3$.

Putting the boundary condition for neutron star with the initial values $u = u_i$, $p = p_i$ at $r = 0$

to the radius $r = r_o$ where $\rho = 0$ at the surface boundary of the star he obtained a new type solution i.e. there is a void at $r = 0$ (i.e. at the centre of the neutron star) though the solutions satisfied the Einstein's equation i.e. from $r = 0$ to $r = r_i$, $\rho = 0$ and $p = 0$. This means that there is no particles and thus pressure is zero in this void region. Beyond this void, from $r = r_i$ to $r = r_o$ $\rho \geq 0$ and $p \geq 0$ the density and pressure increase first from zero at $r = r_i$. After reaching a maximum, ρ and p then decrease. At the outer boundary $r = r_o$, p and ρ reach zero i.e.

- a) after reaching a maximum, p decreases to zero at the outer radius r_o ,
- b) the structure parameter 't' increases from zero at the inner radius r_i to a maximum and then decreases to zero at the outer radius $r = r_o$ (see figure 1)
- c) His solutions without maximum mass limit (i.e., no upper mass limit of these objects) do not depend on a special property of the equation of state for the star matter.
- d) The outer physical surface of these stars (i.e., neutron stars) is always situated above the corresponding event horizon.
- e) The solutions with void inside the center of the star emerge naturally,
- f) In the case when the void is very small, even then the particle number 'N' and the mass 'm' of the star do not have a maximum limit.
- g) His solution leads to a mechanism for neutron stars to avoid collapse into black holes but remain regular.

3. Vacuum Energy Concept

The recent observations of gravitational waves (GWs) open a new direction for exploring fundamental physics. It has been speculated earlier that there may be a new phase of nuclear matter at the core of the neutron stars [9 – 11]. If such a phase exists in the interior of the neutron

star then

- (a) it is expected vacuum energy (VE) at the center of it [12];
- (b) the presence of vacuum energy in the inner core of neutron star indicates a new QCD phase at large densities as well as appearance of vacuum energy in the equation of state (EOS) for a new phase.

Not only that, this will make neutron star the only known objects where vacuum energy might make up a non-negligible fraction of the total mass [13].

Using the solutions of the Tolman-Oppenheimer-Volkoff (TOV) equations [5,6] for spherically symmetric and mass distribution of the star (neutron star) and assuming the presence of vacuum energy in the inner most layer of the neutron star Csáki et al [13] recently investigated for finding the contribution of a vacuum energy in exotic high density phases in different EOSs for modeling and the maximum stable mass of neutron stars. In that study their significant findings are:

- i. the compactness of the neutron star can be characterized by the radius of a neutron star with a fixed mass ;
- ii. For a given mass, there are two possible types of stars — a) one with no exotic phase in the core and another with a significant portion of the star in the new phase.

In other words, introducing the non-zero vacuum energy in the interior of of neutron star can create a disconnected branch of stable neutron star solutions that allow the possibility of having two neutron stars of the same mass but with significantly different radii.

Based on the Ni's solutions of EFE Neslušan [14] estimated the distribution of matter inside

the star for constructing the neutron star model. The rest mass M_o (i.e. the sum of the rest masses of all neutrons constituting the object) of the neutron star he estimated using the traditional theory of these objects (i.e. Misner et al [2]) as

$$M_o = 4\pi m_n \int_{R_{in}}^{R_{out}} n r^2 / \sqrt{1 - 2u/r} . dr \quad (8)$$

where $n = n(r)$ the number density of neutrons given by the relation

$$n = (8\pi / 3 h^3) (m_n c)^3 \sinh^3 t/4 \quad (9)$$

Using the Ni's solutions the significant results obtained by Neslušan are :

- I. There is a variety of neutron stars constituted by the same number of neutrons i.e. for a given rest mass a variety of neutron star models can be constructed with the different other characteristics. For example, for the rest mass $0.669 M_{\square}$ they obtained two neutron star models with the same distance r_o as

Table 1. Model description

r_o (km)	NS model 1		NS model 2	
	r_{in} (km)	r_{out} (km)	r_{in} (km)	r_{out} (km)
0.001	2.824×10^{-11}	12.119	8.981×10^{-11}	6.8558
0.01	2.824×10^{-8}	12.119	8.981×10^{-8}	6.8558
0.1	2.820×10^{-5}	12.119	8.926×10^{-5}	6.8538
0.5	4168×10^{-3}	12.159	1.0301×10^{-2}	6.6126
1.0	4738×10^{-2}	12.391	8.178×10^{-11}	4.7028

- II. Neutron star model is a hollow sphere with a cavity in its interior and the gravitational attraction in the cavity is outward oriented.
- III. The models of compact objects are with no singularity other than the Big Bang type

singularity which can exist in the universe i.e. the conditions are realistic.

- IV. The concept of hollow sphere, as already claimed by Ni, is able to a model of stable compact object of whatever large mass i.e. this model is valid not only for the full sphere concept of neutron stars but also valid for in the astrophysics of real objects

4. Normal Neutron Star Vs Central Hollow Sphered Neutron Star

In 1967 the discovery of Pulsars [15, 16] provided the first evidence of the existence of neutron star. Pulsars are neutron stars neutron stars that pulses of radiation once per rotation. Neutron stars are a class of extremely dense, compact stars, though to be composed primarily of neutrons. Normal neutron stars are typically about 20Km (12 miles) in diameter having mass ranges between $1.8 M_{\square}$ and $1.97 M_{\odot}$ but most are $1.35 M_{\odot}$. The average densities of these stars are extremely high — about $10^{14} \text{ g.cm}^{-3}$. The boundary surface of this star is solid, intermediate layers are mostly neutrons and are probably in the super-fluid, super-conductor states [17, 18]. It is not yet known definitely what is at the centre of the neutron star [19]. Theoretical studies suggest [20 – 22] hyperons, kaons, pions, quark matter are at the inner core of the normal neutron star.

In 2011 Jun Ni [7] published the solution of Tolmann- Oppeneimer- Volkoff equations and suggested this new kind stable neutron star which has a hollow sphere with the inner physical surface and cavity inside. Ni's solution based compact stars are neutron stars having a hollow sphere at the inner core this star. This type compact stars are unknown today but observations of LIGO and VIRGO as well as space based NICER [23,24] will definitely detect this kind of compact stars and theoretical studies will increase our knowledge of understanding such type neutron stars. A comparison of physical properties of the normal neutron star and hollow sphere neutron star are given below :

UNDER PEER REVIEW

Table 2. Comparative analysis between Normal and Hollow Neutron Star

Normal Neutron Star	Hollow Neutron Star
<p>1) Normal massive neutron stars are formed from supernova explosions.</p> <p>2) Solutions of EFE correspond to neutron stars without a void at their centers and offer maximum density in the core.</p> <p>3) There is uncertainty in maximum mass limit of neutron stars, exceeding which it would collapse into blackholes and disappear. As per observation the maximum mass of neutron star obtained till date is $2.14 M_{\odot}$ (PSR J0740+6620) [25,26]. On the other hand, observations by LIGO and VIRGO claims the mass limit for the compact object detected in the binary merger GW190814 is in the range $2.5 - 2.67 M_{\odot}$ [24,27]. But the detected object is either the most massive neutron star or the least massive blackhole. Theoretical studies [28] extend the mass limit up to $3.2 M_{\odot}$.</p> <p>4) The maximum mass limit of the neutron star depends on the choice Equation of States (EoS) of the star matter.</p>	<p>1) Neutron stars with Hollow sphere are formed possibly from binary neutron stars merger</p> <p>2) Ni's solutions [7] of EFE correspond to neutron stars with a void at their centers and maximum density occurs beyond the core.</p> <p>3) Ni's solutions show without maximum mass limit for TOV equations i.e. there is no upper mass limit of these objects. These compact objects (Neutron stars) can accommodate infinite number of neutron particles. Even in the case that the void is very small, the particle number N and the mass 'm' of the star do not have a maximum limit.</p> <p>4) Solutions without maximum mass limit do not depend on the special property of EoS of the star matter.</p>

<p>5) In the case of normal neutron star matter density gradually increases from the surface boundary of the star and then reaching maximum at the centre of the star.</p> <p>6) Asymptotic freedom behavior appears in the core of the star [29,30]</p> <p>7) Exceeding the maximum mass limit normal neutron star collapses into black-hole and disappear.</p>	<p>5) In this case the stars do not have a maximum mass limit which leads to a mechanism for neutron stars to avoid collapse into black holes and remain regular.</p> <p>6) No appearance of asymptotic freedom phase in the interior of this neutron star.</p> <p>7) Matter density starts increasing from the boundary surface , after a few km away maximum density arises and then again decreases gradually upto the inner radius of the hollow sphere / core.</p>
--	---

Conclusion

Regarding compact objects (such as white dwarf, neutron star) we generally understand that its internal density increases from the star's boundary surface towards the centre of the object and density is maximum at the core due strong gravitational attraction. Ni's solutions of EFE [7] provide a breakthrough idea in understanding the compact objects with hollow sphere in its interior at the core. Introducing the presence of vacuum energy in the inner cores of neutron stars Csáki etal [13] stress the existence of the new phase, due QCD phase transition , which requires a modification of the traditional concept of compact object i.e. a non-zero vacuum energy can create a disconnected branch of stable neutron star solutions which allow the possibility of having two neutron star models for a given mass with

significantly different radii —

- a) Neutron star with no exotic phase in the core , and
- b) Another one with a significant portion of it in the new phase.

Neslušan's study [14], based on Ni's solutions, points out that the distribution of matter which is observed as spherically symmetric when the observer is in its center, is no longer observed as spherically symmetric by the observer aside the center in a curved space-time of general relativity. His analysis suggest that in the interior of such type hollow sphere neutron stars there is a critical distance in which the partial gravitational actions of both upper and lower layers equal each other but in the stable configuration this gravity is again balanced by the gradient of pressure which is always oriented against the gravity. Since the concept of hollow sphere model is applicable for stable compact object of whatever large mass, this is also valid only for full-sphere concept of neutron star, but also valid in the astrophysics of real objects [31] such as pulsars with extraordinary large mass e.g. J1748-2021B, J1311-3430, B 1957 + 20 with masses $(2.5 - 2.9) M_{\odot}$, $(2.1 - 2.8) M_{\odot}$ and $(2.1 - 2.7) M_{\odot}$, respectively [32] ; ultra-luminous objects like quasars etc. Now question is detection of such hollow sphere neutron stars. As neutron star mergers chirp about vacuum energy (as suggested by Csáki etal [13]) so observations of electromagnetic counterparts from neutron star mergers through LIGO and VIRGO offer the access to detect such type objects. Numerical simulations of neutron star mergers can provide the required information in this matter [33- 35]. Besides observation, theoretical studies of hollow sphere neutron stars can also increase our knowledge of this new type unknown relativistic compact object (neutron star) in astrophysics.

The significance of this central hollow neutron star is ----

- a) for equilibrium mass distribution it has no upper limit mass i.e., without a maximum mass

limit.

ii) These stars are with a void inside their centers. But one of the most important predictions of general relativity is the existence of maximum mass for any static matter configuration.

Therefore, several problems arise regarding hollow neutron stars :

- i) As these neutron stars have no upper mass limit these cover up the mass gap region $(2-5) M_{\odot}$ although masses of heaviest neutron stars, hybrid stars and high mass strange quark stars cover up to $2.6 M_{\odot}$ [27] and Triaxial type [36,37] quark star up to $3.3 M_{\odot}$ [38] provided the secondary compact star in the binary merger GW190814 be the heaviest neutron star. The remnant of GW190814 remains debated whether it is a heaviest neutron star or lightest black hole ?
- ii) A new evolutionary theory is required to explain the formation of black holes from massive neutron stars.
- iii) The central part of the hollow neutron star is vacuum theoretically, but in realistic case may contain which type deconfined quark matter not yet certain.
- iv) How to distinguish between the normal neutron star and hollow neutron star through their properties ?
- v) Is the hollow neutron star be a good source of gravitational waves ? etc.


Ethical statement : None

References


- 1 .a) Einstein A. *Die Feldgleichungen der Gravitation, Sitzungsberichte der Königlich Preßischen Academie der Wissenschaften* (Berlin), 1915 : p. 844-847 ;
b) Einstein A. *Annalen der Physic.* 1916 ; 354 : 769
2. Misner C W, Thorne K S, Wheeler J A. *Gravitation*, 2nd Edition, (W H Freeman, NY, 1973)
3. Hewett L D. Ph.D. Thesis “A Solution to Einstein Gravitational Field equations for a Spherically Symmetric static perfect fluid” , (Missouri S & T Library and Learning Resources , 1965)
4. Schwarzschild K. Sitzungsberichte der Königlich Preßischen Academie der Wissenschaften zu Berlin, *Phys. Math. Classe* 1916 ; 1 : 189 – 196
5. Oppenheimer J R , Volkoff G M. On the Massive Neutron Core *Phys. Rev.* 1939 ; 55 : 74
6. Tolman R C. Static Solutions of Einstein's Field Equations for Spheres of Fluid (1939)
Phys. Rev. 1939 ; 55 : 364
7. Ni J. Solutions without a maximum mass limit of the general relativistic field equations for neutron stars. *Science China* . 2011 ; 54 : 1304
8. Abbott B P. et al. {LIGO and VIRGO collaboration}. GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral. *Phys. Rev. Lett.*, 2016 ; 119 : 161101
9. Ivanenko D D, Kurdgelidze D F. *Astrofizika* 1965 ; 1 : 479
10. Pacini F. High-energy Astrophysics and a Possible sub-nuclear Energy Source. *Nature* 1966 ; 209: 389
11. Itoh, N. Hydrostatic Equilibrium of Hypothetical Quark Stars. *Prog. Theor. Phys.* 1970 ; 44, 291
12. Ellis J R, Kapusta J I, Olive K A. Phase transition in dense nuclear matter with quark and gluon condensates . *Phys. Lett. B.* 1991; 273 : 123
13. Csáki C, Eröncel C, Hubisz J., Rigo G, Terning J. *J. High Energy Phys.* 2018; 9 : 087
14. Neslušan, L. The Ni's Solution for Neutron Star and Outward Oriented Gravitational Attraction in Its Interior. *J. Mod. Phys.* 2015 ; 6, 2164
15. Hewish A, Bell S J, Pilkington J D, Scott P F, Collins R A . Observation of a Rapidly Pulsating Radio Source. *Nature* 1968 ; 217 : 709
16. Hewish A. *Pulsating Star* (Plenum Press, NY, 1968) p.vii & 5.

17. Baym G, Pethick C J, Sutherland P. The Ground State of Matter at High Densities: Equation of State and Stellar Models . *Astrophys. J.* 1971 ; 170 : 299
18. Haensel P, Potekhin A Y, Yakovlev D G. *Neutron Star I: Equation of State and Structure* (,Springer, NY., 2006)
19. Khanna, K M, Kandie D K, Tonui J K, Cherop H K. Incommensurate Crystallization of neutron matter in Neutron Star. *East Euro. J. Phys.* 2020 ; 2 : 57
20. Chamel N, Haensel P. Physics of Neutron star Crusts. *Living. Rev. Relativity.* 2008 ; 1 : 10
21. Mann, A. The golden age of neutron-star physics has arrived. *Nature.* 2020; 579 : 20
22. Potekhin A Y. *The Physics of Neutron Star* . 2011 ; arxiv: 1102.5735
23. Weber F, et al. Phases of Hadron-Quark Matter in (Proto) Neutron Stars. *Universe* 2019 ; 5:169
24. Godzieba D A, Radice D, Bernuzzi S. On the Maximum Mass of Neutron Stars and GW190814. *Astrophys. J.* 2021 ; 908 : 122
25. Cromartie H T E, Fonseca S M, Ransom . et al. Relativistic Shapiro delay measurements of an extremely massive millisecond pulsar . *Nature Astronomy.* 2020; 4 : 72
26. Roupas Z. Secondary component of gravitational-wave signal GW190814 as an anisotropic neutron star. *Astrophys. Space Sci.* 2021 ; 366 : 9
27. Abbott R P. et al. GW190814: Gravitational Waves from the Coalescence of a 23 Solar Mass Black Hole with a 2.6 Solar Mass Compact Object. *Astrophys. J.* 2020 ; 896 : L44
28. Rhoades C E, Ruffini R. Maximum Mass of a Neutron Star. *Phys. Rev. Lett.* 1974 ; 32 : 324
29. Gross D J, Wilczek F. Ultraviolet Behavior of Non-Abelian Gauge Theories *F. Phys. Rev. Lett.* 1973; 30 : 1343
30. Gross, D. J. Twenty five years of asymptotic freedom. *Nucl. Phys. Proc. Suppl.* 1999 ; 74 : 426
31. Neslušan, L. Applicability of the NI's solution of the Einstein Field Equations to the real Objects, in *Proc. of IAU Symp.: New Frontiers in Blackholes Astrophysics.* held in Ljubljano, Slovenia , No-324 ; 2018
32. Watt, A. et al. Advancing Astrophysics with the Square Kilometre Array, ((Proc. of the conference held in Giardini, Nxos, Italy, 2014)
33. Seidov Z F. The Stability of a Star with a Phase Change in General Theory of Relativity. *Soviet Astronomy* . 1971 ; 15 : 347
34. Shaeffer R, Haensel P, Zdunik L. Phase transitions in stellar cores. I - Equilibrium configurations. *Astron. Astrophys.* 1983 ; 126 : 121

35. Lindblom L. Phase Transitions and the Mass-Radius Curves of Relativistic Stars.
Phys. Rev. D. 1998 ; 58 : 024008
36. Uryu K, Tsokaros A, Baiotti L. et al. Do triaxial supramassive compact stars exist?
Phys. Rev. D 2016 ; 94 : 101302
37. Parui, R. K. A New Compact Star --- the “Triaxial Star” --- and the Detection of a
Cosmic Baby : A Possibility. Int. Astron. Astrophys. Res. J. 2023 ; 5 : 38
38. Zhou E, Tsokaros A, Rezzolla L, Xu R, Uryu K. Uniformly rotating, axisymmetric and
triaxial quark stars in general relativity. Phys. Rev. D 2018 ; 97, 023013



(A)



(B)

Fig.1 : Schematic diagram showing (A) the structure of a hollow sphere neutron star (not scaled) matching with (B) the density ' ρ ' – radius ' r ' curve as per Ni's solution.

$r_c \rightarrow$ Center $r_{in} \rightarrow$ inner radius, $r_{out} \rightarrow$ outer radius of the hollow Neutron Star