

## Original Research Article

### Efficacy of different plant extracts and insecticide on Red flour beetle, *Tribolium castaneum* (Herbst) in Wheat, *Triticum aestivum*

#### ABSTRACT:

The studies on the efficacy of different plant extracts and insecticide on Red flour beetle, *Tribolium castaneum* (Herbst) in Wheat, *Triticum aestivum* were carried out to know the impact of above plant extracts on percent mortality and Percent weight loss. The research was carried out at Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, U.P. in Department of Entomology [2022]. This experiment was conducted under Complete Randomized Design (CRD). The experiment was conducted on effect of plant extracts and insecticide on the mortality of adult *Tribolium castaneum* at 5, 10 and 15 days of exposure. Among the treatments with botanicals the more number of mortality is observed in the Neem leaf powder (65.55%) followed by Eucalyptus leaf powder (55.55%) and least mortality is observed in the turmeric treatment (20%). Experiments conducted on weight loss of wheat grains at 30, 60 and 90 days revealed that the maximum weight loss is occurred in turmeric treatment (1.2%) followed by ginger rhizome powder (0.85%) and minimum is recorded in neem leaf powder (0.34%). Therefore, the present laboratory studies clearly showed the efficacy of different plant extracts on red flour beetle *Tribolium castaneum* (Herbst) were almost effective when compared with chlorpyrifos (Insecticide). Hence it may be concluded that these plant extracts are eco-Friendly, cost effective and easily available. It can easily be incorporated for the management of *Tribolium castaneum* in wheat.

**KEYWORDS:** Chlorpyrifos, Mortality, Red flour beetle, Weight loss, Wheat

#### Introduction:

Wheat is a grass widely cultivated for its seed, a cereal grain which is a worldwide staple food [Mauseth, J. D. (2014)]. The many species of wheat together make up the genus *Triticum* the most widely grown is common wheat (*T. aestivum*). Wheat is grown on more land area than any other food crop, 220.4 million hectares or 545 million acres, (FAOSTAT). World trade in wheat is greater than for all other crops combined. In 2020, world production of Wheat was 761 million tonnes making it the second most produced cereal after Maize. India's Wheat production during the current crop year (July 2022- June 2023) has been estimated at a record high of 112.18 million tones (mt) – 5 percent higher than 106.84 mt last crop year. Since 1960, world production of

**Comment [CM1]:** This paper required a language revision

**Comment [CM2]:** Abstract appears very short. Hence it may be elaborated little more keeping maximum word limit of 250. A general structure of abstract is as under:  
Initial 1-2 sentences about introduction, importance and relevance of the study; Next 1-2 sentences should tell what was done and what were the objectives; Next 1-2 sentences on methodological approaches followed in the study; Next 2-3 sentences should tell key findings of the study; Next 1-2 sentences should highlight the significance of the current findings; Last sentence as conclusion/recommendations/importance of the findings of this study

**Comment [CM3]:** Make the problematic and the objective more clear

**Comment [CM4]:** PLAGIAS hear you must reform this centence (source/Top 24 papers published in the topic of Rice weevil in 2021 (typeset.io))

**Comment [CM5]:** Observation when you put full stop this mean that the idea is finished an citation my necessary

**Comment [CM6]:** REFORMULE AND INSERT THE REFERENCE: there is a plagiat here from this source: <https://eol.org/pages/108051/articles>

**Comment [CM7]:** Plagiat source <https://eol.org/pages/108051/articles>

**Comment [CM8]:** Source required: <https://simple.wikipedia.org/wiki/Wheat>

**Comment [CM9]:** CITATION AND REFORMULATION IS REQUIRED: <https://tuscany2018.iot.ieee.org/about/read-abo-ut-...>

wheat and other grain crops has tripled and is expected to grow further through the middle of the 21<sup>st</sup> century. Global demand for wheat is increasing due to the unique viscoelastic and adhesive properties of gluten proteins, which facilitate the production of processed foods, whose consumption is increasing as a result of the worldwide industrialization process and the westernization of the diet. (Shewry and Hey, 2015). Wheat is an important source of carbohydrates. Globally, it is the leading source of vegetable protein in human food, having a protein content of about 13%, which is relatively high compared to other major cereals but relatively low in protein quality for supplying essential amino acids. When eaten as the whole grain, wheat is a source of multiple nutrients and dietary fiber. (Shewry and Hey, 2015). The red flour beetle (*Tribolium castaneum*) is a species of beetle in the family Tenebrionidae, the darkling beetles. It is a worldwide pest of stored products, particularly food grains, and a model organism for ethological and food safety research (Grunwaldt et al., 2013). Both larvae and adults feed on grain dust and broken grain, but not the undamaged whole grains and spend its entire life cycle outside the grain kernels (Karunakaran et al., 2004). In severe infestation, the flour turns greyish and has a pungent, disagreeable odour- making it unfit for human consumption. This insect causes substantial loss in storage because of its high reproductive potential (Prakash et al., 2008). The control of arthropod pests on stored products has been made by using of fumigants and residual chemical insecticides. Fumigation by phosphine is the main approach for the operative control of the insect pests (Chadda, 2016). The recent emergence of heritable high level resistance to phosphine in stored grain pests is a serious concern among major grain growing countries around the world. (Jagadeesan et al., 2012) Botanicals have low mammalian toxicity, easy to use, biodegradable and moreover no harm to the environment. Over 120 plants and plant products can be used for the control of stored grain Insect pests. (Ali et al., 2014). Reported that botanicals can be used as effective tool against *Tribolium castaneum* along with other IPM tactics. Application of plant-derived materials (i.e., plants powders, extracts, oils) as fumigants for the control of stored commodities insect pests and their products is potential substitute approach for the control of insect pests (Rehman et al., 2020). In this present investigation efficacy of different botanicals and insecticide application to the red flour beetle is observed.

### Materials and Methods:

Studies on the “Efficacy of different plant extracts and insecticide on Red flour beetle *Tribolium castaneum* (Herbst) in Wheat.” were conducted at Entomology department Shuats, Prayagraj during 2022. To assess the effect of different plant extracts and insecticide on percent mortality of *Tribolium castaneum* and weight loss of wheat grains.

### Insect culture : Rearing of test insect *Tribolium castaneum*

Comment [CM10]: Reformulation And Citation Is Required From <https://eol.org/pages/108051/articles>

Comment [CM11]: Reformulation And Citation Is Required From <https://identainternational.info/>

Comment [CM12]: Reformulation And Citation Is Required From <https://evergrowfert.com/knowledge/wheat/>

Comment [CM13]: Reformulation And Citation Is Required From <https://www.inaturalist.org/taxa/232865-Tribolium-...>

Comment [CM14]: Reformulation Is Required To Avoid Plagiat

Comment [CM15]: Introduction section is very important, it should be made more specific and relevant the current study. Most up-to-date review of literature should form the basis for necessity of this work, particularly in AREA context. Please add 2-4 lines as last para, what was the lacuna in existing scientific knowledge which motivated you to undertake this study. What is the current status of studies ON Impact of Efficacy of different plant extracts and insecticide on Red flour beetle?? It very important to clarify why you use the pesticides and plants extract

Comment [CM16]: Enumerate titles

The cultures of *Triboliumcastaneum* were obtained from local wheat godowns of Prayagraj. Plastic containers of 1.5 kg capacity were used for insect rearing. About 500gm of Wheat grains were kept in each container and about 600 adults of insects were released separately. They were allowed to lay eggs for 3 to 5 days and removed after 7 days, These containers were kept at room temperature for the adult emergence of *Triboliumcastaneum*.

### Preparation of plant products

After collecting the leaves, washed thoroughly and then air-dried under shade. The dried leaves were ground to a fine powder with the help of grinding machine or pestle and mortar. The powder was passed through one mm sieve to get the fine powder of uniform particle size in all cases along with chemical, Chlorpyrifos and seeds.

**Comment [CM17]:** CITE THE place where you get your plants leaves

### Mixing of grain protectants

100gm of healthy and uninfested healthy grains will be taken in plastic jars and different botanicals will be mixed according to required quantities. Freshly emerged 10 beetles are released in the jars. Jars were covered with muslin cloth and tied with rubber bands. Observations were recorded at 5 and 10 and 15 days for Mortality. Similarly, Observations were made for Grain weight loss for 30,60 and 90 days.

**Comment [CM18]:** Clarify what is your source or reference you base on as protocol

### Observations to be recorded:

- Mortality percent of red flour beetle *Triboliumcastaneum*(Herbst)

$$\text{Percent mortality} = \frac{\text{Number of dead insects}}{\text{Total number of insects release}} \times 100$$

- Percent of weight loss of treated grain

$$\text{Percent weight loss} = \frac{\text{Initial weight of seeds} - \text{Final weight of seeds}}{\text{Initial weight of seeds}} \times 100$$

**Comment [CM19]:** It is strongly recommended to give the citation of formal

### Statistical Analysis

The data averaged into respective parameter requisite will be subjected to suitable transformation. After analysis, data will be accommodated in the table as per the needs of objectives for interpretation of results. The standard procedures in agriculture statistics given by **Gomez and Gomez (1976)** were consulted throughout. The interpretation of data will be done by using the critical difference value calculated at 0.05 probability level.

### Results and Discussion

### **Efficacy of different plant extracts and insecticide on Percent adult mortality of *Triboliumcastaneum***

The efficacy of different botanicals on mortality of *Triboliumcastaneum* was tested. Freshly emerged 10 beetles are released in the plastic jars. The mortality of released adults were recorded at 5, 10 and 15 days after treatment.

The data given in the table 1 showed that the percent mortality of released adults was highest in treatments viz., Neem leaf powder (65.55 percentage) followed by Eucalyptus leaf powder (55.55 percentage), Tulsi leaf powder (50 percentage), Garlic powder, Ginger rhizome powder, Turmeric powder recorded 41.11 percentage, 31.11 percentage and 20 percentage mortality of adults respectively. For check treatment 100 percentage mortality of adults occurred where as in untreated control 0 percentage mortality recorded..

similar findings were recorded that neem leaf powder was the most effective treatment, observed the mean adult mortality of (67.22) per cent Whereas, turmeric rhizome powder was the least effective in comparison to control against adult mortality (30.56 %). (Sekaret *et al.*,2021)

**Table 1. Effect of different botanicals and insecticide on adult mortality of *Triboliumcastaneum***

Treatments	Mortality after 5 days	Mortality after 10 days	Mortality after 15 days	Over all mean
Neem leaf powder	46.667 (43.07)	66.667 (54.78)	83.333 (66.14)	65.556 (54.57)
Tulsi leaf powder	33.333 (35.21)	53.333 (46.92)	63.333 (52.77)	50.000 (44.96)
Eucalyptus leaf powder	36.667 (37.22)	56.667 (48.84)	73.333 (59.00)	55.556 (48.33)
Turmeric powder	3.333 (7.24)	23.333 (28.78)	33.333 (35.21)	20.000 (24.88)
Ginger powder	16.667 (23.85)	33.333 (35.21)	43.333 (41.15)	31.111 (33.50)
Garlic powder	23.333 (28.78)	43.333 (41.15)	56.667 (48.84)	41.111 (39.62)
Chlorpyrifos 48EC	100.000 (88.34)	100.000 (88.34)	100.000 (88.34)	100.000 (88.34)

control	0.000 (1.65)	0.000 (1.65)	0.000 (1.65)	0.000 (1.65)
F test	S	S	S	S
SEm±	2.887	2.887	2.887	8.174
CD(0.05)	8.655	8.655	8.655	24.508
CV	15.385	10.619	8.824	31.175

S= Significant;

Note: Figures in the table are mean values and those in parenthesis are angular transformed values

#### Efficacy of different plant extracts and insecticide on Weight loss in percentage of wheat grains

The efficacy of different botanicals on percent weight loss of grains was tested Freshly emerged 10 beetles are released in the plastic jars. The weight loss of grains were recorded at 30, 60 and 90 days after treatment.

The data given in the table 2 showed that the percent weight loss of grains was highest in treatments viz., Turmeric 1.2 percentage followed by Ginger 0.85 percentage, Garlic 0.76 percentage, Tulsi 0.65 percentage, Eucalyptus 0.46 percentage, Neem 0.34 percentage. Where as in control highest weight loss is observed 1.7 percentage and least weight loss is observed in check treatment 0.24 percentage.

Similar findings were recorded that the minimum weight loss recorded in Neem followed by Eucalyptus and Tulsi. (Sekaret *et al.*, 2021).

**Table 2. Effect of different botanicals and insecticide on weight loss of Wheat grains**

Treatments	Weight loss after 30 days	Weight loss after 60 days	Weight loss after 90 days	Over all mean
Neem leaf powder	0.133 (2.06)	0.333 (3.30)	0.567 (4.31)	0.344 (3.21)
Tulsi leaf powder	0.267 (2.94)	0.667 (4.68)	1.033 (5.83)	0.656 (4.46)
Eucalyptus leaf powder	0.20 (2.56)	0.433 (3.76)	0.767 (5.02)	0.467 (3.77)

Comment [CM20]: Table must be cited in the above text

<b>Turmeric powder</b>	0.833 (5.23)	1.233 (6.37)	1.567 (7.19)	1.211 (6.25)
<b>Ginger powder</b>	0.433 (3.76)	0.867 (5.34)	1.267 (6.46)	0.856 (5.17)
<b>Garlic powder</b>	0.467 (3.91)	0.667 (4.68)	1.167 (6.19)	0.767 (4.19)
<b>Chlorpyriphos 48EC</b>	0.083 (1.65)	0.233 (2.76)	0.417 (3.70)	0.244 (2.68)
<b>control</b>	0.933 (5.54)	1.867 (7.85)	2.467 (9.03)	1.756 (7.46)
<b>F test</b>	S	S	S	S
<b>SEm±</b>	0.031	0.031	0.031	0.235
<b>CD(0.05)</b>	0.087	0.095	0.095	0.708
<b>CV</b>	11.950	6.979	4.753	51.964

S= Significant;

Note: Figures in the table are mean values and those in parenthesis are angular transformed values

### Conclusion:

In the research, it was concluded that neem leaf powder was the most effective treatment tailed by eucalyptus and tulsi, whereas, turmeric rhizome powder was the least effective in comparison to control against adult mortality, grain weight loss and owing to *T. castaneum*. Other than this the chemical insecticide, Chloropyriphos used as a check showed a greater result than botanical powders. Even though chemical insecticides have an excellent track record of protecting seeds against storage pests, their negative impact on environment and human health necessitates the development of a new technique for their safe use. These issues can be solved using botanicals such as plant powders they are extremely beneficial for safeguarding seeds from stored product insects in an environmentally friendly manner.

### References

**Comment [CM21]:** In conclusion you must show all your results (values)

1. **Ali, S., Sagheer, M., Hassan, M., Abbas, M., Hafeez, F., Farooq, M., Hussain, D., Saleem, M. and Ghaffar, A. (2014).** Insecticidal activity of turmeric (*Curcuma longa*) and garlic (*Allium sativum*) extracts against red flour beetle, *Triboliumcastaneum*: A safe alternative to insecticides in stored commodities. *Journal of Entomology and Zoology Studies*, **2**(3): 201-205.
2. **Chadda, I.C. (2016).** Fumigation with phosphine-a perspective. *Indian Journal of Entomology*, **78**: 39-44.
3. **Gomez KA, Gomez AA.** Statistical procedures for agricultural research with emphasis on rice. Statistical procedures for agricultural research with emphasis on rice; 1976.
4. **Grünwald, S., Adam, I. V., Gurmai, A. M., Bauer, L., Boll, M. and Wenzel, U. (2013).** The red flour beetle *Triboliumcastaneum* as a model to monitor food safety and functionality. *Yellow Biotechnology InsectBiotechnologie in Drug Discovery and Preclinical Research*, 111-122.
5. **Jagadeesan R, Collins PJ, Daghish GJ, Ebert PR, and Schlipalius DI (2012)** Phosphine Resistance in the Rust Red Flour Beetle, *Triboliumcastaneum* (Coleoptera: Tenebrionidae): Inheritance, Gene Interactions and Fitness Costs. <https://doi.org/10.1371/journal.pone.0031582>
6. **Karunakaran, C., Jayas, D. S. and White, N. D. G. (2004).** Identification of wheat kernels damaged by the red flour beetle using X-ray images. *Biosystems Engineering*, **87**(3), 267-274.
7. **Rehman, H. U., Rasul, A., Farooqi, M. A., Aslam, H. M. U., Majeed, B., Sagheer, M. and Ali, Q. (2020).** Compatibility of some botanicals and the entomopathogenic fungus, *Beauveria bassiana* (Bals.), against the red flour beetle, *Triboliumcastaneum* (Herbst)(Coleoptera: Tenebrionidae). *Egyptian Journal of Biological Pest Control*, **30**(1), 1-7.
8. **Sekar, G.C., Singh, K.H.I., Nagaraju, N., Loganathan, R., Singh, N.O. and Singh, L.N. (2021).** Botanical Management of Rust Red Flour Beetle, *Triboliumcastaneum* (Herbst) on Stored Wheat. *International Journal of Agriculture Environment and Biotechnology* **14**(03): 381-386.
9. **Shewry, P. R., and Hey, S. J. (2015).** The contribution of wheat to human diet and health. *Food and energy security*, **4**(3): 178-202.