

**Cost Effective of Gram pod borer *Helicoverpa armigera* (Hubner) on green gram
at Allahabad**

Abstract

During *kharif* season of 2022, an experiment was conducted to evaluate the cost benefit ratio by using different insecticidal application *viz.*, Chlorantraniliprole 18.5 SC (0.5ml/l), Spinosad 45 SC (0.4ml/l), Nisco sixer plus (2ml/l), *Bacillus thuringiensis* 4% WSP (2gm/l), *Beauveria bassiana* 1.15% WP (5gm/l), Neem oil @2% (5ml/l), ½ dose Chlorantraniliprole + Nisco sixer plus (2.25ml/l) and Control plot against gram pod borer, *Helicoverpa armigera* (Lepidoptera, Noctuidae) on green gram with three replications. Results revealed that highest grain yield was recorded in (T₁) Chlorantraniliprole 18.5 SC (15.6 q/ha) followed by (T₂) Spinosad 45 SC (14.8 q/ha), (T₇) Half dose Chlorantraniliprole + Nisco sixer plus (13.8 q/ha). Insecticidal treatment with (T₁) Chlorantraniliprole 18.5 SC (1:2.87), followed by (T₂) Spinosad 45 SC (1:2.79), (T₇) Half dose Chlorantraniliprole + Nisco sixer plus (1:2.45), Nisco sixer plus (1:2.36), *Beauveria bassiana* 1.15% WP (1:2.30), *Bacillus thuringiensis* 4% WSP (1:2.00) and Neem oil @2% (1:1.83) is found to be least effective but comparatively superior over the control (1:1.51). End of the experiment it reported that Chlorantraniliprole is the best for management of gram pod borer.

KEY WORDS: Biopesticides, Chlorantraniliprole, Cost benefit ratio, green gram, *Helicoverpa armigera*, Insecticides.

INTRODUCTION

Pulses, also known as legumes, they are the edible seeds of leguminous plants cultivated for food. Pulses constitute an excellent supplement of protein in the vegetarian diet of human being and plays a significant role in correcting the widespread malnutrition all over the world. Pulses are known as the “poor man’s meat” because they are rich in nutrition and low in cost **Umbarkar et al. (2010).**

Mung bean (*Vigna radiata*) is a plant species of Fabaceae which is also known as green gram. The green gram is an annual vine with yellow flowers and fuzzy brown pods. There are three subgroups of *Vigna radiata*, including one cultivated (*Vigna radiata* subsp. *radiata*) and two wild ones (*Vigna radiata* subsp. *sublobata* and *Vigna radiata* subsp. *glabra*). It has a height of about 15–125 cm. Mung bean has a well-developed root system. The lateral roots are many and slender, with root nodules grown. Stems are much branched, sometimes twining at the tips. Young stems are purple or green, and mature stems are greyish yellow or brown **Meena et al. (2021).**

India is the world’s largest producer as well as consumer of green gram. It produces about 1.5 to 2.0 million tonnes of mung bean annually from about 3 to 4 million hectares of area with an average productivity of 500 kg per hectare. Green gram output accounts for about 10- 12% of total pulse production in the country. the mung bean production in India was 1.39 million tonnes in which, Maharashtra’s contribution was about 20%, while Rajasthan was highest having 26% of the total production. Mung bean production in the country is largely concentrated in five states viz., Rajasthan, Maharashtra, Andhra Pradesh, Gujarat and Bihar. These five states together contribute for about 70% of total Mung production in the country. **(Anonymus, 2015).**

The major insect pests during different growth stages are thrips, whitefly, leafhopper and stem fly caused appreciable damage. But, worldwide, over 30 species of lepidoptera feed on pods and seeds **Shanower et al. (1999).**

The insect pests are one of the major biotic constraints for reduced yield of green gram. About 17 insect pests which are regarded as key pests are reported to cause significant yield losses in green gram **Cheema et al. (2017).** Pod borer, *Helicoverpa armigera* (Hubner), is a key pest found to cause pod damage up to 27.49 % **Joshi et al. (2019).**

Materials and Methods:

The trail was conducted during *kharif* season 2022 at Sam Higginbottom University of Agriculture Technology And Sciences Prayagraj UP, Central research farm (CRF), Prayagraj, Uttar Pradesh, India, in a randomized block design with eight treatments replicated three times using Malini variety in a plot size of (2m×1m) at a spacing of (30×10cm) with a recommended package of practices excluding plant protection. The treatments used in experiment are *viz.*, Chlorantraniliprole 18.5SC (0.5ml/l), Spinosad 45 SC (0.4ml/l), Nisco sixer plus (2ml/l), *Bacillus thuringiensis* 4% WSP (2gm/l), *Beauveria bassiana* 1.15% WP (5gm/l), Neem oil @2% (5ml/l), ½ dose Chlorantraniliprole + Nisco sixer plus (2.25ml/l) and Control were evaluated against gram pod borer. Each treatment was replicated thrice. All the agronomic practices were followed as per the recommended package of practices. Two sprays were given for all treatments when the crop is at 25 days old except control plot and second spray after 15 days later. The observations were recorded on five randomly selected plants in each replication.

2.1 Yield: (q/ha)

The green gram pods were picked from all the plants per plot and pods were shelled. The average weight of picked pods was used to calculated by the following formula

$$Yield = \frac{Yield\ per\ plot}{Plot\ size} \times 100$$

2.2 Benefit Cost Ratio:

Gross return was calculated by multiplying total yield with the market price of the produce. Cost benefit ratio by following formula

$$B:C\ Ratio = \frac{Gross\ returns}{Total\ Cost\ of\ cultivation}$$

Where,

B: C = Benefit Cost Ratio

Results and Discussion:

The yield among the different treatments were significant. All the treatments were superior over control. The highest increased yield over control was recorded in Chlorantraniliprole 18.5 SC (15.6q/ha) followed by Spinosad 45 SC (14.8 q/ha), ½ dose Chlorantraniliprole + Nisco sixer plus (13.8q/ha), Nisco sixer plus (12.5 q/ha), *Beauveria bassiana* 1.15% WP (11.4 q/ha), *Bacillus thuringiensis* 4% WSP (10.8 q/ha) and Neem oil (10.3 q/ha) is found to be least effective but comparatively superior over the control (7.1 q/ha).

The increased per cent yield over control treatment was different. All treatments were superior over control. The highest increase yield over control was recorded in Chlorantraniliprole 18.5 SC (8.7 q/ha) followed by Spinosad 45 SC (7.7 q/ha), ½ dose Chlorantraniliprole + Nisco sixer plus (6.7 q/ha), Nisco sixer plus (5.4 q/ha), *Beauveria bassiana* 1.15% WP (4.3 q/ha), *Bacillus thuringiensis* 4% WSP (3.7 q/ha) and Neem oil (3.2 q/ha).

When cost benefit ratio was worked out, interesting result was achieved. Among the treatments studied, the best and most economical treatment was Chlorantraniliprole 18.5 SC (1:2.87). Similar finding made by **Hanumanth and kumar (2022)**, followed by Spinosad 45 SC (1:2.79) is found to be the next best treatment which is in line with the findings of **Lakshmikanth and Kumar (2018)**, ½ dose Chlorantraniliprole + Nisco sixer plus (1:2.45), Nisco sixer plus (1:2.36), *Beauveria bassiana* 1.15% WP (1:2.30), *Bacillus thuringiensis* 4% WSP (1:2.00) and Neem oil (1:1.83) is found to be least effective and this finding is supported **Sravani and Kumar (2022)** but comparatively superior over the control (1:1.51).

From the Table 3. It shows that higher yield comes from Chlorantraniliprole insecticides and also more benefit is seen as compared to other treatments used in the experiment. Among all the treatments Chlorantraniliprole is effective.

Table.1. Cost of agronomical practices of cultivation/ha

| S.No | Particular | Requirement | Rate/unit (₹) | Cost (₹) |
|-------------|---------------------------|---------------------|----------------------|-----------------|
| (A) | Land preparation | 2.5 hours | 500₹ /hours | 1250 |
| | Ploughing | 2 hours | 500₹ /hours | 1000 |
| | Harrow | 10 labours | 340₹ /labour | 3400 |
| | Layout of field | 2 labours | 340₹ /labour | 680 |
| (B) | Manures and fertilizer | | | |
| | FYM | 10 tons | 200₹ /ton | 2000 |
| | Urea | 30 Kg | 10₹ /Kg | 300 |
| | SSP | 60 Kg | 24₹ /Kg | 1440 |
| | MOP | 30 Kg | 18₹ /Kg | 480 |
| | Labour | 4 labours | 340₹ /Labour | 1360 |
| (C) | Seed sowing | | | |
| | Seed material | 30 Kg | 160₹ /Kg | 4800 |
| | Sowing and transplanting | 7 labours | 340₹ /Labour | 2380 |
| (D) | Weed management | 8 labours x 2 times | 340₹ /labour | 5440 |
| (E) | Water management | 4 labours x 3 times | 340₹ /labour | 4080 |
| (F) | Harvesting | 8 labours | 340₹ /labour | 2720 |
| (G) | Total cost of cultivation | | | 31330 |

Table.2. Economics of the Treatments

| S.No | Treatments | Use of Chemical (2 times spray) | Cost of Chemical (₹) | Total Cost of Chemical (₹/ha) | Total labour cost (₹) | Total cost of treatment (₹) |
|------|---|---------------------------------|----------------------|-------------------------------|-----------------------|-----------------------------|
| 1 | Chlorantraniliprole 18.5 SC | 500ml/ha | 8800 ₹/lit | 4400 | 680 | 5080 |
| 2 | Spinosad 45 SC | 400ml/ha | 8800 ₹/lit | 3520 | 680 | 4200 |
| 3 | Nisco sixer plus | 2 Litre/ha | 1700 ₹/lit | 3400 | 680 | 4080 |
| 4 | <i>Bacillus thuringiensis</i> 4% WSP | 2kg/ha | 435 ₹/kg | 870 | 680 | 1550 |
| 5 | <i>Beauveria bassiana</i> 1.15% WP (1X10 ⁸ CFU/gm) | 4kg/ha | 550₹/kg | 1100 | 680 | 1780 |
| 6 | Neem oil 2% | 5litre/ha | 320 ₹/lit | 1600 | 680 | 3030 |
| 7 | ½ dose Chlorantraniliprole + Nisco sixer plus | 250ml+ 2 litre/ha | 8800 + 1700 ₹/lit | 5600 | 680 | 5668 |
| 8 | Control | – | – | – | – | – |

Table.3. Effect of treatments on green gram

| S.NO | TREATMENTS | Yield Q/ha | Increase yield over control Q/ha |
|----------------------|---|-------------------|---|
| T₁ | Chlorantraniliprole 18.5SC | 15.6 | 8.7 |
| T₂ | Spinosad 45SC | 14.8 | 7.7 |
| T₃ | Nisco sixer plus | 12.5 | 5.4 |
| T₄ | <i>Bacillus thuringiensis</i> 4% WSP | 10.8 | 3.7 |
| T₅ | <i>Beauveria bassiana</i> 1.15% WP (1X10 ⁸ CFU/gm) | 11.4 | 4.3 |
| T₆ | Neem oil 2% | 10.3 | 3.2 |
| T₇ | ½ dose Chlorantraniliprole + Nisco sixer plus | 13.8 | 6.7 |
| T₈ | Control | 7.1 | — |




Fig.1. Assessment of yield on the efficacy of Chlorantraniliprole and biopesticides on gram pod borer

Table.4. Economics and cost benefit ratio of the cultivation

| Treatment symbol | Treatment | Yield (q/ha) | Cost of yield q/₹ | Total cost of yield in ₹ | Common cost of cultivation (₹) | Total treatment cost (₹) | B:C Ratio |
|-------------------------|---|---------------------|--------------------------|---------------------------------|---------------------------------------|---------------------------------|------------------|
| T₁ | Chlorantraniliprole 18.5 SC | 15.6 | 6700 | 104520 | 31330 | 5080 | 1:2.87 |
| T₂ | Spinosad 45 SC | 14.8 | 6700 | 99160 | 31330 | 4200 | 1:2.79 |
| T₃ | Nisco sixer plus | 12.5 | 6700 | 83750 | 31330 | 4080 | 1:2.36 |
| T₄ | <i>Bacillus thuringiensis</i> 4% WSP | 10.8 | 6700 | 72360 | 31330 | 1550 | 1:2.00 |
| T₅ | <i>Beauveria bassiana</i> 1.15% WP (1X10 ⁸ CFU/gm) | 11.4 | 6700 | 76380 | 31330 | 1780 | 1:2.30 |
| T₆ | Neem oil 2% | 10.3 | 6700 | 69010 | 31330 | 3030 | 1:1.83 |
| T₇ | ½ dose Chlorantraniliprole + Nisco sixer plus | 13.8 | 6700 | 92460 | 31330 | 5668 | 1:2.45 |
| T₈ | Control | 7.1 | 6700 | 47570 | 31330 | 0 | 1:1.51 |




Fig.2. Cost benefit ratio of treatments

4. Conclusion:

Results revealed that the maximum yield and cost benefit ratio is recorded at Chlorantraniliprole 18.5 SC, followed by Spinosad 45 SC, half dose of Chlorantraniliprole + Nisco sixer plus can be suitably incorporated in pest management schedule against gram pod borer as an effective tool under chemical control.

References:

- Cheema, H.K., Pratap, A. and Sujayanand, G. K. (2017).** Breeding for Insect Resistance in Mung Bean and Urd Bean. *In: Breeding Insect Resistant Crops for Sustainable Agriculture Springer*, Singapore, (pp. 353-385).
- Hanumant, P. A. and Kumar, A. (2022).** Field evaluation of chemicals and bio-pesticides against chickpea pod borer [*Helicoverpa armigera* (Hubner)]. *The Pharma Innovation Journal*, **11**(7): 3405-3410.
- Joshi, N., Shera, P., Sangha, K. and Sharma, S. (2019).** Bioformulations for management of pod borer, *Helicoverpa armigera* (Hübner) in Mungbean (*Vigna radiata* L.). *Journal of Biological Control*. **33**(1): 76-79.
- Lakshmikanth, R. and Kumar, A. (2018).** Comparative efficacy of selected chemicals and Biopesticides against gram pod borer [*Helicoverpa armigera* (Hubner)] (Lepidoptera: Noctuidae) on cowpea [*Vigna unguiculata* (L.) Walp.]. *Journal of pharmacognosy and phytochemistry*, **7**(3): 3307-3309.
- Meena, V. P., Khinchi, S. K., Kumawat, K. C. and Choudhary, S. (2022).** Seasonal incidence of gram pod borer, *Helicoverpa armigera* (Hubner) and spotted pod borer, *Maruca testulalis* (Geyer) on green gram in relation to weather parameters, *Journal of Experimental Agriculture International*, **19**(1), 1-8.
- Shanower, T. G., Romeis, J. M. E. M. and Minja, E. M. (1999).** Insect pests of pigeonpea and their management. *Annual review of entomology*, **44**(1), 77-96.
- Sravani, M. and Kumar, A. (2022).** Field efficacy and economics of selected insecticides against pod borer [*Helicoverpa armigera* (Hubner)] on green gram [*Vigna radiata* (L.) Wilczek]. *The Pharma Innovation Journal*, **11**(7): 3920-3923.
- Umbarkar, P. S., Parsana, G. J. and Jethva, D. M. (2010).** Seasonal incidence of gram pod borer, *Helicoverpa armigera* (Hubner) on greengram. *Legume Research-An International Journal*, **33**(2): 148-149.