

# Morphological characterization of Recombinant Inbred lines of Soybean [*Glycine max* (L.) Merrill]

## Abstract

Morphological characterization facilitates the seed production agencies to identify the varieties on phenotypic basis by their distinguished characters. The presence of sufficient variability for various agro-morphological traits, as well as the improvement of these traits through selection and introduction of new alleles, lines, and accessions into breeding activity from both exotic and indigenous collections, aids in the identification of potential donors. In the present study, 100 genotypes including 96 RIL (recombinant inbred lines), two parental and two national checks were planted at Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur. Fourteen phenological traits viz. growth type, growth habit, flower colour, leaf size of lateral leaflet, leaf shape of lateral leaflet, leaf intensity of green colour, pod hairiness, pod colour of hair, seed size, seed shape, seed coat colour of testa, seed coat lustre, hilum colour and hilum funicle were observed at different growth stages of plants. Among these traits, Semi-determinant growth type (97 %), Semi-erect growth habit (98 %), white flower colour (99 %), Pod colour of hair (100%) and hilum funicle same as testa (97%) were most dominating. Whereas Seed coat colour of testa [yellow (74 %), yellow-green (24 %) and Green (2 %)], Hilum colour [black (48 %) and brown (43 %) and imperfect black (9%)] and Seed shape [spherical (48 %), spherical flattened (33%), flattened (16 %) and elongated (3%)] were varied among genotypes.

**Keywords:** Heterogeneity, exotic collection and determinant growth

## Introduction

Soybean [*Glycine max* (L) Merrill] currently ranks first in the world and third in India among the major oilseed crops after groundnuts and rapeseed-mustard with an area of 12.5 million hectares and a production of 13.2 million tonnes and yield of 1.06 t/ha (Director's Report, IISR, 2021-22). Soybean is belonging to the family Leguminosae syn. Fabaceae, subfamily Faboidae, and tribe Phaseoleae. *Glycine* (Wild.) is further subdivided into two subgenera: *Glycine* and *Soja*. There are 26 wild perennial species in the *Glycine* subgenus (Hermann 1962; Newell and Hymovitz 1980). The seed of soybean consist very high range of protein (36.1 – 41.2 %) with considerable amount of edible oil (16.8 – 20.2 %) and several nutritional properties (Amrate *et al.* 2020; Uikey *et al.* 2022). The amount of oil, protein, or other favourable substances was shown to be connected to the size and shape of the seeds (Cober and Voldeng 2008; Barela *et al.* 2022(c); Yang *et al.* 2019; Uikey *et al.* 2022).

Crop genotype exploitation is mostly dependent on the presence of **genetic variability** within or between them and the crop genetic resources have been categorised in large part by qualitative traits (Verma *et al.* 2021). On the other hand, morphological characterization facilitates the seed production agencies to identify the varieties on phenotypic basis by their distinguished characters. To a large part, the commercial value of soybean seeds is determined by their visual quality, which comprises seed size, shape, colour and texture. In addition to the direct effect of seed size on yield, many studies have found a link between the external appearance of soybean seeds and their inside quality. Morphological characterisation is an important stage in the description and classification of accessions or genotypes because a breeding programme is largely dependent on the amount of genetic variability.

Multiple superior genotypes should be managed simultaneously in the seed production and varietal development programme. Numerous opportunities exist for genetic duplication and admixture. The Protection of Plant Varieties and Farmers Rights Act (PPV&FRA), 2001, intends to safeguard plant varieties based on distinctiveness, uniformity, and stability (DUS). In soybean and other crops, varieties and genotypes are morphologically recognised using descriptors based on distinctiveness, uniformity, and stability (DUS) (Ramteke and Muralitharan. 2012; Dhaliwal *et al.*, 2020, Banerjee *et al.*,2022). It shows to be a solid method to define genotypes since the qualitative traits demonstrate stability over a range of environment and generations. Looking to all these aspects, an investigation was carried out to characterise the recently developed RILs of soybean at IISR, Indore on the basis of

phenotype and to draw important selection indices by estimating genetic variability, heritability and genetic advance with respect to the different yield attributing traits.

### **Materials and methods**

This experiment was conducted at breeding field of J.N.K.V.V., Jabalpur, during Kharif 2021. Jabalpur is most prominent place of soybean cultivation in India and all the varieties or genotypes express their characters fully. Two blocks were used in the experiment's randomized complete block design (RCBD), which helped to minimize experimental error and field variation. Each block was planted with two replications of 96 genotype RIL lines, each of which had two parental and two national checks from the IISR in Indore and the JNKVV in Jabalpur. The List of RILs, along with their pedigree showed in Table 1. Individual plant observations as well as net plot observations for morphological features were kept track of fourteen phenological trait viz. **growth type: determinate and semi-determinate**, growth type, growth habit, flower colour, leaf size of lateral leaflet, leaf shape of lateral leaflet, leaf intensity of green colour, pod hairiness, pod: colour of hair, seed size, seed shape, seed coat colour of testa, seed coat lustre, hilum colour and hilum funicle were used to record the observations of the lines proposed for this investigation.

### **Results and discussion**

There were 98 genotypes that were show semi-erect, and 2 genotypes erect type growth habit. Out of 100 genotypes only one genotype was violet, while the ninety-nine genotypes were white. Small leaves were found in 4 genotypes, medium leaves in 8 genotypes, and large leaves in 88 genotypes.


**The RILs were classified into two groups based on the shape of the lateral leaflet: round ovate, pointed ovate. 23 genotypes were round ovate and 77 genotypes were pointed ovate leaves. 97 genotype expressed green colour and 3 genotypes expressed dark green colour.** RILs were classified depending on the presence or absence of hair on the pod: pubescent (hairs present) and glabrous (hairs absent). Ninety-nine genotypes had pubescence (either grey or tawny), while the one was glabrous. At the advanced pod filling stage, the pod colour was observed to be tawny brown. One hundred genotypes indicated tawny pubescence. Genotypes are divided into three groups based on seed size: small (less than 10

g), medium (10-13 g), and bold/large (more than 13 g). Forty-five genotypes had small seeds, forty-four had medium seeds, and the remaining eleven had bold/large seeds.


Out of 100 genotype, 48 genotypes had spherical seeds, 33 had spherical flattened seeds, 3 had elongated seeds, and the remaining 16 genotypes had flattened seeds. The colour of testa excluding hilum were shown as yellow, yellow green, green. 74 genotypes had yellow seed coats, 2 genotypes had yellow-green seed coats, and the remaining 2 genotypes had green seed coats. Seed coat lustre of ten genotypes was shining, 81 genotypes were intermediate, and the other nine genotypes had dull seed coat lustre. Barela *et al.*, 2022(b), Soni *et al.*, (2022) Banerjee *et al.*, (2022), and Rahangdale *et al* (2023) revealed that there was a lot of variation in the soybean genotypes for DUS characters and qualitative traits provide stability over changing environments and generations.

Hilum colour were classified into four categories: brown, imperfect black, and black. 43 genotypes had brown hilums, 9 had imperfect black hilums, and 48 had black hilums. Hilum funicle colour was found to be either the same as testa or different from testa. Ninety-seven genotypes had hilum funicle colour that was the same as the testa, while the remaining three genotypes had hilum funicle colour that was different from the testa.

To establish phenotypic distinctiveness among soybean genotypes, morphological features must be used to classify them, making genotype identification easier in the future seed production programme. Since the DUS recommendations provided under the Protection of Plant Varieties and Farmers' Rights Act 2001 (PPVFRA) have become mandatory for the provision of protection to varieties and germplasm, etc. As a result, morphological features must be used to characterise the RILs. These distinct characteristics can assist breeders and seed producers in identifying genotypes on phenotypic basis. For morphological features, Barela *et al.* 2022(a), Soni *et al.*, (2022), Painkra *et al.*, (2019), Uikey (2022) and Mehra (2019) used a similar characterisation study.



**Figure 1:** Graphical representation of particular frequency of different traits



**Figure 1:** Different destructive features of soybean (a) Pubescent (b) Glabrous, Leaf shape of lateral leaflet (c) Pointed leaf (d) Ovate leaf, Hilum colour (e) Black Hilum (f) Brown hilum (g) Imperfect black hilum (h) Seed size.

UNDER

## CONCLUSION

To meet the growing global demand for food security, genetic resources offer the basis for breeding selection and enhancement. The material in the soybean development programme is used more effectively when it is systematically characterized. The wide variety of seed hilum colors present in the RILs accounts for the high market price and profitability for farmers. Genotypes with this morphology can be chosen as a donor in the crossing operation after the stability and heritability of the traits have been determined.

## References

- Amrate PK, Shrivastava MK, Pancheshwar, Dinesh K, Stuti S. Charcoal rot and yellow mosaic virus disease of soybean under hotspot condition: symptoms, incidence and resistance characterization. *International Journal of Bio-Resource and Stress Management*. 2020; 11(3): 268–273.
- Barela A, Jain S, Tiwari S, Rahangdale S, Singh P. Impact of gamma radiations on seed germination and morphological characteristics of pea (*Pisum sativum* L.) *The Pharma Innovation Journal*. 2022(a);11(8): 464-467.
- Barela A, Shrivastava MK, Mohare S, Rahangdale S, Jawarkar S, Amrate PK and Singh Y., Morphological Characterization and Recognition of New Traits of Soybean [*Glycine max* (L.) Merrill] *International Journal Environment and Climate Change* 2022(b);12(12):1497-1504.
- Barela, A., Jain, S., Khandalkar, V.S., Singh, P., Rahangdale, S., Behra, K. and Singh, J. Ameliorations of Gamma Radiations on Chlorophyll Spectrum and Morphological Characteristics of Peas. *International Journal of Environment and Climate Change*. 2022(c); 12(12), pp.1724-1731.
- Cober ER, Voldeng HD. Mass selection for small seed size in natto soybean populations and the resulting effect on seed yield. *Crop Sci* .2000; 48,1337–1340.
- Dhaliwal I, Khosla G, Singh TP, Gill GS, Kaushik P. DUS characterization of some released varieties and advanced breeding lines of soybean (*Glycine max* L.) under Punjab

Agroclimatic Conditions. International Journal of Current Microbiology and Applied Sciences. 2020; 92124-35.

Director report ( AICRP Soybean), Annual Progress Report. ICAR-Indian Institute of Soybean Research, Khandwa Road, Indore 452001 (M.P.).2022.

Banerjee J, Shrivastava MK, Amrate PK, Singh Y, Upadhya A, Soni M. Genetic variability and association of yield contributing traits in advanced breeding lines of soybean. Electronic Journal of Plant Breeding. 2022;13(2).

Soni M, Shrivastava MK, Singh Y, Sharma S, Amrate PK, and Banerjee J, Evaluation the Genetic Variability and Association Analysis of Soybean (*Glycine max* (L.) Merrill for some Physiological Parameters. International Journal of Plant & Soil Science. 2022; 34(10): 17-26.

Uikey S, Sharma S, Amrate Pawan K, Shrivastava MK. Identification of rich oilprotein and disease resistance genotypes in soybean [*Glycine max* (L.) Merrill]. International Journal of Bio-resource and Stress Management. 2022;13(5):497-506.

Hermann FJ. A revision of the genus *Glycine* and its immediate allies. US Department of Agriculture, 1962; pp: 1268.

Mehra S, Shrivastava MK, Amrate PK, Yadav RB. Studies on variability, correlation coefficient and path analysis for yield associated traits in soybean [*Glycine max* (L.) Merrill]. Journal of Oil seeds Research. 2020;37(1):56–59.2.

Newell, C.A. and Hymowitz. A Taxonomic Revision in the Genus *Glycine* Subgenus *Glycine* (Leguminosae). Springer 1980; 32:63-69

Painkra P, Shrivastava R and Nag SK. Argo-morphological characterization of soybean germplasm. Journal of Oilseeds Research 36 (2),85,2019.

PPV & FRA. Guidelines for the Conduct of Test for Distinctness, Uniformity and Stability on Soybean (*Glycine max* (L.) Merrill); 2018.

Rahangdale, S., Lakhani, J.P., Singh, S.K. and Barela, A., Phenotyping and reveal novel traits in mung bean (*Vigna radiata* L. Wilczek) genotypes. *South African Journal of Botany*. 2023. 156, pp.65-71.

Ramteke R, Murlidharan P. Characterization of soybean (*Glycine max*) varieties as per DUS guidelines. Indian Journal of Agricultural Sciences. 2012;82(7):572–577.

Verma V., Shrivastava MK, Mehra S, Amrate PK. and Yadav RB. Estimation of Genetic Parameters for Yield Associated Traits and Principal Component in Advance Breeding Lines of Soybean. (*Glycine max*. (L.) Merrill). International Journal Current Microbiology and Applied Science.2021;10 (01): 2704-2710.

Yang H, Wang W, He Q *et al.* 2019. Identifying a wild allele conferring small seed size, high protein content and low oil content using chromosome segment substitution lines in soybean. *Theor Appl Genet* 132:2793–2807.

**Table.1** List of Genotypes and their parentage.

S.No.	Name of RILs	Pedigree	S.No.	Name of RILs	Pedigree
1	<b>RIL-107-1</b>	JS 97-52 × NRC 37	51	<b>RIL-107-95</b>	JS 97-52 × NRC 37
2	<b>RIL-107-3</b>	JS 97-52 × NRC 37	52	<b>RIL-107-96</b>	JS 97-52 × NRC 37
3	<b>RIL-107-5</b>	JS 97-52 × NRC 37	53	<b>RIL-107-97</b>	JS 97-52 × NRC 37
4	<b>RIL-107-6</b>	JS 97-52 × NRC 37	54	<b>RIL-107-98</b>	JS 97-52 × NRC 37
5	<b>RIL-107-7</b>	JS 97-52 × NRC 37	55	<b>RIL-107-100</b>	JS 97-52 × NRC 37
6	<b>RIL-107-8</b>	JS 97-52 × NRC 37	56	<b>RIL-107-101</b>	JS 97-52 × NRC 37
7	<b>RIL-107-9</b>	JS 97-52 × NRC 37	57	<b>RIL-107-102</b>	JS 97-52 × NRC 37
8	<b>RIL-107-11</b>	JS 97-52 × NRC 37	58	<b>RIL-107-103</b>	JS 97-52 × NRC 37
9	<b>RIL-107-12</b>	JS 97-52 × NRC 37	59	<b>RIL-107-105</b>	JS 97-52 × NRC 37
10	<b>RIL-107-18</b>	JS 97-52 × NRC 37	60	<b>RIL-107-106</b>	JS 97-52 × NRC 37
11	<b>RIL-107-19</b>	JS 97-52 × NRC 37	61	<b>RIL-107-109</b>	JS 97-52 × NRC 37
12	<b>RIL-107-21</b>	JS 97-52 × NRC 37	62	<b>RIL-107-110</b>	JS 97-52 × NRC 37
13	<b>RIL-107-23</b>	JS 97-52 × NRC 37	63	<b>RIL-107-111</b>	JS 97-52 × NRC 37
14	<b>RIL-107-24</b>	JS 97-52 × NRC 37	64	<b>RIL-107-114</b>	JS 97-52 × NRC 37
15	<b>RIL-107-27</b>	JS 97-52 × NRC 37	65	<b>RIL-107-115</b>	JS 97-52 × NRC 37
16	<b>RIL-107-28</b>	JS 97-52 × NRC 37	66	<b>RIL-107-116</b>	JS 97-52 × NRC 37
17	<b>RIL-107-29</b>	JS 97-52 × NRC 37	67	<b>RIL-107-118</b>	JS 97-52 × NRC 37
18	<b>RIL-107-31</b>	JS 97-52 × NRC 37	68	<b>RIL-107-121</b>	JS 97-52 × NRC 37
19	<b>RIL-107-32</b>	JS 97-52 × NRC 37	69	<b>RIL-107-123</b>	JS 97-52 × NRC 37
20	<b>RIL-107-35</b>	JS 97-52 × NRC 37	70	<b>RIL-107-124</b>	JS 97-52 × NRC 37
21	<b>RIL-107-36</b>	JS 97-52 × NRC 37	71	<b>RIL-107-125</b>	JS 97-52 × NRC 37

22	<b>RIL-107-41</b>	JS 97-52 × NRC 37	<b>72</b>	<b>RIL-107-126</b>	JS 97-52 × NRC 37
23	<b>RIL-107-46</b>	JS 97-52 × NRC 37	<b>73</b>	<b>RIL-107-127</b>	JS 97-52 × NRC 37
24	<b>RIL-107-48</b>	JS 97-52 × NRC 37	<b>74</b>	<b>RIL-107-131</b>	JS 97-52 × NRC 37
25	<b>RIL-107-49</b>	JS 97-52 × NRC 37	<b>75</b>	<b>RIL-107-132</b>	JS 97-52 × NRC 37
26	<b>RIL-107-51</b>	JS 97-52 × NRC 37	<b>76</b>	<b>RIL-107-133</b>	JS 97-52 × NRC 37
27	<b>RIL-107-52</b>	JS 97-52 × NRC 37	<b>77</b>	<b>RIL-107-134</b>	JS 97-52 × NRC 37
28	<b>RIL-107-55</b>	JS 97-52 × NRC 37	<b>78</b>	<b>RIL-107-135</b>	JS 97-52 × NRC 37
29	<b>RIL-107-56</b>	JS 97-52 × NRC 37	<b>79</b>	<b>RIL-107-136</b>	JS 97-52 × NRC 37
30	<b>RIL-107-57</b>	JS 97-52 × NRC 37	<b>80</b>	<b>RIL-107-137</b>	JS 97-52 × NRC 37
31	<b>RIL-107-58</b>	JS 97-52 × NRC 37	<b>81</b>	<b>RIL-107-138</b>	JS 97-52 × NRC 37
32	<b>RIL-107-59</b>	JS 97-52 × NRC 37	<b>82</b>	<b>RIL-107-140</b>	JS 97-52 × NRC 37
33	<b>RIL-107-60</b>	JS 97-52 × NRC 37	<b>83</b>	<b>RIL-107-141</b>	JS 97-52 × NRC 37
34	<b>RIL-107-62</b>	JS 97-52 × NRC 37	<b>84</b>	<b>RIL-107-143</b>	JS 97-52 × NRC 37
35	<b>RIL-107-64</b>	JS 97-52 × NRC 37	<b>85</b>	<b>RIL-107-151</b>	JS 97-52 × NRC 37
36	<b>RIL-107-65</b>	JS 97-52 × NRC 37	<b>86</b>	<b>RIL-107-154</b>	JS 97-52 × NRC 37
37	<b>RIL-107-67</b>	JS 97-52 × NRC 37	<b>87</b>	<b>RIL-107-157</b>	JS 97-52 × NRC 37
38	<b>RIL-107-68</b>	JS 97-52 × NRC 37	<b>88</b>	<b>RIL-107-159</b>	JS 97-52 × NRC 37
39	<b>RIL-107-70</b>	JS 97-52 × NRC 37	<b>89</b>	<b>RIL-107-161</b>	JS 97-52 × NRC 37
40	<b>RIL-107-71</b>	JS 97-52 × NRC 37	<b>90</b>	<b>RIL-107-168</b>	JS 97-52 × NRC 37
41	<b>RIL-107-73</b>	JS 97-52 × NRC 37	<b>91</b>	<b>RIL-107-172</b>	JS 97-52 × NRC 37
42	<b>RIL-107-76</b>	JS 97-52 × NRC 37	<b>92</b>	<b>RIL-107-173</b>	JS 97-52 × NRC 37
43	<b>RIL-107-79</b>	JS 97-52 × NRC 37	<b>93</b>	<b>RIL-107-174</b>	JS 97-52 × NRC 37
44	<b>RIL-107-80</b>	JS 97-52 × NRC 37	<b>94</b>	<b>RIL-107-177</b>	JS 97-52 × NRC 37
45	<b>RIL-107-83</b>	JS 97-52 × NRC 37	<b>95</b>	<b>RIL-107-178</b>	JS 97-52 × NRC 37
46	<b>RIL-107-84</b>	JS 97-52 × NRC 37	<b>96</b>	<b>RIL-107-181</b>	JS 97-52 × NRC 37
47	<b>RIL-107-85</b>	JS 97-52 × NRC 37	<b>97</b>	<b>JS-97-52</b>	<b>PK 327 X L129</b>
48	<b>RIL-107-88</b>	JS 97-52 × NRC 37	<b>98</b>	<b>NRC 37</b>	<b>Gaurav X Punjab 1</b>

<b>49</b>	<b>RIL-107-91</b>	JS 97-52 × NRC 37	<b>99</b>	<b>JS 335</b>	<b>JS 78-77 X JS 71-05</b>
<b>50</b>	<b>RIL-107-92</b>	JS 97-52 × NRC 37	<b>100</b>	<b>JS 2029</b>	<b>JS 97-52 X JS 95-56</b>

UNDER PEER REVIEW

**Table 2.** Morphological characterization of Soybean RILs

<b>Trait</b>	<b>Classes</b>	<b>Frequency</b>	<b>Note</b>
<b>Growth type</b>	Determinant	3	1
	Semi- determinant	97	2
<b>Growth habit</b>	Erect	2	1
	Semi erect	98	3
<b>Flower colour</b>	White	99	1
	Violet	1	2
<b>Leaf size of lateral leaflet</b>	Small	4	3
	Medium	88	5
	Large	8	7
<b>Leaf shape of lateral leaflet</b>	Pointed ovate	77	3
	Round ovate	23	4
<b>Leaf intensity of green colour</b>	Dark green	3	7
	Green	97	5
<b>Pod hairiness</b>	Pubescent	99	1
	Glabrous	1	0
<b>Pod: colour of hair</b>	Tawny brown	100	2
<b>Seed size</b>	Medium	44	5
	Small	45	3
	Large	11	7
<b>Seed shape</b>	Spherical	48	1
	Spherical-flattened	33	2
	Flattened	16	4
	Elongated	3	3
<b>Seed coat colour of testa</b>	Yellow	74	1
	Green	2	3
	yellow-green	24	2
<b>Seed coat lustre</b>	Intermediate	81	5
	Shiny	10	3
	Dull	9	7

<b>Hilum colour</b>	Black	48	6
	Brown	43	3
	Imperfect Black	9	5
<b>Hilum funicle</b>	Same as testa	97	1
	Different as testa	3	2

UNDER PEER REVIEW