

Original Research Article

Compatibility of fungicides with potent *Trichoderma* isolates

Abstract

The sensitivity of biological control agents were tested against fungicides that are used to treat plant diseases. Application of fungicides should not interfere with biological control agents ability to effectively manage disease. *Trichoderma* is an aggressive coloniser of soil and plant roots that can flourish in a variety of environmental circumstances. It also functions as a natural bioagent to shield plants against infection by soil-borne fungal infections. Evaluation of the various fungicides against five potent *Trichoderma* isolates *in vitro* was done in order to determine the compatibility of bioagents with the fungicides was performed at the Department of Plant Pathology, College of Agriculture, JNKVV, Jabalpur. *Trichoderma* was compatible with the following contact, systemic, and combi fungicides: Imbrex (Fluxapyroxad EC), Curzate M8 (Cymoxanil 8%+ Mancozeb 64% WP), Vitavax (Carboxin 37.5% + Thiram 37.5% DS) and Seedkot (Thiram 75% WS), while it proved just moderately compatible with Amistar top (Azoxystrobin 18.2% + Difenconazole 11.4% SC) and Headline (Pyraclostrobin 20% WG). It showed least compatibility with Nativo (Tebuconazole 50%+ Trifloxystrobin 25% WG), Taqat (Captan 70%+ Hexaconazole 5% WP), Folicur (Tebuconazole 25.9% EC), and Saaf (Carbendazim 12% + Mancozeb 63% WP)

Keywords: Compatibility, Fungicides, *Trichoderma*, ppm

Introduction

Plant diseases can be managed using a variety of biological control agents (BCAs). These consist of bacteria, fungi, and actinomycetes. The most significant BCAs are found in the genus *Streptomyces*, *Bacillus* species, *Pseudomonas* species, and *Trichoderma* species. A promising alternative to decreasing modern agriculture's heavy reliance on expensive chemical fungicides, which not only pollute the environment but also encourage the emergence of resistant strains, is biological control of plant infections [Harman *et al.* 2004].

Mycoparasitism, antibiosis, competition for nutrients or space, increased root and plant development, induced resistance, solubilization and sequestration of inorganic nutrients, and inactivation of the pathogens' enzymes are just a few of the most current mechanisms [Lewis and Lumsden, 2001]. In addition to their capacity for biocontrol, BCAs also exhibit traits like competence in the rhizosphere, resistance to fungicides, saprophytic competitiveness, tolerance of extremes in temperature, adaptability to various environmental conditions, good searching abilities, host specificity, high reproduction rates, short life cycles, adaptability, and the capacity to maintain

themselves while reducing host populations [Okigbo and Ikediugwu, 2000] have demonstrated that *Trichoderma viride* displaced the yam tuber's naturally occurring microflora.

Achieving sustainable global food security will be a challenging task with the growing human population and shifting global food consumption patterns brought on by climate change [Kumar *et al.* 2021]. Therefore, management of plant diseases should focus on eco-friendly measures and use of bioagents for plant disease management will certainly serve as a crucial strategy to achieve it. However, combined use of bioagent along with need-based fungicide may result in better control of plant diseases. Therefore, compatibility of different fungicide with bioagent shall lead to identification of safe fungicide to be used in combination with potential bio-agent like *Trichoderma* spp.. IDM strategies that combine fungicides and suitable bio agents shield seeds and seedlings from soil- and seed-borne pathogens [Dubey and Patil, 2001]. It may be more effective to treat disease and manage soil-borne diseases if appropriate bio agents are used with fungicides [Papavizas and Lewis, 1981]. Similar disease control would be accomplished by combining BCAs and fungicides as opposed to using more fungicides alone [Monte, 2001]. By combining antagonists with synthetic compounds, the potential for resistance development is avoided, and the need for fungicide application is decreased. Therefore, it is suggested to determine whether prospective bio agents are compatible with widely used fungicides for the environmentally friendly approach to treat tea diseases. Understanding the effects of fungicides on the pathogen and the antagonists would help in the selection of fungicides and fungicide resistant antagonists through compatibility studies *in vitro*. Fungicides should have an inhibitory effect on the pathogen but shouldn't have a detrimental effect on the antagonists. Additionally, this approach might show even better control of resistant fungal pathogen strains and aid commercial producers in using less fungicides, reducing the amount of chemical residue in the marketed goods. Combining BCA sprays with small doses of fungicides afterward may help the formulations' antagonists and relative cost [Houdam and Dutta, 2014].

Trichoderma spp. is a globally identified successful bioagent which not only control plant diseases [Kumar *et al.* 2013a; Kumar *et al.* 2014; Jain *et al.* 2017; Kharte *et al.* 2022] but can also be used as biofertilizer (Srivastava *et al.* 2009) and in production of several secondary metabolites [Kumar *et al.* 2009]. They are also helpful in plant growth promotion [Kumar and Sahu, 2014; Kumar *et al.* 2019] and bioremediation [Kumar *et al.* 2015]. Further, their use as a native isolate have proven better potential in the local area for successful bio-control agent after proper identification and characterization [Kumar *et al.* 2013b; Kumar and Sahu, 2015; Kumar *et al.* 2016]. On a variety of crops, *Trichoderma* species have been shown to inhibit root infection by soil-borne diseases such as *Macrophomina phaseolina*, *Rhizoctonia solani*, *Fusarium* species, and *Pythium* species [Benítez *et al.* 2004]. *Trichoderma* species can promote growth, which may or may not be essential for biological control [Dubey *et al.* 2007]. Effective suppression of root-infecting fungi and root-knot nematodes has been demonstrated by *Trichoderma harzianum* [Spiegel and Chet, 1998; Sun and Liu, 2006].

Rhizome rot was inhibited by *Trichoderma harzianum*, which was isolated from the soil [Ram *et al.* 1999]. It also promoted plant growth and yield. According to reports, numerous *Trichoderma* species exhibit varying degrees of innate and/or induced resistance to various fungicides. In the context of this, research was done to see if it was possible to combine *Trichoderma* species with fungicides in a lab conditions. The long-term objective is to create an effective IDM package for controlling soil-borne plant diseases and arresting the spread of pathogens resistant to fungicides. A crucial component of integrated disease management is including chemical-resistant *Trichoderma* species. Utilising tolerant species that keep infections under sufficient pressure to limit their growth can help to prevent disease to a greater extent. Tr-1 (*Trichoderma sp.*), Tr-5 (*T. harzianum*), Tr-7 (*T. yunnanense*), Tr-13 (*T. asperellum*) and Tr-15 (*T. asperellum*) were used as test organisms for the *in vitro* compatibility of different fungicides with the BCA.

Material and Methods

Isolation of *Trichoderma spp.* from rhizosphere soil

The top two centimeters of surface soil were removed in order to collect the rhizosphere soil from the several agro climatic zones of Madhya Pradesh, India, at a depth ranging from 15-20 cm. One g of soil was removed from the composite soil sample and mixed with 9 ml of sterilized distilled water to create a working sample. The serial dilution method was used to separate *Trichoderma spp.* from this sample. One ml suspension was added to each of the three Petri plates containing the potato dextrose agar media (PDA) (Elad *et al.*, 1980) from the 10^{-3} and 10^{-4} dilutions. The plates were gently whirled, and the mixture was then incubated at room temperature. After *Trichoderma* had grown, the most probable colonies were plucked from the culture and examined under a microscope. To obtain pure cultures, the resulting *Trichoderma* isolates were subcultured using single spore isolation. These pure cultures were then transferred to potato dextrose agar slants and kept in the refrigerator at 4°C for future research. All 16 *Trichoderma* isolates were evaluated for their ability to control the important soil-borne plant pathogens such as *Fusarium oxysporum* f. sp. *ciceri*, *Sclerotium rolfsii*, *Sclerotinia sclerotiorum*, *Rhizoctonia solani*, and *Rhizoctonia bataticola*. Five of the best performed *Trichoderma* isolates from this screening were chosen to assess their fungicide compatibility. Through ITS sequencing, the identity of these five chosen *Trichoderma* cultures as Tr-1 (*Trichoderma sp.*-OR150423), Tr-5 (*T. harzianum*- OR150424), Tr-7 (*T. yunnanense*- OR150425), Tr-13 (*T. asperellum*-OR150426) and Tr-15 (*T. asperellum*- OR150427) was further validated.

Table 1: List of fungicides used for testing the compatibility with different *Trichoderma spp.*

Fungicides	Composition	Manufacturing company	Category of fungicide

Amistar Top	(Azoxystrobin 18.2% + Difenoconazole 11.4% SC)	Syngenta	Systemic
Saaf	Carbendazim 12% + Mancozeb 63% WP	UPL India	Systemic and contact
Seedkot	Thiram 75% WS	National Pesticides & Chemicals	Non-systemic
Taqat	Captan 70%+ Hexaconazole 5% WP	ADS Agrotech Pvt. Ltd.	Contact
Curzate M8	Cymoxanil 8%+ Mancozeb 64% WP	Corteva Agri Sciences Pvt. Ltd.	Preventive and locally systemic
Nativo	Tebuconazole 50%+ Trifloxystrobin 25% WG	Bayer Cropscience Ltd.	Systemic
Headline	Pyraclostrobin 20% WG	BASF Pvt. Ltd.	Broad spectrum, locally systemic
Imbrex	Fluxapyroxad EC	BASF Pvt. Ltd.	Broad spectrum
Folicur	Tebuconazole 25.9% EC	Bayer Cropscience Ltd.	Systemic
Vitavax	Carboxin 37.5% + Thiram 37.5% DS	Dhanuka Agritech Limited	Systemic and contact

Compatibility of *Trichoderma sp.* with fungicides under *in vitro* conditions

Ten fungicides were tested for compatibility with the five *Trichoderma* isolates using the poisoned food method Dhingra and Sinclair (1995). For each treatment, 150 ml of PDA was taken in a 250 ml conical flask and sterilized. To achieve the appropriate concentration at lukewarm temperature, the test chemical was added to this medium in the quantity according to recommended doses and thoroughly mixed by shaking the flasks. Aseptically poured into sterilized Petri plates, the poisoned medium was allowed to solidify. Three replications were maintained for each treatment. With a sterile cork borer, mycelial discs (5 mm in diameter) of the test isolates of *Trichoderma*, Tr-1, Tr-5, Tr-7, Tr-13, and Tr-15 culture, were removed from the colony's periphery and placed in

the center of poisoned media in each Petri plate. *Trichoderma spp.* discs (5mm) were placed onto Petri plates with untreated medium (i.e., without chemicals) to maintain appropriate controls. At 28°C in a BOD, all of the inoculated Petri plates were incubated. After five days of inoculation, the colony diameter of all isolates of *Trichoderma spp.* was measured in, with the average value being recorded. The following formula (Vincent, 1927) was used to compute the percentage of inhibitions of the fungal mycelial growth.

$$PI = \frac{C - T}{C} \times 100$$

C

PI = Percentage inhibition

C = Radial growth of the *Trichoderma* in control plate (cm)

T = Radial growth of the *Trichoderma* in treatments (cm)

Results

It is highly challenging to manage soil-borne plant diseases using just one method of control. In order to control the diseases carried on by several soil borne pathogens, integrated disease management (IDM) is a viable strategy. Since biological control has grown to be one of the key elements of IDM, it is inevitable that biocontrol agents will frequently be applied alongside fungicides to plants, soil, or both, which could cause either synergism or antagonistic interactions between them. The compatibility of *Trichoderma sp.* with various fungicides regularly used in annual crops was assessed in this work *in vitro* because most commonly used fungicides have either positive, negative, or neutral effects on biocontrol agents.

Compatibility of *Trichoderma sp.* with fungicides

Total five (Tr-1, Tr-5, Tr-7, Tr-13, and Tr-15) *Trichoderma* isolates were tested for compatibility with systemic, contact and broad spectrum fungicides at three distinct concentrations: 50 ppm, 100 ppm, and 150 ppm.

1. Amistar Top (Azoxystrobin 18.2% + Difenoconazole 11.4% SC)

Amistar Top was compatible (100%) with *Trichoderma* isolates Tr-15 and Tr-7 at 50, 100 and 150 ppm concentrations, and it inhibits the mycelial growth to 5.55% at 150 ppm in Tr-7, according to an *in-vitro* compatibility investigation of *Trichoderma* isolates with this fungicide (Table 2, Plate 1 & 2). However, in Tr-1, Tr-5, and Tr-13, the percentage inhibition of mycelial growth of *Trichoderma* rises from 50 to 150 ppm. Tr-13 exhibited inhibition of 19.63% (50 ppm), 24.81% (100 ppm), and 28.88% (150 ppm). In Tr-1 inhibition % ranges from 36.66 to 54.81%, and similarly in Tr-5 it

ranges from 21.11 to 34.81%. Comparing Tr-1 to other *Trichoderma* isolates and the control, this fungicide was found to be less compatible with Tr-1.

Table 2: Compatibility of *Trichoderma* isolates with Amistar Top fungicide

Amistar Top (Azoxystrobin 18.2% + Difenoconazole 11.4% SC)						
Isolates	Concentrations					
	50 ppm		100 ppm		150 ppm	
	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)
Tr-1	57.00	36.66	53.67	40.36	40.67	54.81
Tr-5	71.00	21.11	66.00	26.66	58.67	34.81
Tr-7	90.00	0.00	90.00	0.00	85.00	5.55
Tr-13	72.33	19.63	67.67	24.81	64.00	28.88
Tr-15	90.00	0.00	90.00	0.00	90.00	0.00
Control	90.00	-	90.00	-	90.00	-
SEm±	0.72	-	1.43	-	0.90	-
CD 0.01%	3.11	-	6.16	-	3.89	-

2. Saaf (Carbendazim 12% + Mancozeb 63% WP)

Saaf significantly suppressed the development of *Trichoderma* isolates Tr-1, Tr-7, and Tr-15 at all tested concentrations, according to an *in vitro* compatibility investigation of *Trichoderma* isolates with this fungicide (Table 3, Plate 1 & 2). As compared to isolate Tr-5, which grew up to 16 mm at 50 and 100 ppm and 2.33 mm at 150 ppm. The mycelial growth in Tr-13 was recorded 13.33 mm (50 ppm), 10.67 mm (100 ppm), and 0 mm (150 ppm). This reveals that even at concentrations of 50 ppm and higher, none of the isolates were compatible with Saaf.

SAAF (Carbendazim 12% + Mancozeb 63% WP)						
Isolates	Concentrations					
	50 ppm		100 ppm		150 ppm	
	Mycelial Growth (mm)	Percent inhibition (%)	Mycelial Growth (mm)	Percent Inhibition (%)	Mycelial growth (mm)	Percent Inhibition (%)
Tr-1	0.00	100.00	0.00	100.00	0.00	100.00
Tr-5	16.00	82.22	16.00	82.22	2.33	97.41
Tr-7	0.00	100.00	0.00	100.00	0.00	100.00
Tr-13	10.67	88.14	13.33	85.18	0.00	100.00
Tr-15	0.00	100.00	0.00	100.00	3.00	96.66

Control	90.00	-	90.00	-	90.00	-
SEm±	0.36	-	0.59	-	1.55	-
CD 0.0 1%	1.56	-	2.56	-	6.70	-

Table 3: Compatibility of *Trichoderma* isolates with SAAF fungicide

3. Seedkot (Thiram 75% WS)

The compatibility tests of *Trichoderma* isolates with the various tested concentrations of thiram (Table 4, Plate 1 & 2) revealed excellent results at all concentrations of 50 ppm, 100 ppm, and 150 ppm. compared to the control, where Tr-2 at all doses showed growth of 90.00 mm. The isolate Tr-15 had 100% compatibility (90 mm growth) up to 100 ppm, but at 150 ppm, it began to fall a bit, with 85.33 mm (5.18% inhibition) of mycelial growth. The isolates Tr-13 and Tr-7 were shown to be 100% (90 mm) compatible at 50 ppm, however Tr-1 showed an inhibition of 18.88% mycelial growth. However, as the concentration of Thiram was increased from 100 ppm to 150 ppm, the percentage of growth inhibition for both isolates of *Trichoderma* continued to rise, reaching up to 3.33-15.18% inhibition in isolate Tr-13, 17.77-39.25% in isolate Tr-7, and 19.63-29.63% in isolate Tr-1, respectively.

Table 4: Compatibility of *Trichoderma* isolates with Seedkot fungicide

Seedkot (Thiram 75% WS)						
Isolates	Concentrations					
	50 ppm		100 ppm		150 ppm	
	Mycelial Growth (mm)	Percent Inhibition (%)	Mycelial Growth (mm)	Percent Inhibition (%)	Mycelial Growth (mm)	Percent Inhibition (%)
Tr-1	73.00	18.88	72.33	19.63	63.33	29.63
Tr-5	90.00	0.00	90.00	0.00	90.00	0.00
Tr-7	89.00	1.11	74.00	17.77	54.67	39.25
Tr-13	90.00	0.00	87.00	3.33	76.33	15.18
Tr-15	90.00	0.00	90.00	0.00	85.33	5.18
Control	90.00	-	90.00	-	90.00	-
SEm±	0.53	-	2.63	-	1.86	-
CD 0.0 1%	2.27	-	11.35	-	8.04	-

4. Taqat (Captan 70%+ Hexaconazole 5% WP)

The compatibility assay of *Trichoderma* isolates was tested with three concentrations of Taqat, it revealed that Taqat significantly inhibited the growth of *Trichoderma* isolates at 50 ppm concentration, with percent inhibitions of Tr-5, Tr-13, Tr-1, Tr-15, and T7 recorded as 87.77%, 86.30%, 83.70%, 79.63, and 78.88%, respectively, in comparison to control (Table 5, Plate 1 & 2). All *Trichoderma* isolates showed decreased growth when Taqat concentration was increased from 100 to 150 ppm, and inhibition was observed for Tr-1(87.77-88.88 %), Tr-5(91.11-94.07 %), Tr-7(86.30-90.37%), Tr-13(91.11-94.44%), and Tr-15(88.52-92.58%), revealed incompatibility of *Trichoderma* with Taqat at all the concentrations tested.

Table 5: Compatibility of *Trichoderma* isolates with Taqat fungicide

Taqat (Captan 70%+ Hexaconazole 5% WP)						
Isolates	Concentrations					
	50 ppm		100 ppm		150 ppm	
	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)
Tr-1	14.67	83.70	11.00	87.77	10.00	88.88
Tr-5	11.00	87.77	8.00	91.11	5.33	94.07
Tr-7	19.00	78.88	12.33	86.30	8.66	90.37
Tr-13	12.33	86.30	8.00	91.11	5.00	94.44
Tr-15	18.33	79.63	10.33	88.52	6.67	92.58
Control	90.00	-	90.00	-	90.00	-
SEm±	0.62	-	0.56	-	0.47	-
CD 0.01%	2.69	-	2.42	-	2.04	-

5. Curzate M8 (Cymoxanil 8%+ Mancozeb 64% WP)

Curzate M8 was found to be compatible with all *Trichoderma* isolates up to 100 ppm in an *in vitro* compatibility testing of *Trichoderma* (Table 6, Plate 1 & 2). With the exception of Tr-5, which showed inhibition of 6.30% (50 ppm) and 15.92% (100 ppm), whereas, all isolates of *Trichoderma* showed full Petri plate growth of 90 mm at 50 and 100 ppm concentrations. This reveals their 100% compatibility at 50 and 100 ppm concentrations. However, the compatibility continued to decline when Curzate M8 concentration was further increased. It was found that *Trichoderma* isolates Tr-15, Tr-1, and Tr-5, respectively, each showed 10.74%, 11.47%, and 22.96% growth inhibition at 150 ppm concentration when compared to the control, while Curzate M8 was found to be 100% compatible with isolates Tr-7 and Tr-13 at 150 ppm concentration.

Table 6: Compatibility of *Trichoderma* isolates with Curzate M8 fungicide

Curzate M8 (Cymoxanil 8%+ Mancozeb 64% WP)						
Isolates	Concentrations					
	50 ppm		100 ppm		150 ppm	
	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)
Tr-1	90.00	0.00	90.00	0.00	79.67	11.47
Tr-5	84.33	6.30	75.67	15.92	69.33	22.96
Tr-7	90.00	0.00	90.00	0.00	90.00	0.00
Tr-13	90.00	0.00	90.00	0.00	90.00	0.00
Tr-15	90.00	0.00	90.00	0.00	80.33	10.74
Control	90.00	-	90.00	-	90.00	-
SEm±	0.59	-	0.59	-	0.41	-
CD 0.01%	2.56	-	2.56	-	1.76	-

6. Nativo (Tebuconazole 50%+ Trifloxystrobin 25% WG)

Nativo's compatibility study showed that *Trichoderma* isolates could not grow well in the presence of this fungicide, even at lower concentration of 50 ppm (Table 7, Plate 1 & 2). Inhibition rates for Tr-1, Tr-5, Tr-7, Tr-13, and Tr-15 isolates were calculated to be 81.85%, 83.33%, 91.11%, 85.18%, and 75.55%, respectively. However, as the concentration of Nativo was increased from 50 ppm to 100 ppm, greater percentage of *Trichoderma* isolates' development was inhibited, reaching as high as 84.44% (Tr-1), 89.25% (Tr-5), 94.44% (Tr-7), 95.92% (Tr-13), and 88.14% (Tr-15), respectively. The colony growth of the *Trichoderma* isolates Tr-5 and Tr-13 was thoroughly examined and fully inhibited upon raising the concentration of Nativo from 100 ppm to 150 ppm, while Tr-1, Tr-15, and Tr-7 each shown 90.74%, 90.74%, and 94.44% inhibition, respectively.

Table 7: Compatibility of *Trichoderma* isolates with Nativo fungicide

Nativo (Tebuconazole 50%+ Trifloxystrobin 25% WG)						
Isolates	Concentrations					
	50 ppm		100 ppm		150 ppm	
	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)
Tr-1	16.33	81.85	14.00	84.44	8.33	90.74
Tr-5	15.00	83.33	9.67	89.25	0.00	100.00
Tr-7	8.00	91.11	5.00	94.44	5.00	94.44
Tr-13	13.33	85.18	3.67	95.92	0.00	100.00
Tr-15	22.00	75.55	10.67	88.14	8.33	90.74
Control	90	-	90	-	90	-
SEm±	1.86	-	1.65	-	0.38	-
CD 0.01%	8.06	-	7.13	-	1.66	-

7. H
eadlin
e

(Pyraclostrobin 20% WG)

Headline compatibility test (Table 8, Plate 1 & 2) showed that there was a 1.85-48.52% reduction in growth of the isolates of *Trichoderma* starting at 50 ppm concentration compared to control. The colony diameter continued to decline as Headline concentration was raised. As compared to the control, *Trichoderma*'s growth was inhibited by a percentage ranging from 1.47 to 57.77% at 100 ppm and 6.66 to 58.88% at 150 ppm. This demonstrates that Headline and isolates of *Trichoderma* are only moderately compatible.

Table 8: Compatibility of *Trichoderma* isolates with Headline fungicide

Headline (Pyraclostrobin 20% WG)						
Isolates	Concentrations					
	50 ppm		100 ppm		150 ppm	
	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)
Tr-1	88.33	1.85	86.67	3.70	84.00	6.66

Tr-5	46.33	48.52	38.00	57.77	37.00	58.88
Tr-7	74.00	17.77	65.67	27.03	37.33	58.52
Tr-13	73.67	18.14	72.00	20.00	57.00	36.66
Tr-15	88.33	1.85	88.67	1.47	77.67	13.70
Control	90.00	-	90.00	-	90.00	-
SEm±	1.01	-	0.94	-	1.56	-
CD 0.0 1%	4.36	-	4.07	-	6.73	-

8. Imbrex (Fluxapyroxad EC)

Fluxapyroxad was found to be highly compatible with all *Trichoderma* isolates in an *in vitro* compatibility study, with the exception of Tr-5 isolate, which exhibits mycelial growth of 84 mm, 77.33 mm, and 67.67 mm at 50 ppm, 100 ppm, and 150 ppm, respectively (Table 9, Plate 1 & 2). From a concentration of 100 ppm (89 mm) to 150 ppm (83 mm), Fluxapyroxad began to inhibit *Trichoderma* growth in Tr-7. At doses of 100 to 150 ppm, it was found that the growth rates of the *Trichoderma* isolates Tr-1, Tr-13, and Tr-15 did not show any significant reduction. The results revealed that *Trichoderma* and Fluxapyroxad are extremely compatible up to 150 ppm concentrations with most of the isolates.

Table 9: Compatibility of *Trichoderma* isolates with Imbrex fungicide

Imbrex (Fluxapyroxad EC)						
Isolates	Concentrations					
	50 ppm		100 ppm		150 ppm	
	Mycelial Growth (mm)	Percent Inhibition (%)	Mycelial Growth (mm)	Percent Inhibition (%)	Mycelial Growth (mm)	Percent Inhibition (%)
Tr-1	90	0.00	88	2.22	90	0.00
Tr-5	84	6.66	77.33	14.07	67.67	24.81
Tr-7	90	0.00	89	1.11	83	7.77
Tr-13	90	0.00	89.33	0.74	90	0.00
Tr-15	90	0.00	89.67	0.36	90	0.00
Control	90	-	90	-	90	-
SEm±	0.47	-	1.20	-	0.86	-
CD 0.0 1%	2.04	-	5.19	-	3.72	-

9. Folicur (Tebuconazole 25.9% EC)

With all of the *Trichoderma* isolates, the fungicide Folicur did not work very well. According to Table 10, (Plate 1 & 2) the percent growth inhibition ranged from 75.18 to

91.47% at 50 ppm, 84.44 to 94.07% at 100 ppm, and 90.36 to 100% at 150 ppm. At all concentrations, isolate wise compatibility falls off begins from at Tr-5, Tr-15, Tr-1, Tr-7, and Tr-13, respectively.

Table 10: Compatibility of *Trichoderma* isolates with Folicur fungicide

Folicur (Tebuconazole 25.9% EC)						
Isolates	Concentrations					
	50 ppm		100 ppm		150 ppm	
	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)
Tr-1	10.67	88.14	8.33	90.74	0.00	100
Tr-5	22.33	75.18	14.00	84.44	8.67	90.36
Tr-7	8.67	90.36	7.33	91.85	0.00	100
Tr-13	7.67	91.47	5.33	94.07	0.00	100
Tr-15	20.33	77.41	11.33	87.41	0.00	100
Control	90.00	-	90.00	-	90.00	-
SEm±	0.73	-	0.49	-	0.36	-
CD 0.0 1%	3.16	-	2.12	-	1.56	-

10. Vitavax (Carboxin 37.5% + Thiram 37.5% DS)

Trichoderma might develop with Vitavax at lower concentrations of 50 ppm (Table 11, Plate 1 & 2) according to a compatibility investigation. It was determined that the growth of the *Trichoderma* isolates Tr-13, Tr-15, Tr-1, Tr-7, and Tr-5 was inhibited by 0%, 0.36%, 1.11%, 1.85%, and 33.33%, respectively. However, when Vitavax concentration increased from 50 ppm to 150 ppm, the percent inhibition of *Trichoderma* isolates also increased, reaching values of 0.74-88.88% and 0.33-92.58%, respectively. Among all the isolates there was no significant decrease in mycelium growth was seen in Tr-15, indicating that isolate Tr-15 performed remarkably well against Vitavax and it was extremely compatible with Vitavax.

Table 11: Compatibility of *Trichoderma* isolates with Vitavax fungicide

Vitavax (Carboxin 37.5% + Thiram 37.5% DS)						
Isolates	Concentrations					
	50 ppm		100 ppm		150 ppm	
	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)	Mycelial growth (mm)	Percent inhibition (%)
Tr-1	89	1.11	63	30.00	36	60.00
Tr-5	60	33.33	10	88.88	6.67	92.58

Tr-7	88.33	1.85	74	17.77	60.33	32.96
Tr-13	90	0.00	80	11.11	54.67	39.25
Tr-15	89.67	0.36	89.33	0.74	87	3.33
Control	90	-	90	-	90	-
SEm±	1.25	-	0.64	-	0.79	-
CD 0.0 1%	5.42	-	2.76	-	3.43	-

DISCUSSION

Maheshwary *et al.*, (2020) observed that *Trichoderma* is incompatible with different contact, systemic, and combination fungicides may be related to the greater quantities applied as well as the test native *Trichoderma* under capacity for tolerance. Tebuconazole 50% + Trifloxystrobin 25% WG, Azoxystrobin 18.2% + Difenconazole 11.4% SC, and Carbendazim 12% + Mancozeb 63% WP were all determined to be completely incompatible with *Trichoderma*. These conclusions are supported by the current study's findings. The results support previous research from various researchers (Madhusudhan *et al.*, 2010; Rakholiya *et al.*, 2010; Ranganathswamy *et al.*, 2012; Sreeja and Girija, 2015; Rai *et al.*, 2016). The direct harmful chemical's impact on *Trichoderma* cells and spores may be the cause of the fungicides' inhibitory effects. Due to their capacity to break down compounds and innate resistance to most fungicides, antagonistic microorganisms may react differently to different fungicides (Papavizas, 1985). Azoxystrobin was moderately compatible with systemic fungicides, but tebuconazole, propiconazole, and carbendazim were incompatible. Their findings concur with those made by Bagwan (2010) and Bindu *et al.* (2011), who revealed that tebuconazole is incompatible with *Trichoderma*.

Azoxystrobin was shown to be extremely compatible with *T. harzianum* and *T. viride* among all the fungicides (0.00% inhibition at three tested concentrations). The fungicide with the highest inhibition was carbendazim. Ranganathswamy *et al.*, (2012) also observed that *Trichoderma* sp. was compatible with azoxystrobin.

Manadhar *et al.*, (2020) tested compatibility with fungicides where they found that Carbendazim, Saaf had complete inhibitory effect on all *Trichoderma*, irrespective of the isolates tested. While the fungicide Aver green (Cymoxanil 8%+ Mancozeb 64% WP) which has similar active ingredient as Dupont curzate M8 fungicide used in present study was found to be compatible with all *Trichoderma* isolates. These results are similar with the results found in present study.

Kumar *et al.*, (2019) findings indicate fungicide compatibility analysis of *Trichoderma* which revealed that with thiram 75%WP *T. viride* is more than 70 % compatible upto 100 ppm. According to earlier findings, agrochemicals and bio control agents that can withstand a particular degree of fungicides were combined to eradicate illnesses [De Cal *et al.*, 1994]. Similar to this, Bagwan (2010) observed that *Trichoderma harzianum* and *Trichoderma viride* are compatible with thiram, copper

oxychloride, and Mancozeb at 0.2%. The use of microorganisms that combat plant pathogenic fungus is risk-free because fungicides frequently have negative effects on non-target species [Bentez *et al.*, 2004].

Compatibility tests were conducted under *in vitro* condition to find out safer fungicides. For this different fungicide were tested against *Trichoderma* isolates, results indicate that among the fungicides tested, *Trichoderma* was most sensitive to captan, tebuconazole, vitavax, propiconazole and chlorothalonil (Bagwan, 2010).

Laboratory tests were done by Wedajo (2015) to determine whether certain *Trichoderma* species may be used with fungicides. The findings showed that both *Trichoderma* species were 50% compatible with both fungicides at the curzate (400 ppm) and sancozeb (600 ppm) concentrations that were chosen. As the concentration of both fungicides got higher, it was found that the percent inhibition of radial growth in AUT1 and AUT2 gradually increased. At concentrations greater than the 1000 ppm employed in their investigation, both fungicides were capable of totally suppressing the growth of both *Trichoderma* species.

It might be possible to reduce disease in a similar way to how more fungicides are used if biological control agents and fungicides were combined. A fungicide application is decreased and the possibility of resistance developing is eliminated when antagonists are used with synthetic and non-synthetic chemicals. Mancozeb 75%WP and pyraclostrobin were the two fungicides that performed best in earlier experiments on *T. viride* compatibility. In keeping with Somasekhara *et al.* (1998) findings, carbendazim 50%WP was discovered to be very inhibiting.

Since *Trichoderma* species have no negative environmental effects, it is crucial to combine them with fungicides at lower concentrations rather than administering these chemicals alone to effectively manage fungal pathogens. Similar to this, [Srinivas and Ramakrishnan (2002)] stated that the combination of biological control agents and frequently used fungicides demonstrated favourable relationship by reducing the seed infection as compared to fungicide and the fungal antagonists independently. According to [Silimela and Korsten (2001)], the effectiveness of the biological control agent could be increased still further when used in conjunction with the suggested fungicide and at a lower concentration. In context with this, the antagonistic potential of *Trichoderma* species in terms of improved modes of action as well as increased hyper parasitism activity in the current investigation. The results of this screening will aid in choosing biological control agents that can be employed in association with specific fungicides at decreased doses to control plant pathogenic fungus.


Plate 1: Compatibility of *Trichoderma* isolates with different fungicides at various concentrations

Fungicides	Tr-1		Tr-5		Tr-7	
	50 150	100	50 150	100	50 150	100
Amistar Top						
Saaf						
Seedkot						
Taqat						
Curzate M8						
Nativo						
Headline						
Imbrex						

Folicur			
Vitavax			
Control			

Plate 2: Compatibility of *Trichoderma* isolates with different fungicides at various concentrations

Fungicides	Tr-13			Tr-15		
Conc. (ppm)	50	100	150	50	100	150
Amistar Top						
Saaf						
Seedkot						
Taqat						
Curzate M8						
Nativo						
Headline						



Conclusion

The current research demonstrates that various chemical fungicides both support and limit the growth of *Trichoderma* spp. For the integrated disease management of agricultural crops, it is possible to choose which fungicides are compatible with *Trichoderma* spp. and employ them in combination. It is conceivable for the chemical responses of *Trichoderma* isolates to vary, hence it is preferable to verify the compatibility of individual *Trichoderma* spp. before using them in integration rather than generalising the impact of chemicals on *Trichoderma* growth. Pathogens can be instantly controlled and kept under control for a longer amount of time by using a combination of bioagent and chemical. To determine the impact on disease control, field tests combining the *Trichoderma* spp. and suitable fungicides from this study are required.

References

- Adekunle AT, Cardwell KF, Florini DA, Ikotun T. Seed treatment with *Trichoderma* species for control of damping-off of cowpea caused by *Macrophomina phaseolina*. *Biocontrol Sci Tech*. 2001;11: 449-457.
- Bagwan NB. Evaluation of *Trichoderma* compatibility with fungicides, pesticides, organic cakes and botanicals for integrated management of soil borne diseases of soybean (*Glycine max* (L.) Merrill). *Int J Plant Prot*. 2010;3: 206-209.
- Benítez T, Rincón AM, Limón MC, Codón AC. Biocontrol mechanisms of *Trichoderma* strains. *Int Microbiol*. 2004;7: 249-260.

Bindu MG, Bhattiprolu SL, Balireddy V. Compatibility of biocontrol agent *Trichoderma viride* with various pesticides. *Journal Horticulture Science*. 2011;6(1):71- 73.

De Cal A, Pascual S, Melgarejo P. *In vitro* studies on the effects of fungicides on beneficial fungi of peach twig mycoflora. *Mycopathologia*. 1994;126: 15-20.

Dhingra, OD and Sinclair, JB. Basic plant pathology methods. CBS Publications and Distribution, New Delhi. 1995;335.

Dubey SC, Patil B. Determination of tolerance in *Hanetophorus cucumeris*, *Trichoderma viride*, *Gliocladium virens* and *Rhizobium* sp. to fungicides. *Indian Phytopathol*. 2001;54: 98-101.

Dubey SC, Suresh M, Singh B. Evaluation of *Trichoderma* species against *Fusarium oxysporum* f.sp. *ciceris* for integrated management of chickpea wilt. *Biological Control*. 2007;40:118-127.

Ehteshamul-Haque S, Zaki MJ, haffar A. Biological control of root rot diseases of okra, sunflower, soybean and mungbean. *Pak J Bot*. 1990;22: 121-124.

Elad Y, Chet I, Katan I. *Trichoderma harzianum* a biocontrol agent effective against *Sclerotium rolfisii* and *Rhizoctonia solani*. *Phytopathology*. 1980;70:119-121.

Gomez, K.A. and Gomez, A.A. Statistical procedures for agricultural research 2nd Ed., John Wiley and Sons, New York.1984

Harman GE, Howell CR, Viterbo A, Chet I, Lorito M. *Trichoderma* species--opportunistic, avirulent plant symbionts. *Nat Rev Microbiol*. 2004;2: 43-56.

Hjeljord L, Tronsmo A. *Trichoderma* and *Gliocladium* in biological control: an overview. In: *Trichoderma* and *Gliocladium*-Enzymes, Biological Control and Commercial Applications. (Eds.): Harma GE and Kubicek CP. Taylor & Francis Ltd, London, Great Britain. 1998;131-151.

Houdam R, Dutta BK. Compatibility of *Trichoderma atroviride* with fungicides against black rot disease of tea: an *in vitro* study. *J Int Academic Research for Multidisciplinary*. 2014;2: 25-33.

Jain, AK, Kumar, A, Chouhan, SS and Tripathi, SK. Cultural characteristics and evaluation of *Trichoderma* isolates against *Rhizoctonia solani* Kühn causing banded leaf and sheath blight of Little Millet. *Annals of Plant Protection Sciences*. 2017;25(1): 140-143.

Kharte, S, Kumar, A, Sharma, S, Ramakrishnan, RS, Kumar, S, Malvi, S, Singh Y. and Kurmi S. In vitro Evaluation of Fungicides and Bio-agents for the Management of Lentil Wilt caused by *Fusarium oxysporum* f. sp. *lentis*. *Biological Forum – An International Journal*. 2022;14(4): 489-495.

Kumar A, Bansal, RD, and Chelak YK. Compatibility of *Trichoderma viride* with Fungicides for Plant Disease Management. *Int. J. Pure App. Biosci.* 2019;7(3): 44-51.

Kumar A, Jain AK, Singh J, Tripathi SK and Tiwari RK. *In-vitro* studies on cultural characterization of a repository of local isolates of *Trichoderma* spp. from Madhya Pradesh. *Indian Phytopathology.* 2016;69(4s), 482-485.

Kumar A, Sahu TK, Bhalla A and Jain AK. Morphological characterization of *Trichoderma harzianum* from Madhya Pradesh. *Annals of Plant Protection Sciences.* 2013b;22(1): 190-239.

Kumar, A. and Sahu, TK. Use of local isolates of *Trichoderma* from Madhya Pradesh against *Rhizoctonia solani* causing wet root rot of chickpea. *Environment and Ecology.* 2015;33(4): 1553-1557.

Kumar, A., and Sahu, TK. Studies on substrate evaluation for mass multiplication of *Trichoderma* species and their plant growth promotion activity in tomato. *International journal of plant protection.* 2014;7(2): 382-388.

Kumar, A., Bohra, A., Mir, RR, Sharma, R., Tiwari, A., Khan, MW and Varshney, RK. Next generation breeding in pulses: Present status and future directions. *Crop Breeding and Applied Biotechnology,* 2021;21(s), e394221S13.

Kumar, A., Govil, M., Singh, S., Sharma, KK, Tripathi, SK, Tiwari, RK, Tripathi, AN and Singh, S. Role of Micro-organisms in Bioremediation: A Comprehensive Model Using *Trichoderma* spp. Handbook of research on uncovering new methods for ecosystem management through bioremediation. (2015). doi:10.4018/978-1-4666-8682-3.ch002.

Kumar, A., Jain, A. K., Sahu, T. K., Singh, T. K. and Shivcharan, S. Exploration of potential biocontrol agent *Trichoderma* spp. from Madhya Pradesh against *Fusarium oxysporum* f. sp. *ciceris* causing wilt of chickpea. *Environment and Ecology.* 2013a;31(2B), 877-882.

Kumar, A., Kumar, S., Srivastava, R. and Sharma, AK. Fungal biocontrol agents (BCAS) and their metabolites. In. *Agricultural Diversification: Problems and Prospects* (Eds. by A.K. Sharma, S. Wahab and R. Srivastava). I. K. International, New Delhi, 2009:44-56.

Kumar, A., Patel, A., Singh, SN and Tiwari, RK. Effect of *Trichoderma* spp. in Plant Growth Promotion in Chilli. *International Journal of Current Microbiology and Applied Science.* 2019;8(3):1574-1581.

Kumar, A., Sahu, TK, Bhalla, A. and Solanki, S. Influence of *Trichoderma* spp. against *Ustilaginoidea virens* inciting false smut of rice. *Environment and Ecology,* 2014;32(1): 163-168.

- Lewis, JA and Lumsden, RD. (2001). Biocontrol of damping off of greenhouse grown crops caused by *Rhizoctonia solani* with a formulation of *Trichoderma* spp. *Crop Protection* 20: 49- 56.
- Madhusudhan P, Gopal K, Haritha V, Sangale UR, Rao SVRK. Compatibility of *Trichoderma viride* with fungicides and efficiency against *Fusarium solani*. *Journal Plant Disease Science*. (2010);5(1):23-26.
- Maheshwary, NP, Gangadhara, NB, Amoghavarsha, C., Naik, MK, Satish KM and Nandish, MS. Compatibility of *Trichoderma asperellum* with fungicides. *The Pharma Innovation Journal* 2020;9(8): 136-140.
- Manandhar, S., Timila, RD, Karkee, A., Gupt, SK, and Baidya, S. Compatibility study of *Trichoderma* isolates with chemical fungicides. *The Journal of Agriculture and Environment*. (2020; 21:1-18
- Monte E. Understanding *Trichoderma*: between biotechnology and microbial ecology. *Int Microbiol*. 2001;4: 1-4.
- Okigbo RN, Ikediugwu FEO. Studies on biological control of postharvest rot of yam with *Trichoderma viride*. *J. Phytopathol*. 2000;148: 351-355.
- Papavizas GC, Lewis JA. Introduction and Augmentation of Microbial Antagonists for the Control of Soil-Borne Plant Pathogens. In: *Biological Control in Crop Production*, Papavizas, G.C. (Ed.). Allanheld and Qsmun, Totowa, New Jersey. 1981;305-322.
- Rai D, Bisht KS, Tewari AK. *In vitro* effect of newer fungicides on mycelia growth in biocontrol fungus *Trichoderma harzianum* (Th 14). *Journal Hill Agriculture*. 2016;7(1):162-164.
- Rakholiya KB. Efficacy of fungicides against *Trichoderma harzianum* and *Sclerotium rolfsii*. *International Journal Plant Protection*. 2010;3(2):406- 407.
- Ram, P., Mathur, K., Lodha, BC. Integrated management of rhizome rot of ginger involving biocontrol agents and fungicides. *J. Mycol. Plant Pathol*. 1999;29(3): 416-420.
- Ranganathswamy, M., Patibanda, AK, Chandrashekhar, GS, Sandeep, D., Mallesh, SB, Halesh kumar, HB. Compatibility of *Trichoderma* isolates with selected fungicides in vitro. *International Journal Plant Protection*. 2012;5(1):12-15.
- Silimela M, Korsten L. Alternative methods for preventing pre and post-harvest diseases and sunburn on mango fruits. *S.A. Mango Growers" Assoc. Yearbook*. 2001; 21: 39-43.
- Somasekhara, YM, Siddaramaiah, AL and Anilkumar, TB. Evaluation of *Trichoderma* isolates and their antifungal extracts as potential biological control agents against pigeonpea wilt pathogen, *Fusarium udum* Butler. *Current Research University of Agricultural Sciences Bangalore*, 1998;27(7-8): 158-160

Spiegel Y, Chet I. Evaluation of *Trichoderma* spp., as a biocontrol agent against soil borne fungi and plant-parasitic nematodes in Israel. *Integrated Pest Management Reviews* (Online) 1998;3: 169-175.

Sreeja SJ, Girija VK. Compatibility of *Trichoderma viride*, *Pseudomonas fluorescens* and *Rhizobium* spp. with selected fungicides. *Plant Disease Research*. 2015;30(2):188-189.

Srinivas P, Ramakrishnan G. Use of native microorganisms and commonly recommended fungicides in integrated management of rice seed borne pathogens. *Annu Plant Prot Sci*. 2002;10: 260-264.

Srivastava, R., Joshi, M., Kumar, A., Pachauri, S. and Sharma, AK. Biofertilizers for sustainable agriculture. In. *Agricultural Diversification: Problems and Prospects* (Eds. By A.K. Sharma, S. Wahab and R. Srivastava). I.K. International, New Delhi, 2009;5:7-71.

Sun MH, Liu XZ. Carbon requirements of some nematophagous, entomopathogenic and mycoparasitic hyphomycetes as fungal biocontrol agents. *Mycopathologia*. 2006;161: 295-305.

Tapwal A, Kumar S, Gautam N, Pandey S. Compatibility of *Trichoderma viride* for Selected Fungicides and Botanicals. *International Journal of Plant Pathology*. 2012;3: 89-94.

Vincent, JM. Distortion of fungal hyphae in the presence of certain inhibitors., *Nature*. 1947;159: 850.

Wedajo, B. Compatibility Studies of Fungicides with Combination of *Trichoderma* Species under *In vitro* Conditions. *Virol-mycol*. 2015;4: 149.

Yedidia II, Benhamou N, Chet II. Induction of defense responses in cucumber plants (*Cucumis sativus* L.) By the biocontrol agent *Trichoderma harzianum*. *Appl Environ Microbiol*. 1999;65: 1061-1070.