

## Review Article

### **Central Venous Catheter-Associated Deep Venous Thrombosis in Critically Ill Children in Tanta Pediatric Intensive Care Unit**

**Comment [DM1]:** Factor VIII predictivity and effectiveness of thromboprophylaxis shall be added as mentioned in the conclusion

#### **Abstract**

**Background:** The presence of a central venous catheter (CVC) and admission to the intensive care unit is the most important risk factors for deep venous thrombosis (DVT) in children. At least 18% of critically ill children with a CVC develop radiologically confirmed catheter-associated thrombosis (CAT). Clinically apparent thrombosis occurs in 17.5% of critically ill children with a CVC.

**Comment [DM2]:** Better sentence explaining the association of CVC, DVT and ICU shall be mentioned here.

**Comment [DM3]:** Reference shall be mentioned

**Methods:** This prospective observational study was carried out on 55 pediatric patients from 2.5 to 108 months (28 males and 27 females) admitted to PICU, Tanta University Hospital. They were subdivided into: Group I: Fifteen patients with FVIII activity <100 % (low normal)  
Group II: Twenty patients with FVIII activity from 100 % to 150 % (high normal)  
Group III: Twenty patients with FVIII activity >150 %.

**Comment [DM4]:** Study period shall be mentioned here, number of cases studied shall be mentioned in the result section

**Comment [DM5]:** 1<sup>st</sup> time mentioning the abbreviation full form shall be mentioned

**Comment [DM6]:** Factor VIII activity units shall be mentioned

After correction of dehydration (if present); All patients in the three groups: FVIII and D-dimer, were collected on the day of insertion of CVC (all patients from the start received thromboprophylaxis in the form of (Enoxaparin Sodium®) with dose (0.5 mg/kg subcutaneous every 12 hours) and after one week of enrollment.

**Comment [DM7]:** Full form shall be written

**Comment [DM8]:** Regarding alteplase is not mentioned here

**Results:** FVIII activity was a significant predictor for central venous line thrombosis with cut-off value >152, sensitivity 100, specificity 80, PPV 93.7, and NPV 100.

**Comment [DM9]:** D-dimer value is not commented here

**Conclusion:** F VIII activity was a significant predictor for central venous line thrombosis.

**Comment [DM10]:** Conclusion shall be in par with the title of the study

Enoxaparin sodium® was effective as thromboprophylaxis with a dose (of 0.5 mg/kg subcutaneously every 12 hours) within the normal range of factor VIII (up to 150%).

Alteplase® dosing for CVC occlusion based on patient weight category and the volume of the affected catheter lumen with an efficacy of 73.33%.

## Introduction

Deep venous thrombosis is a significant problem in critically ill children. The most important risk factor for DVT is the presence of a CVC. DVT develops in approximately 17.5% of critically ill children with CVC. The risk for CVC-related DVT is low within 24 hours after insertion of the CVC but significantly increases thereafter. Although routinely used in critically ill, pharmacologic thromboprophylaxis is not recommended in children with CVC due to the paucity of studies to support this practice. It would be ideal to provide thromboprophylaxis only to critically ill children at high risk for CVC-related DVT. No test has been known to identify these children [1].

The no tunneled type of CVC is usually inserted when children are critically ill and likely hypercoagulable. Coagulation factors, particularly factor VIII, increase during critical illness. Factor VIII activity >150 IU/dL is associated with 2.6 to 4.8-fold odds of developing DVT in adults [2].

There are numerous risk factors that contribute to the development of an abnormal thrombus. Identification of the causative factors of VTE as well as determining the reversibility of these factors is very important in planning the therapy, in terms of length and intensity of the interventions, and assessment of the risk of recurrence of the thrombosis [3].

The goals of VTE therapy are to prevent clot extension, prevent embolism, restore venous patency, limit the long-term sequelae, and reduce the risk of recurrence. These goals are achieved by rebalancing the hemostatic system with anticoagulant and thrombolytic therapy and, when possible, reversal of the causative factors[4].

## Methods

This prospective observational study was carried out on 55 pediatric patients from 2.5 to 108 months (28 males and 27 females) admitted to PICU, Tanta University Hospital.

**Comment [DM11]:** Aim and Objectives of the study shall be mentioned after the introduction.

**Comment [DM12]:** Reference shall be mentioned

**Comment [DM13]:** These numbers shall be mentioned in the results, duration of the study shall be mentioned

The duration of research was conducted from December 2020 to December 2022. Written informed consent was obtained from the guardians of all patients included in the study. The study was approved by the ethics committee of the Faculty of Medicine, Tanta University.

Inclusion criteria: Ages from 2.5 months to 108 months with a no tunneled CVC inserted within the previous 24 hours were enrolled. Patients on mechanical ventilation., Patients with hemodynamic instability and Patients received inotropes.

**Comment [DM14]:** Any reason of choosing this age group shall be mentioned. If this is the statistics of age group of present study, these numbers shall be mentioned in the result.

**Comment [DM15]:** Sentence shall be completed

Children with documented thromboembolic events within the past 3 months. Children received anticoagulants within the past 3 months. Children received hemostatic support, particularly fresh frozen plasma or cryoprecipitate were excluded.

**Comment [DM16]:** Sentence shall be reframed

Patients were divided into 3 groups:

**Comment [DM17]:** Reason for dividing these patients based on FVIII shall be mentioned.

Group I: Fifteen patients with FVIII activity <100 % (low normal)

Group II: Twenty patients with FVIII activity from 100 % to 150 % (high normal)

Group III: Twenty patients with FVIII activity >150 %.

All patients received thromboprophylaxis in the form of (Enoxaparin Sodium<sup>®</sup>) with a dose (of 0.5 mg/kg subcutaneously every 12 hours).

**Comment [DM18]:** In the introduction, it has been mentioned that routinely in pediatric patients pharmacological thrombolytic prophylaxis is not recommended.

**Comment [DM19]:** Shall be removed.

All patients enrolled in this study were evaluated by Full history taking. Presence of significant immobility (limitation in independent purposeful physical movement of body or one or more extremities from baseline > 48 hours). Presence of chronic VTE risk factors during major surgery, major injury, or during periods of infection and inflammation, Inflammatory bowel disease., Nephrotic syndrome. Cystic fibrosis., Need for PICU., Need for mechanical ventilation. and Use of total parenteral nutrition.

**Comment [DM20]:** Complete the sentence. Meaning not conveyed

Routine investigations including Complete blood count, Prothrombin time and INR, Partial Thromboplastin time, Liver function tests (Alanine transferase, Aspartate transferase and serum albumin), C-reactive protein and erythrocyte sedimentation rate [5,6].

**Comment [DM21]:** Complete the sentence. Meaning is not conveyed

Scoring systems for patients: Pediatric Risk for mortality scoring III: obtained within 24 hr. of admission [7]. Pediatric Sequential Organ Failure Assessment scoring: obtained from each patient daily and recorded 1<sup>st</sup> day and after one week of admission [8].

All patients enrolled in this study After correction of dehydration (if present): FVIII and D-dimer, were collected on the day of insertion of CVL (all patients from the start received thromboprophylaxis in the form of (Enoxaparin Sodium<sup>®</sup>) with dose (0.5 mg/kg subcutaneous every 12 hours) and after one week of enrollment[9].

**Comment [DM22]:** Small letter a shall be used

**Comment [DM23]:** Repetition of same information shall be avoided

On the day of enrollment, Ultrasound imaging was performed on the central vein where the CVC was inserted and on both lower extremities, the most common sites of non-CVC-related DVT[10]. The US was repeated on the site of insertion of the CVC within 24 hours of its removal or one week after insertion of the CVC, when the clinical team suspected CVC-related DVT, or when the subject was unlikely to survive for the next 24 hours.

Blinded, certified radiology technicians at each hospital performed the US using the EDAN model (DUS 60) ISO 9000, ISO 13485 Origin: CHINA Ultrasound Imaging Platform with 7-15 MHz linear transducers[11].

#### **Statistical analysis of the data**

Data were fed to the computer and analyzed using IBM SPSS software package version 20.0. (Armonk, NY: IBM Corp). (128) Qualitative data were described using numbers and percentages. The Kolmogorov-Smirnov test was used to verify the normality of distribution Quantitative data were described using range (minimum and maximum), mean, standard deviation, and median. The significance of the obtained results was judged at the 5% level[12].

The used tests were: Chi-square test (For categorical variables, to compare between different groups). Student t-test (For normally distributed quantitative variables, to compare between two studied groups). ANOVA with repeated measures (For normally distributed quantitative variables, to compare between more than two periods or stages, and Post Hoc test (LSD) for pairwise

comparisons).Mann Whitney test (For abnormally distributed quantitative variables, to compare between two studied groups).Linear correlation coefficient (r): For detection of correlation between two quantitative variables in one group.The power of significance was evaluated as follows: Probability level (P-value) >0.05=Non significant (NS). P-value <0.05=Significant.

## **Results**

The demographic and anthropometric data of the studied patients was presented in table (1):There was no significant difference throughout the study between the studied groups.

Regarding the comparison of US between the studied groups:There was significant increase in the incidence of IV echoic material in group III compared with groups I, and II. Otherwise, there was no statistically significant difference throughout the study figure (1).

Regarding the Comparison of D-dimer between the studied groups: There was significant decrease of D-dimer on the 7th day compared with the 1st day in groups I, and II compared with group III.

Regarding the Comparison of FVIII activity between the studied groups: There was significant increase in FVIII activity in group III compared with groups I, and II on 1st day. Also, there was significant increase in FVIII activity in group III compared with groups I, and II on the 7th day)figure (2).

Regard Effect of Enoxaparin sodium® between the Studied Groups: There was significant increase in CVC occlusion in group III compared with groups I, and II Table (2).

Regard Effect of Alteplase® between the Studied Groups: There was significant increase in Patients who received Alteplase® in group III compared with groups I, and II.

Regarding Pediatric Risk for Mortality III Scoring Score in Studied Groups:There was significant increase in the risk of mortality in group III compared with groups I, and II.Otherwise, there was no significant difference throughout the study.

Regarding correlation of FVIII with other parameters: There was positive correlation between FVIII activity with ESR on the 1st and 7th days. There was negative correlation between FVIII activity with PT and PTT on the 1st and 7th daysTable (3).

Regarding the ROC curve for FVIII activity for predicting catheter-related thrombosis in critically ill children:FVIII activity was a significant predictor for central venous line thrombosis with cut-off value>152, sensitivity100, specificity 80, PPV 93.7, and NPV 100figure (3).

### Discussion

The present study showed that regarding age, weight, and sex there was no significant difference in age and sex throughout the study between the studied groups. The non-significant difference found in age, weight, and sex demonstrates the non-bias choice of the study cases.

The present study showed that Hb, WBCs, PLTs, PT, PTT, and ESR there was no significant difference between the studied groups during the study period. The non-significant difference proved in Hb, WBCs, PLTs PT and ESR demonstrates the non-bias choice of the cases.

**Comment [DM24]:** Shall be mentioned once for all the patient particulars, haematology and biochemistry parameters

The present study showed that ALT, AST, Albumin, and CRP there was no significant difference between the studied groups during the study period. The non-significant difference proved in ALT, AST, Albumin, and CRP demonstrates the non-bias choice of the cases.This was in accordance with, Kamphuisen et al. [13] conducted two groups of patients with or without pulmonary embolism had factor FVIII activity <100 % - from 100 % to 150 % - activity >150 %. They found that CRP is not a reliable parameter to determine whether the FVIII level is the consequence of an acute phase reaction. Similarly, Tsai et al. [14] conducted a prospective study on a total of 159 venous thromboembolism patients and measured FVIII and CRP levels, in blood samples. They found that there was no association between venous thromboembolism and C-reactive protein levels

**Comment [DM25]:** Reframe the sentence

The present study showed that there was no difference in central line type and size throughout the study between the studied groups. This was in accordance with, Huibonhoa et al. [15] conducted a post hoc study on 236 children admitted to the PICU, who had a non-tunneled CVC. They observed that the size and duration of CVC were not different among the studies group. This was in contrast with, Li et al. [16] enrolled a retrospective study on patients aged from birth to 18 (months) who were admitted to PICU and underwent at least one CVC placement. The patient's catheter types and sizes were significantly associated with the occurrence of CADVT.

Comment [DM26]: Shall be removed

Comment [DM27]: Grammar shall be checked

The present study showed that regarding the comparison of the US between studied groups. There was an increase in the incidence of IV echoic material in group III compared with groups I, and II. Consistent with these results, Faustino et al. [17] showed that DVT was significantly diagnosed on the day of discharge or within 24 hr of its removal.

Comment [DM28]: Full form shall be used

The present study showed that regarding the effect of Enoxaparin sodium®, there was a significant increase in CVC occlusion in group III compared with groups I, and II. This was in accordance with, Kearon C et al. [18] collected and analyzed retrospectively Group 1 of Enoxaparin sodium®, at prophylactic dose of 50 anti-Xa IU/kg/dose every 12hr subcutaneously. This was in contrast with, Merli G.J et al. large randomized National institute of allergy and infectious diseases trial was bringing together three clinical trial platforms covering four continents and over 300 hospitals. Researchers suggest that therapeutic doses of anticoagulants are not only safe but also superior to preventive ones. [19]

The present study showed that regarding risk factors there was an increase in the incidence of risk factors (TPN or MV) in group III compared with groups I, and II. These may also predispose to thrombosis since they can induce endothelial cell damage with the release of pro-coagulant factors and platelet activation, leading to thrombus formation and deep vessel occlusion. [20]. This was in accordance with, Lasagni et al. [21] collected and analyzed retrospectively a total of 78 CVC-VTEs children (median age at onset was 19 and 17 months). They found that TPN was the most frequent

reason for CVC insertion. This was in contrast with, Sandoya et al. [22] conducted an observational, cross-sectional study on 35 patients to identify factors that are associated with the development of venous thrombosis in patients with CVC admitted to the intensive care unit. They found that adequate nutritional status or malnutrition was not associated with a higher prevalence of thrombosis.

The present study showed that regarding There was a significant decrease of D-dimer on the 7th day compared with the 1st day in groups I, and II compared with group III. This was in accordance with, Evensen et al., [23] who conducted a prospective cohort study to measure circulating levels of FVIII and D-dimer. They found that D-dimer and FVIII showed significant linear associations with VTE risk.

**Comment [DM29]:** Regarding shall be removed, small t shall be used after comma

The present study showed that regarding Alteplase®, there was a significant increase in patients who received Alteplase® in group III compared with groups II, and I. The catheter function was regained in 80% and 73.33% respectively in group II and group III. This was in accordance with Shen et al. [24] who published a retrospective pediatric subgroup analysis. This study also indicated that shorter times from observation of occlusion to intervention improved catheter patency outcomes. [25]

The present study showed that regarding a comparison of FVIII activity between the studied groups. There was an increase in FVIII activity in group III compared with groups I, and II on the 1st day. Also, there was an increase in FVIII activity in group III compared with groups I, and II. This was in accordance with, Faustino et al. [26] who conducted a prospective cohort study on 85 children who were admitted to the PICU within 24 hours after the insertion of a central venous catheter. They found that FVIII activity is significantly associated with incident CVC-related DVT.

**Comment [DM30]:** Shall be removed, comma shall be put after 'that'

The present study showed that regarding the ROC curve; FVIII activity was a significant predictor for central venous line thrombosis with cut-off value >152, sensitivity 100%, specificity of 80%,

PPV of 93.7%, and NPV of 100%. This was in accordance with, Huibonhoa et al. [15] found that the sensitivity and specificity of physical examination with CVC dysfunction included in detecting CADVT diagnosed with ultrasound were 29.2 % and 80.2 %, respectively.

**Conclusion:**

- F VIII activity was a significant predictor for central venous line thrombosis.
- Enoxaparin sodium® was effective as thromboprophylaxis with a dose (of 0.5 mg/kg subcutaneously every 12 hours) within the normal range of factor VIII (up to 150%).
- Alteplase® dosing for CVC occlusion based on patient weight category and the volume of the affected catheter lumen with an efficacy of 73.33%. Conflict of interest: none to declare.

**List of abbreviations:**

CADVT: Catheter associated deep venous thrombosis CVC: central venous catheter, DVT: deep venous thrombosis, ESR: erythrocytic sedimentation rate, PICU: pediatric intensive care unit, TPN: total parenteral nutrition, US: ultrasound, VTE: venous thromboembolism

**Declarations**

Ethics approval and consent to participate: Local ethics committee of Faculty of Medicine, Tanta University approved the study. The study is in accordance with the ethical standards of institutional research committee and with the 1964 Helsinki declaration and its later amendments. Written informed consent was obtained from the parents of all subjects of the study before enrolment.

Consent for publication: not applicable.

**References**

- 1- Alhazzani W, Lim W, Jaeschke RZ, Murad MH, Cade J and Cook DJ. Heparin thromboprophylaxis in medical-surgical critically ill patients: a systematic review and meta-analysis of randomized trials. *Crit Care Med.* 2013;41(9):2088-98.
- 2- Faustino EV, Lawson KA, Northrup V and Higerson RA. Mortality-adjusted duration of mechanical ventilation in critically ill children with symptomatic central venous line-related deep venous thrombosis. *Crit Care Med.* 2011;39(5):1151-6.
- 3- Ryan ML, Van Haren RM, Thorson CM, Andrews DM, Perez EA, Neville HL, et al. Trauma induced hypercoagulability in pediatric patients. *J Pediatr Surg.* 2014;49(8):1295-9.
- 4- Tichelaar V, Mulder A, Kluin-Nelemans H and Meijer K. The acute phase reaction explains only a part of initially elevated factor VIII:C levels: a prospective cohort study in patients with venous thrombosis. *Thromb Res.* 2012;129(2):183-6.
- 5- Schwaberg B, Schlatzer C, Freidorfer D, Bruckner M, Wolfsberger CH, Mileder LP, et al. The use of a disposable umbilical clamp to secure an umbilical venous catheter in neonatal emergencies-an experimental feasibility study. *Children (Basel).* 2021;8(12):8-10.
- 6- Liu YY, Wang YP, Zu LY, Zheng K, Ma QB, Zheng YA, et al. Comparison of intraosseous access and central venous catheterization in Chinese adult emergency patients: A prospective, multicenter, and randomized study. *World J Emerg Med.* 2021;12(2):105-10.
- 7- Kaur A, Kaur G, Dhir SK, Rai S, Sethi A, Brar A, et al. Pediatric Risk of Mortality III Score - Predictor of Mortality and Hospital Stay in Pediatric Intensive Care Unit. *J Emerg Trauma Shock.* 2020;13(2):146-50.
- 8- Matics TJ and Sanchez-Pinto LN. Adaptation and Validation of a Pediatric Sequential Organ Failure Assessment Score and Evaluation of the Sepsis-3 Definitions in Critically Ill Children. *JAMA Pediatr.* 2017;171(10):e172352.

- 9- Pietroboni PF, Carvajal CM, Zuleta YI, Ortiz PL, Lucero YC, Drago M, et al. Landmark versus ultrasound-guided insertion of femoral venous catheters in the pediatric intensive care unit: An efficacy and safety comparison study. *Med Intensiva (Engl Ed)*. 2020 Mar. 44 (2):96-100.
- 10- Bravo-Pérez C, de la Morena-Barrio ME, de la Morena-Barrio B, Miñano A, Padilla J, Cifuentes R, et al. Molecular and clinical characterization of transient antithrombin deficiency: A new concept in thrombophilia. *Am J Hematol*. 2022;97(2):216-25.
- 11- Luo YB, Gao Q, Guo J, Zhang Y, Sun J, Li WY, et al. Ultrasound-guided co-axial introducer needle biopsy in the diagnosis of eosinophilic cystitis in children. *Chin Med J (Engl)*. 2021;134(15):1872-3
- 12- Kirkpatrick LA, Feeney BC. et al; A simple guide to IBM SPSS statistics for version 20.0. Student ed. Belmont, Calif.: Wadsworth, Cengage Learning; 2013.
- 13- Kamphuisen PW, Ten Wolde M, Jacobs EM, Ullmann EF and Büller H. Screening of high factor VIII levels is not recommended in patients with recently diagnosed pulmonary embolism. *J ThrombHaemost*. 2003;1(10):2239-40.
- 14- Tsai AW, Cushman M, Rosamond WD, Heckbert SR, Tracy RP, Aleksic N, et al. Coagulation factors, inflammation markers, and venous thromboembolism: the longitudinal investigation of thromboembolism etiology (LITE). *Am J Med*. 2002;113(8):636-42.
- 15- Huibonhoa RMT, Pinto MG, Faustino EVS, Study E-D, Study FV and Investigators CT. Accuracy of physical examination in detecting central venous catheter-associated thrombosis in critically ill children. *Thromb Res* 2022;218:192-8. Davalos A, Castillo J, Alvarez-Sabin J, Secades JJ, Mercadal J, López S, Cobo E, Warach S, Sherman D, Clark WM, Lozano R. Oral citicoline in acute ischemic stroke. *Stroke* 2002; 33:2850–57.

- 16- Li H, Lu Y, Zeng X, Feng Y, Fu C, Duan H, et al. Risk factors for central venous catheter-associated deep venous thrombosis in pediatric critical care settings identified by fusion model. *Thromb J.* 2022;20(1):1-11.
- 17- Faustino EVS, Spinella PC, Li S, Pinto MG, Stoltz P, Tala J, et al. Incidence and acute complications of asymptomatic central venous catheter-related deep venous thrombosis in critically ill children. *J Pediatr.* 2013;162(2):387-91.
- 18- Kearon C, Ginsberg J.S, Julian, J.A, Douketis J Solymoss, S Ockelford P, et al. Fixed-Dose Heparin (FIDO) Investigators; et al. Comparison of fixed-dose weight-adjusted unfractionated heparin and low-molecular-weight heparin for acute treatment of venous thromboembolism. *JAMA* 2006, 296, 935–942.
- 19- Merli, G.J. and Groce, J.B. Pharmacological and Clinical Differences Between Low-Molecular-Weight Heparins. *P. T.* 2010, 35, 95–105 Overgaard K. The effects of citicoline on acute ischemic stroke: a review. *J Stroke Cerebrovasc Dis* 2014;23(7):1764–69.
- 20- Cortejoso L, Manrique-Rodríguez S, Fernández-Llamazares CM and Sanjurjo-Sáez M. Treatment and prophylaxis of catheter-related thromboembolic events in children. *Int J Pharm Pharm Sci* 2012;15(5):632-41.
- 21- Lasagni D, Nosadini M, Molinari AC, Saracco P, Pelizza MF, Piersigilli F, et al. Systemic catheter-related venous thromboembolism in children: data from the Italian registry of pediatric thrombosis. *Front Pediatr.* 2022;10:843643-50.
- 22- Sandoya A and Esthefany P. Risk factors for the development of venous thrombosis in patients with a central venous catheter admitted to the intensive care unit of the Roberto Gilbert Elizalde children's hospital. *Revista Ecuatoriana de Pediatría.* 2021;4:1-6.
- 23- Evensen LH, Folsom AR, Pankow JS, Hansen JB, Allison MA, Cushman M, et al. Hemostatic factors, inflammatory markers, and risk of incident venous thromboembolism: The Multi-Ethnic Study of Atherosclerosis. *J Thromb Haemost* 2021;19(7):1718-28.

- 24- Shen V, Li X and Murdock M. Recombinant tissue plasminogen activator (alteplase) for restoration of function to occluded central venous catheters in pediatric patients. *J Pediatr Hematol Oncol* 2003;25:38-45.
- 25- Ponc D, Irwin D and Haire WD. Recombinant tissue plasminogen activator (alteplase) for restoration of flow in occluded central venous access devices: a double-blind placebo-controlled trial. The cardiovascular thrombolytic to open occluded lines (COOL) efficacy trial. *J Vasc Interv Radiol* 2001;12:951-5
- 26- Faustino EVS, Li S, Silva CT, Pinto MG, Qin L, Tala JA, et al. Factor VIII may predict catheter-related thrombosis in critically ill children: a preliminary study. *Pediatr Crit Care Med*. 2015;16(6):49-60.

**Table (1): Comparison of Demographic Data between the Studied Groups**

		Groups						ANOVA	
		Group I		Group II		Group III		F	P
Age (Months)	<b>Range</b>	2.5	- 90	3	- 100	2.5	- 108	0.28	0.757
	<b>Mean</b>	22.467	± 29.330	16.400	± 23.525	20.950	± 24.388		
	<b>±SD</b>								
Weight (kg)	<b>Range</b>	5	- 25	5	- 30	3	- 25	0.04	0.99
	<b>Mean</b>	10.1	± 6.50	9.55	± 5.66	9.875	± 5.08		
	<b>±SD</b>								
Chi-Square		N	%	N	%	N	%	X <sup>2</sup>	p
Gender	<b>Male</b>	7	46.67	10	50.00	10	50.00	0.05	0.96
	<b>Female</b>	8	53.3	10	50.0	10	50.0		

$\chi^2$ , p:  $\chi^2$ , and p values for the **Chi-square test** for comparing the three groups, F, p: F, and p values for the **ANOVA test**.

**Table (2): Effect of Enoxaparin Sodium® with a dose (of 0.5 mg/kg subcutaneously every 12 hours). on the Studied Groups**

Enoxaparin sodium®	Groups						Chi-Square	
	Group I		Group II		Group III		X <sup>2</sup>	p
	N	%	N	%	N	%		
<b>CVC occlusion</b>	0	0.00	5	25.00	15	75.00	22.589	<0.001*
<b>No CVC occlusion</b>	15	100.00	15	75.00	5	25.00		
<b>Total</b>	15	100.00	20	100.00	20	100.00		

CVC: central venous catheter

**Table (3): correlation of FVIII with other parameters throughout the study**

	FVIII activity (%) 1 <sup>st</sup> day		FVIII activity (%) 7 <sup>th</sup> day	
	R	P	R	p
<b>Age (Months)</b>	0.056	0.686	0.002	0.990
<b>Weight (kg)</b>	0.107	0.438	0.058	0.674
<b>Hb (g/dL)</b>	0.003	0.985	0.048	0.730
<b>PLT (k/μL)</b>	-0.224	0.100	-0.221	0.105
<b>TLC (k/μL)</b>	0.041	0.768	0.001	0.996
<b>PT (second)</b>	-0.271	0.045*	-0.279	0.039*
<b>INR</b>	-0.120	0.383	-0.159	0.246
<b>PTT (second)</b>	-0.344	0.010*	-0.316	0.019*
<b>ESR 1ST HR (mm/hr.)</b>	0.301	0.025*	0.249	0.066
<b>S. Albumin (g/dl)</b>	-0.221	0.104	-0.200	0.143

<b>CRP</b>	0.166	0.226	0.155	0.260
<b>Central size</b>	0.150	0.275	0.091	0.510
<b>D-dimer &lt;0.5ug/ml 1<sup>st</sup> day</b>	0.237	0.082	0.221	0.105

PLT: platelets TLC: total leucocytic count PT: prothrombin time INR: international normalized ratio PTT: partial thromboplastin time ESR: erythrocytic sedimentation rate CRP: C-reactive protein




Figure (1): Comparison of Ivs between the Studied Groups.




Figure (2): Effect of Enoxaparin sodium® between the Studied Groups.




Figure (3): ROC curve for FVIII activity for predicting catheter-related thrombosis in critically ill children.