

# Effect of Phosphorus and Biofertilizers on Yield and Economics of Chickpea (*Cicer arietinum* L.)

## ABSTRACT

At Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj, (U.P. India), a field experiment was carried out during *Rabi* season of 2022, to study the yield and economics of chickpea (*Cicer arietinum* L.). The soil texture of the experimental plot was Sandy loam, with a pH 7.2, that was neutral with EC- 0.26 (dS/m), organic carbon (0.72%), available N (178.48kg/ha), available P (27.80kg/ha) and available K (233.24kg/ha). *Rhizobium sp.* (5g/kg seed), *Pseudomonas striata* (10g/kg seed), PSB (20g/kg seed) were used as biofertilizers, and phosphorus (40kg P<sub>2</sub>O<sub>5</sub>/ha, 50kg P<sub>2</sub>O<sub>5</sub>/ha, 60kg P<sub>2</sub>O<sub>5</sub>/ha) were used as fertilizers in 3 levels. The experiment was laid out in Randomized Block Design (RBD) with 10 treatments and replicated thrice. The treatment of Phosphorus (60kg/ha) + PSB (20g/ha) resulted in highest No. of pods/plant (30.90), No. of seeds/pod (1.53), seed yield (2570.74kg/ha) and stover yield (3583.33kg/ha), were all determined to be significant. The treatment (T<sub>9</sub>) with application of Phosphorus (60kg/ha) + PSB (20g/kg seed) also observed maximum gross returns (1, 54,165.87 INR/ha), net returns (1,13,725.57 INR/ha) and benefit cost ratio (2.81) in chickpea crop.

**Keywords:** Biofertilizers, Chickpea, PSB, *Rhizobium*, *Pseudomonas striata*, Yield, *Rabi*.

## 1. INTRODUCTION

Chickpea (*Cicer arietinum* L.) is the third most cultivated legume in the world. India stands out as one of the largest producers and consumers of its seeds (or seed); however, its production is insufficient to meet the needs of the internal market, and chickpea imported. It is recognized as a legume with high concentrations of proteins, nutrients, and carbohydrates. It is adaptable to wide climatic variation, has low production cost, and promotes biological fixation of atmospheric nitrogen [1]. India is the leading producer of chickpea contributing to about 70% of the world's chickpea production. In India, Madhya Pradesh (39%), Maharashtra (14%), Rajasthan (14%), Uttar Pradesh (7%), Karnataka (6%), and Gujarat (5%) are the major chickpea growing states. In India pulses are grown nearly in 28.83 m ha with an annual production of 25.72 m t and productivity of 0.8 t ha. Some of the states like Uttar Pradesh is about 8.24 m ha with an annual production of 9.97 m t and productivity of 1.08 t ha major producer of chickpea in India as advocated by Ministry of agriculture and Farmers Welfare". [2]

Phosphorus fertilization is among the main practices of crop management; yet, it is considered complex in tropical soils because of the high capacity of phosphorus for covalent adsorption to soil oxides and because of low natural availability of P to plants. In chickpea, balanced phosphorus nutrition is fundamental for establishing symbiosis with N<sub>2</sub> fixing rhizobacteria, and it stimulates nodulation, initial development of roots, plant growth, and seed yield and quality, among other aspects. It is affirm that fertilization with 60 kg ha<sup>-1</sup> of P<sub>2</sub>O<sub>5</sub> is sufficient for maximum production of chickpea seeds in *Vertisols* in India. A positive effect on relative growth rate, dry matter accumulation, nodulation, yield, and harvest index with application of 60 kg/ha of P<sub>2</sub>O<sub>5</sub>. Economic efficiency and high degree of protein are also reported by using this same phosphorus application rate [1]. The response of phosphorus

Comment [IA1]: affirmed

depends upon many factors like climate, variety and soil type and availability of nutrients during the period of growth. The requirement of phosphorus in legumes like chickpea is higher than other crops for their root development and metabolic activities. Phosphorus is the vital component of DNA, RNA, ATP and photosynthetic system and catalyses a number of bio chemical reactions from the beginning of seedling growth through to the formation of grain at maturity [3].

Biofertilizers are low cost, renewable sources of plant nutrients, which supplement chemical fertilizers. Biofertilizers solubilize plant nutrients like nitrogen and phosphorus through their activities in the soil or rhizosphere and make available to plants in a gradual manner. PSB solubilizes the insoluble forms of phosphates like tricalcium, iron and aluminium phosphates into available forms. PSB solubilizes insoluble inorganic phosphorus compounds by exerting organic acids, which is the primary mechanism of solubilizing of insoluble inorganic phosphates. Besides organic acids, production of chelating substances, mineral acids, siderophores and proton extrusion mechanism are also involved [4]. Biofertilizers may colonize the rhizosphere and promote growth by increasing the availability and supply of nutrients and growth stimulus to crop. Nitrogen fixer and phosphate solubilizing microorganisms play an important role in supplementing nitrogen and phosphorus to the plant, allowing a sustainable use of nitrogen and phosphate fertilizers. Some important strains are mentioned as plant growth promoting rhizobacteria (PGPR) and that can be used as biofertilizers i.e., *Rhizobium*, *Pseudomonas*, *Azospirillum*, *Azotobacter*, *Bacillus*, *Burkholderia*, *Erwinia*, *Mycobacterium*, *Flavobacterium* etc [5]. Phosphate-solubilizing bacteria exist in most soils. In *in vitro* conditions, they can improve P bioavailability by lowering the soil pH, solubilizing Pi, activating synthesized phosphatases, mineralizing organic P, and/or chelating P from Al<sup>3+</sup>, Ca<sup>2+</sup>, and Fe<sup>3+</sup> [6]. Phosphate solubilizing bacteria, when inoculated, secrete acidic substances and solubilize otherwise unavailable soil phosphorus. The culture can hence prove broad spectrum biofertilizers which may increase yield of crops (Legumes, vegetables etc) by 10-30%. Use of PSB culture increases nodulation, crop growth, nutrient uptake and crop yield [7]. Certain microorganisms such as phosphate solubilizing bacteria and fungi associated with the plant rhizosphere are known to convert insoluble inorganic phosphorus (P) into a soluble form that could be utilized by the plants [8]. Thus, adopting proper nutrient management practices in conjunction with PSB will help to improve the yield and quality of chickpea besides maintaining the soil fertility [9]. The present study was therefore, carried out to study the effect of phosphorus and biofertilizers on yield and economics of chickpea.

## 2. MATERIAL AND METHODS

The experiment was conducted during *Rabi* season of 2022, at Crop Research Farm, Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, U.P. India which is located at 25° 24' 42" N latitude above the mean sea level (MSL). The soil has sandy loam texture, with a pH of 7.2, organic carbon (0.72%), available N (178.48kg/ha), available P (27.80kg/ha) and available K (233.24kg/ha). The experiment was laid out in Randomized Block Design with ten treatments each replicated thrice. The experiment comprised three biofertilizer treatments and three levels of phosphorus when applied in combinations as follows;

Comment [IA2]: shown in table 1.

Table 1. Treatment combination

| S. No. | Treatment combination                           |
|--------|---|
| 1.     | Phosphorus 40kg/ha + <i>Rhizobium</i> sp.       |
| 2.     | Phosphorus 40kg/ha + <i>Pseudomonas striata</i> |
| 3.     | Phosphorus 40kg/ha + PSB                        |
| 4.     | Phosphorus 50kg/ha + <i>Rhizobium</i> sp.       |

5. Phosphorus 50kg/ha + *Pseudomonas striata*
6. Phosphorus 50kg/ha + PSB
7. Phosphorus 60kg/ha + *Rhizobium sp.*
8. Phosphorus 60kg/ha + *Pseudomonas striata*
9. Phosphorus 60kg/ha + PSB
10. Control (20-50-20 NPK kg/ha)

Five plants were tagged randomly at third week of germination for recording observations. At maturity of the pods, chickpea crop was harvested **treatment wise**. Followed by bundles and sun dried and later weighed by weighing balance for calculating biological yield. The seed yield was recorded after threshing of plants from each treatment and then converted into kg/ha. The observation like number of pods/plant, number of seeds/pod, seed yield and stover yield were recorded from each plot. Net returns, gross returns and benefit cost ratio were computed based on the cost of cultivation with their prevailing market price. Data was statistically analyzed by using Gomez and Gomez [10].

**Comment [IA3]:** .

### 3. RESULTS AND DISCUSSION

#### 3.1. Yield Parameters

According to yield attributes and yield that was collected at harvest, significant and higher number of pods/plant (30.90), number of seeds/pod (1.53), seed yield (2570.74kg/ha) and stover yield (3583.33kg/ha) was observed in treatment 9 [Phosphorus 60kg/ha + PSB] in (Table 2).

Application of phosphorus had significantly influenced the number of pods/plant (30.90), number of seeds/pod (1.53), seed yield (2570.74kg/ha) and stover yield (3583.33kg/ha) in contrast to control [11] [12].

The crop sown with treatments of phosphorus with inoculation of PSB significantly increased the values of yield attributing characters and yields viz., number of pods/plant (30.90), number of seeds/pod (1.53), seed yield (2570.74kg/ha) and stover yield (3583.33kg/ha) of chickpea compared to control [7] [13].

**Comment [IA4]:** Data interpretation should be provided in results and discussions. For example, treatment 6 produced higher yield than treatment 7. Indicating that the 60 kg phosphorus/ha could not be effectively used by the plant without the present of phosphorus solubilizing bacteria, etc.

Table 2. Influence of phosphorus and biofertilizers on yield attributes of chickpea.

| S.No. | Treatments                                      | No. of pods/plant | No. of seeds/pod | Seed yield (kg/ha) | Stover yield (kg/ha) |
|-------|---|-------------------|------------------|--------------------|----------------------|
| 1.    | Phosphorus 40kg/ha + <i>Rhizobium sp.</i>       | 26.27             | 1.13             | 1521.34            | 2606.67              |
| 2.    | Phosphorus 40kg/ha + <i>Pseudomonas striata</i> | 25.90             | 1.13             | 1465.56            | 2570.00              |
| 3.    | Phosphorus 40kg/ha + PSB                        | 28.07             | 1.20             | 1758.56            | 2983.33              |
| 4.    | Phosphorus 50kg/ha + <i>Rhizobium sp.</i>       | 27.30             | 1.20             | 1658.17            | 2783.33              |

|             |   |       |      |         |         |
|-------------|---|-------|------|---------|---------|
| 5.          | Phosphorus 50kg/ha + <i>Pseudomonas striata</i> | 26.87 | 1.13 | 1533.27 | 2746.67 |
| 6.          | Phosphorus 50kg/ha + PSB                        | 28.53 | 1.47 | 2276.39 | 3470.00 |
| 7.          | Phosphorus 60kg/ha + <i>Rhizobium</i> sp.       | 28.10 | 1.33 | 1975.50 | 3256.67 |
| 8.          | Phosphorus 60kg/ha + <i>Pseudomonas striata</i> | 27.70 | 1.20 | 1701.20 | 2836.67 |
| 9.          | Phosphorus 60kg/ha + PSB                        | 30.90 | 1.53 | 2570.74 | 3583.33 |
| 10.         | Control (20-50-20 NPK kg/ha)                    | 25.13 | 1.13 | 1396.27 | 2496.67 |
| F test      |   | S     | S    | S       | S       |
| SEm(±)      |   | 0.93  | 0.05 | 101.85  | 79.42   |
| CD (p=0.05) |   | 2.77  | 0.16 | 302.63  | 235.99  |

### 3.2. Economics.

The total cost of cultivation (40,440.30 INR/ha) was observed to be maximum in treatment 9 (Phosphorus 60kg/ha + PSB) and gross returns (1, 54,110.00 INR/ha), net returns (1, 13,669.70 INR/ha) and B: C ratio (2.81) was also recorded highest in treatment 9 (Table 3).

It is evident from the Table 3 that the highest net returns (1, 13,669.70 INR/ha) and B: C ratio (2.81) were obtained with the application of Phosphorus 60kg/ha. This could be attributed to higher yields of chickpea [3] [14].

Application of phosphorus with inoculation of PSB exhibited excessive values of gross returns (1, 54,110.00 INR/ha), net returns (1, 13,669.70 INR/ha) and B: C ratio (2.81). It also recorded highest growth attributes, yield attributes and yield. [14] [15].

**Table 3. Influence of phosphorus and biofertilizers on yield attributes and yield of chickpea.**

| S.No. | Treatments                                      | Total cost of cultivation (INR) | Gross return (INR/ha) | Net return (INR/ha) | Benefit cost ratio (B:C) |
|-------|---|---------------------------------|-----------------------|---------------------|--------------------------|
| 1.    | Phosphorus 40kg/ha + <i>Rhizobium</i>           | 40,208.30                       | 93,610.00             | 53,401.70           | 1.32                     |
| 2.    | Phosphorus 40kg/ha + <i>Pseudomonas striata</i> | 40,214.30                       | 90,760.00             | 50,545.70           | 1.25                     |
| 3.    | Phosphorus 40 kg/ha + PSB                       | 40,220.30                       | 1,08,180.00           | 67,861.70           | 1.68                     |
| 4.    | Phosphorus 50kg/ha + <i>Rhizobium</i>           | 40,318.30                       | 1,01,880.00           | 61,561.70           | 1.52                     |
| 5.    | Phosphorus 50kg/ha + <i>Pseudomonas striata</i> | 40,324.30                       | 94,840.00             | 54,515.70           | 1.35                     |
| 6.    | Phosphorus 50 kg/ha + PSB                       | 40,330.30                       | 1,34,480.00           | 94,149.70           | 2.33                     |
| 7.    | Phosphorus 60kg/ha + <i>Rhizobium</i>           | 40,428.30                       | 1,21,240.00           | 80,811.70           | 1.99                     |
| 8.    | Phosphorus 60kg/ha + <i>Pseudomonas striata</i> | 40,434.30                       | 1,04,300.00           | 63,865.70           | 1.57                     |

|     |                          |           |             |             |      |
|-----|--------------------------|-----------|-------------|-------------|------|
| 9.  | Phosphorus 60kg/ha + PSB | 40,440.30 | 1,54,110.00 | 1,13,669.70 | 2.81 |
| 10. | Control                  | 39,624.38 | 86,700.00   | 47,075.62   | 1.18 |

#### 4. CONCLUSION

It was concluded that application of Phosphorus 60kg/ha in combination with PSB (Treatment 9) recorded highest seed yield and benefit cost ratio in chickpea crop.

#### REFERENCES

1. R.F. Pegoraro, M.N.D. Almeida Neta, C.A.D. Costa (2018) Chickpea production and soil chemical attributes after phosphorus and molybdenum fertilization. *Ciência e Agrotecnologia*, 42(5):474-483
2. GOI. Economics and statistics of Indian agriculture [annual report]. New Delhi: Government of India; 2020-21.
3. Gulpadiya, V.K, Chhonkar, D.S (2014) Effect of phosphorus on growth, productivity and economics of chickpea varieties. *Annals of Plant and Soil Research* 16(4): 334-337
4. B.J. Prajapati, Nitin Gudadhe, V.R. Gamit and H.J. Chhaganiya (2017) Effect of integrated phosphorus management on growth, yield attributes and yield of chickpea. *Farming and Management* 2(1): 36-40
5. Kennedy IR, Choudhury ATMA, Kecskes ML. (2004) Nonsymbiotic bacterial diazotrophs in crop-farming system: can their potential for plant growth promotion be better exploited? *Soil Biology and Biochemistry*. 2004; 36(8): 1229-1244
6. H. Etesami, B.R. Jeong, B.R. Glick (2021) Contribution of arbuscular mycorrhizal fungi, phosphate-solubilizing bacteria and silicon to P uptake by plant. *Frontiers in Plant Science*, Vol. 12, 699618
7. Shayam Das, B.L. Pareek, Amit Kumawat1 and Shish Ram Dhikwal (2013) Effect of Phosphorus and Bio fertilizers on productivity of chickpea ( *Cicer arietinum* L.) in North Western Rajasthan, India. *Legume Research*, 36 (6): 511 - 514, 2013
8. Jyoti Saxena, Amita Saini, Kirti Kushwaha and Agustin Arino (2016) Synergistic effect of plant growth promoting bacterium *Pseudomonas fluorescens* and phosphate solubilizing fungus *Aspergillus awamori* for growth enhancement of chickpea. *Indian Journal of Biochemistry & Biophysics* Vol.53, June-August 2016, pp 135-143
9. Singh Umesh and Singh Bhagwan (2014) Effect of basal and foliar application of diammonium phosphate in cognizance with phosphate solubilizing bacteria on growth, yield and quality of rain fed chickpea ( *Cicer arietinum* ). *Indian Journal of Agronomy* 59 (3): 427-432
10. K.A. Gomez and A.A. Gomez (1984) Statistical Procedures for Agriculture Research. 2<sup>nd</sup> Edition, John Wiley and Sons, New York, 680 p.

11. S.V.S. Chauhan and Bhoopendra Singh Raghav (2017) Effect of phosphorus and phosphate solubilizing bacteria on growth, yield and quality of chickpea (*Cicer arietinum*). *Annals of Plant and Soil Research* 19(3): 303-306
12. K.K. Pingoliya, A.K Mathur, M.L. Dotaniya, D.K. Jajoria and G.P. Narolia (2014) Effect of phosphorus and iron levels on growth and yield attributes of chickpea (*Cicer arietinum* L.) under agro climatic zone IV A of Rajasthan, India. *Legume Research*, 37 (5): 537-541, 2014.
13. Rajneesh Singh, Tej Pratap, Durgesh Singh, Ghanshyam Singh and Abhinav Kumar Singh (2018) Effect of phosphorus, Sulphur and biofertilizers on growth attributes and yield of chickpea (*Cicer arietinum* L.) *Journal of Pharmacognosy and Phytochemistry* 7(2): 3871-3875
14. Narendra, K. Yadav, R.K. Mahender, S. Dudwe, T.S and Tomar, I.S. (2020) Effect of phosphorus and bioinoculants and their residual effect on succeeding chickpea (*Cicer arietinum* L.) *Indian Journal of Agricultural Sciences* 90(2): 320-325.
15. Yadav, S. Kumar, S. Anshuman, K. Singh, N. and Srivastava, A. (2021) Studies on effect of different biofertilizers on yield and economics of chickpea. *The Pharma Innovation Journal* 10(4):541-545