

Evaluation Of The Antidiabetic Effect Of Aqueous Crude Extract Of Seed, Leaf And Stem Of *L. Usititatissimum* On Streptozotocin-Induced Diabetic Rats

Abstract

Diabetes is a metabolic disorder characterized by persistent high glucose level. *L. usititatissimum* (flaxseed) which is rich in fibre and essential fatty acid is one of the functional foods used in management of diabetes mellitus. The aim of this study is to evaluate the antidiabetic effect of *L. usititatissimum* seed leaf and stem on streptozotocin induced diabetic rats to see which part is more potent in the management of diabetes. Thirty (30) albino rats weighing 170-200g were used for this study and diabetes was induced by intraperitoneal injection of 55mg/kg body weight of streptozotocin. The albino rats were randomly divided into six (6) groups which are as follows: Group A; normal control, B; diabetic control group, C; standard drug (Glibenclamide), D-F were administered different parts of crude plants extract. The phytochemistry analysis showed that the seed possessed more bioactive component followed by the leaf and lastly the stem. The phytochemical found present include: alkaloid, flavonoids, tannins, saponins, terpene, steroid, balsam, carbohydrate and phenol. Across the groups that received crude aqueous extract of *L. usititatissimum* was a significant reduction of blood glucose level, however, the seed (6.60 ± 2.450^{bc}) showed more significant difference when compared to the leaf (7.65 ± 1.100^{bc}) and stem (7.95 ± 2.650^{bc}) respectively. On the other hand, the protein and albumin biomarkers were significantly ($P \leq 0.005$) increased across groups. Crude aqueous extract of *L. usititatissimum* also showed hypolipidemic properties, improved serum enzymes and electrolytes levels. There was also significant positive impact on hematological parameters. In accordance with the results of this investigation, *L. Usititatissimum* seed part is a more potent hypoglycemic and hypolipidemic agent on streptozotocin-induced diabetic rats although, the leaf and stem also demonstrated significant positive parameters on biochemical and haematological parameters. This positive result might be as result of various phytochemical component present in the plant.

Key words: phytochemistry, hypoglycaemic, hypolipidemic, hematology.

1.0 INTRODUCTION

Diabetes mellitus, a non-communicable disease (1), is one of the top five killers in the world (2). This disorder prevents the body from using sugar efficiently (3), which causes an accumulation of too much sugar in the bloodstream and its presence in the urine (4). Furthermore, abnormal lipid metabolism is usually linked to diabetes mellitus (5). Abnormalities in lipid metabolism are frequently seen in both insulin-dependent and non-insulin-dependent diabetes mellitus, and the degree of hyperlipidemia is connected with persistent hyperglycemia (6). In people with pre-diabetes, making small dietary changes and exercising frequently may help postpone or even stop the development of diabetes (7). Omega-3 fatty acids have been associated with enhancements in glycemic and insulin sensitivity (8). Ingesting 1-2 grams of long-chain omega-3 polyunsaturated fatty acids daily can impair glucose intolerance (pre-diabetes) (9;10). Additionally, adding soluble and viscous fiber to the diet can reduce the glucose response to carbohydrate-rich foods by slowing down gastric emptying and glucose absorption (11). Flaxseed contains soluble fiber, α -linolenic acid (ALA), and linoleic acid, which are essential omega-3 fatty acids found in flaxseed and are necessary for normal human growth and development (12).

Considering it has historically been used to treat a variety of medical ailments (13), flaxseed is recognized in Nigeria as a functional food. The objective of this study is to objectively assess the effectiveness of the flax plant in controlling diabetes mellitus. The study precisely examines the flax plant's distinct parts, including the leaves, seeds, and stems, to ascertain which one is most effective in treating diabetes.

The family Linaceae includes flax, also known as *Linum usitatissimum*, which is well-known for its blue blossoms flowers. It is an annual herb that resembles a little palm and has unbranched stems that can grow up to 30 cm tall, though they usually grow shorter. The complex, pinnate leaves are touch-sensitive and close. The flax plant was grown and harvested for this investigation in Nigeria's Plateau State's, Zarmaganda layout. It was subsequently recognized, verified, and given the voucher number JUHN21000351 at the University of Jos' Herbarium Department in Plateau State.



Plate I: Fresh flax plant



Plate II: Dry flax stalk with capsule and flower



Plate III: Flaxseeds

Whole flaxseeds have a lovely nutty flavor and a crispy and chewy texture both as dietary supplements and as ingredients in different meal preparations. It contains Fats, high-quality proteins, and dietary fiber abundantly, flax plant has a sizable amount of the fiber which is water-soluble and viscous in nature (14).

Throughout history, people have gathered a variety of plant resources to satisfy their requirements (15). Numerous items that are used for healing are included in these resources, such as nuts, fruits, vegetables, spices and fibers (16). In contemporary time, many people still depend on gathering plant products for sustenance and as a source of income, especially in underdeveloped nations (17). Traditional medicine continues to be widely used and relied upon, largely because it is more affordable and it is perceived as safer than modern medical procedures. (18). Pharmaceutical companies have developed several medications as a result of the traditional medical system, which uses medicinal herbs. Despite the notable advancements made in

contemporary scientific medicine (19), the use of complementary or alternative medicine has increased significantly on a global scale. This trend is especially apparent in underdeveloped nations, where traditional medicine continues to be the go-to method for treating a variety of illnesses, even among those who have access to contemporary medical procedures (20).

2.0 Materials and method

2.1 Procurement of materials, chemicals and animals

Thirty (30) male albino rats with 170-200 g body weight were purchased from the Animal house in the University of Jos, Nigeria. The rats were allowed free access to standard pellet feed purchased from Grand Cereal and Oil Mills Ltd, Jos Nigeria and water. All chemicals used in this research work were of reagent grade and purchased from the Sigma-Aldrich Company, Germany. Syringes for injections, glucometer to check blood glucose level, and commercial kits to analyze biochemical parameters were purchased from scientific stores and local pharmacies in Jos.

2.2 Preparation of the Plant Extract

After the plant *L. Usitatissimum* (Flaxseed) was collected, the seed, leaf and stem were separated and washed differently to remove the sandy particles and then air dried at room temperature under shade. When it was completely dried, the different parts were pounded to powdery form using local pestle and mortar and sieved to obtain fine powder as described by Hussain et al., (2022) with modifications. The finer powdery form of each part was stored in an air-tight plastic container at ambient conditions until when required. The preparation of the decoction was carried out using warm water at 40°C. 100g of the fine powder of each part was soaked in one (1) litre of distilled warm (40°C) water stirred frequently and allowed to stand for 30 – 60 minutes (to ensure maximum extractions of phytochemicals). The mixture was then filtered using Whatman

filter paper to remove all the larger particles. The filtrate was concentrated in the oven (SM9053A England) at a temperature of 40-45°C for 9 days.

2.3 Determination of Median Lethal Dose (LD₅₀)

To evaluate the hazardous risk of impacts of the Leaf, Seed, and Stem Extract, the Median Lethal Dose was calculated, Using thirteen (13) rats for each extract as described by Chinedu, Arome & Ameh (2013) with slight modification.

The extract of *L. inumusitatissimum* Leaf, Seed, and Stem was given orally to three dosing groups of three rats each during the first part of the experiment, along with one control group that received distilled water. Within four hours of hunger, four hours after feeding, the following twenty-four hours, the following forty-eight hours, and the following two weeks, animals in each group were watched for any immediate indicators of toxicity and mortality.

In the second stage of the acute toxicity test, four rats were split into four groups, and then different higher extract doses—1200, 1600, 2900, and 5000 mg/kg bodyweight—were orally administered based on the results of phase one.

Determination of The Median Lethal Dose (LD₅₀) Of Aqueous Extract of Flax Plant

2.4 Induction of Diabetes and Administration of plant extract

Thirty (30) White albino rats (male Wistar strain) weighing 170-200g were purchased from the animal house of the University of Jos, Nigeria. Diabetes was induced by intraperitoneal injection of streptozotocin at (55mg/kg) to 25 animals in 5 groups, the animals were left for 48hours under controlled conditions after which diabetes was confirmed from the fasting blood glucose using one - touch glucometer. Blood glucose level greater or equal to 160mg/dl, accompanied with hyperglycosuria after 48 hours was considered diabetic. The 5 rats left without induction of diabetes were considered normal

control (23). The plant extract was administered daily at 800mg/kg body weight of each part of the plant for 28 days and the albino rats body weight was obtained weekly using a digital weighing balance.

2.5 Experimental Design

The aqueous extract of *Linum usitatissimum* were administered through oral route. The aqueous extract of seed, leave and stem was administered orally at a dose of 800mg/kg body weight across groups daily. The grouping of the animal are as follows:

Group A_{NC} – Normal control (NC)

Group B_{DC} – Diabetic control (DC)

Group C_{Glib} – Diabetic +800mg/kg bodyweight of Standard drug (Glibenclamide)

Group D_{Seed} – Diabetic + 800mg/kg bodyweight of Seed extract

Group E_{Leaf} – Diabetic + 800mg/kg bodyweight of Leaf extract

Group F_{Stem} – Diabetic + 800mg/kg bodyweight of Stem extract




Fig1: An overview of Study Plan

2.6 COLLECTION OF BLOOD SAMPLE

At the end of the experiment, the rats were starved for 12 hours before they were sacrificed by decapitation and the blood was collected through the jugular vein in plane and heparinise containers for the analysis (24).

2.7 STATISTICAL ANALYSIS

Result values are expressed as mean \pm standard deviation. Analysis of variance (ANOVA) was used for Comparism. Differences were considered significant when values of $p \leq 0.05$. A Graph Pad prism 7.0 in triplicates ($n = 3$) was used to carry out the above analysis

3.0 Discussion

The study evaluated the antidiabetic properties of the aqueous extract and fractions of *L.usititissimum* on STZ induced diabetic rats. According to Bhagarviet al. (2015) and Sharma (2018), diabetes is a disease condition characterized by hyperlipidaemia brought on by the uninhibited actions of lipolytic enzymes and the near absence of insulin. Prolonged hyperlipidaemia leads to serious complications that increase oxidative stress.

3.1 Acute Toxicity

According to the findings of the acute toxicity study, the LD₅₀ of flaxseed leaf, stem, and extract is larger than 5000 mg kg⁻¹ body weight. The limit test dose is typically employed when the experimenter has knowledge that the test substance is probably non-toxic or of low toxicity (27). As a result, the non-lethal effects caused by the large dose of this extracts suggest that acute oral exposure to plant extracts is often harmless. Therefore, it can be said that extracts are safe. Any chemical that has an estimated LD₅₀ value of more than 3000-5000 mg kg⁻¹ (by oral route) may be regarded as being low toxicity and harmless as seen in table 1 Flaxseed is observed to have low acute toxicity (28).

Table 1: LD₅₀ Toxicity Test Result *L. usititissimum*

Group	No of Rats	Mortality Recorded	LD ₅₀
10mg/kg bw	3	0	≥10
100mg/kg bw	3	0	≥100
1000mg/kg bw	3	0	≥1000
1600mg/kg bw	3	0	≥1600

2900mg/kg bw	3	0	≥ 2900
5000mg/kg bw	3	0	≥ 5000

3.2 Phytochemical Assay

Phytochemicals are naturally occurring bioactive substances that can be found in a variety of plant sections, including fruits, vegetables, flowers, leaves, roots of medicinal plants. They provide safeguarding benefits by acting as a protective mechanism against diseases in association with nutrients and fibers (29). Table 2 presents the preliminary phytochemical composition of the aqueous extract obtained from the *L. usitatissimum* plant and its different components. It was observed that the seed part of the plant had more of the phytochemicals followed by the leaf and lastly the stem. The presence of phytochemicals such as alkaloids, terpenes, steroids, carbohydrates, resins, phenols, saponins and flavonoids was identified. The antioxidant capabilities of flavonoids are well known (30), as well as their capacity to alter how the body reacts to dangerous free radicals that are produced during the aberrant metabolic processes frequently seen in diabetes (31). Because of this, flavonoids may have therapeutic advantages (32). Flavonoids have been associated with a number of beneficial attributes on metabolic illnesses, such as cancer, diabetes, obesity, and cardiovascular disease (33). Flavonoids may be helpful in the prevention and treatment of some comorbidities (34). Additionally, they function as antioxidants, reducing the effects of nitrogen and oxygen species on oxidative stress and reducing the risk of disease (35). The ability of flavonoids to control glucose absorption, insulin signaling, insulin secretion, and adipose deposition is related to their anti-diabetic effects (36), to reduce apoptosis and improve hyperglycemia, they focus on a number of molecules that are involved in controlling several pathways, such as those that enhance insulin production, promote β -cell proliferation, and control liver glucose metabolism. This study supports the findings of Luka et al. (2012), who noted that plants with

hypoglycaemic action usually include flavonoids. Furthermore, when digestive enzymes break down dietary polysaccharides, blood glucose levels rises (38), one of these enzymes involved in this process is - amylase, which is present in sizable amounts in saliva and pancreatic juice. Its function is to speed up the dissolution of -1,4-glycosidic bonds found in starch, glycogen, and other oligosaccharides (39). The hydrolase class enzyme α -glucosidase, which is secreted by the epithelial cells lining the brush borders of the small intestine, is another enzyme implicated in this process. Postprandial hyperglycemia is a result of the hydrolytic degradation of oligosaccharides into absorbable monosaccharides, which is aided by α -glucosidase and amylase, and the use of secondary metabolites generated from plants to block these digestive enzymes is a popular strategy for lowering postprandial blood glucose levels (40) and alkaloids have the ability to block these enzymes. Alkaloids have the capacity to attach to either the competitive or noncompetitive sites of the digestive enzymes. This binding stops the enzyme-substrate complex from forming, which eventually lowers enzyme activity (41) Tannins have anti-inflammatory and antioxidant capabilities, tannins, a class of phytochemicals, have demonstrated neuroprotective effects in diabetes complications (42; 43). As a result of saponins' ability to block the enzymes that break down disaccharides into monosaccharides, increased blood glucose levels are reduced (44). In order to reduce high postprandial blood sugar levels, this impact is especially important for managing both Type I and Type II diabetes (45). Saponins also have capacity to precipitate and coagulate red blood cells (RBCs), saponin-containing plants are used in the treatment of wounds (46).

Table 2: Result of the phytochemical screening of the Crude Extracts of leaf, seed and stem of *L. usititatissimum*.

Test	Leaf	Seed	Stem
Alkaloids	-	+	-
Flavonoids	+	+	+

Tannins	+	+	-
Saponins	-	+	-
Terpens/Steroids	-	+	-
Cardiac glycosides	-	-	-
Balsam	+	+	+
Carbohydrates	+	+	+
Phenol	+	+	+
Resins	-	+	-

+ = Detected - = Not Detected

3.3 Biochemical Observations

The findings of this study demonstrate that treatment with *L. usititatissimum* extract at a dosage of 800mg/kg bodyweight significantly ($p \leq 0.05$) reduced blood glucose levels in STZ-induced diabetic rats across all groups when compared to the diabetic group (table 3). The decrease was notably significant compared to both the normal control group and the group treated with the standard drug Glibenclamide. These results align with previous studies (47), suggesting that the plant extract possesses hypoglycemic properties, potentially stimulating insulin release and promoting the uptake of glucose in peripheral tissues, thereby reversing streptozotocin-induced hyperglycemia. These findings indicate that *L. usititatissimum* may be acting through mechanisms such as increasing insulin secretion to enhance glucose uptake by adipose or muscle tissues and inhibiting the intestinal absorption of glucose because of the presence of fibre in the plant (48). It is worthy of note to state that the crude aqueous seed part had more antihyperglycemic effect on the streptozotocin induced diabetic rats even though there was a significant reduction of blood glucose level for the leaf and stem group. Also, Albumin, a major protein

present in human plasma, constitutes approximately 25% of total protein synthesis in the liver and accounts for half of the proteins secreted by the liver. Its production is impeded in various diseases, particularly those affecting the liver (49). Moreover, research indicates that an insulin deficiency leads to reduced synthesis of liver proteins, including albumin and serum proteins (50). Table 3 reveals that the extract treatment group exhibits significantly higher levels of albumin and total serum protein compared to the diabetic control groups. This finding may be attributed to the liver's improved functioning through enhanced metabolism and regulation of protein and albumin, attributable to the beneficial effects of *L. usitatissimum*.

Table 3: Effect of Aqueous Extract of *L. usitatissimum* on Serum Glucose, Serum total Protein and Serum Albumin in STZ induced Diabetic Rats.

Group	Treatment	Glucose (mmol/L)	Total Protein (g/L)	Albumin (g/L)
A	Normal Control	5.85 ± 0.050	74.00±4.400	44.50±1.500
B	Diabetic Control	23.35 ± 4.850 ^b	50.30±0.300 ^a	31.00±3.000 ^a
C	Diabetic + Glibenclamide	5.45 ± 0.450 ^{ac}	72.00±4.00 ^{bc}	43.00 ± 1.000 ^{bd}
D	Diabetic + Extract (Seed)	6.60 ± 2.450 ^{bc}	70.10 ± 3.000 ^{bd}	43.50±0.500 ^{bd}
E	Diabetic + Extract (Leaf)	7.65 ± 1.100 ^{bc}	69.00±1.000 ^{bd}	42.50±3.500 ^{bd}
F	Diabetic + Extract (Stem)	7.95 ± 2.650 ^{bc}	68.50±0.500	40.05±0.500

Values are expressed as mean ± SEM, n=3.

^aValues are significantly lower when compared to normal control (p < 0.05)

^bValues are significantly higher when compared to normal control (p < 0.05)

^cValues significantly lower when compared with diabetic control (p < 0.05)

^dValues significantly higher when compared with diabetic control ($p < 0.05$)

Elevated bilirubin levels may serve as a biomarker for oxidative stress and inflammation in individuals with pre- and new-onset diabetes (51). Studies have demonstrated that serum bilirubin levels are influenced by glucose metabolic status, with individuals with pre-diabetes and new-onset diabetes exhibiting higher bilirubin levels compared to those with normal fasting glucose (52). Consistent with other research findings, table 4 indicates that the aqueous treatment groups and the Glibenclamide treatment group exhibit significantly lower bilirubin levels compared to the diabetic control group. Similar results have been reported by (53).

Table 4: Effect of aqueous extract of *L. usitatissimum* on Serum total bilirubin, direct bilirubin and indirect bilirubin in STZ induced Diabetic rats.

Group	Treatment	TB ($\mu\text{mol/L}$)	DB ($\mu\text{mol/L}$)	IDB ($\mu\text{mol/L}$)
A	Normal Control	9.30 \pm 0.900	2.68 \pm 0.450	6.62 \pm 0.450
B	Diabetic Control	78.05 \pm 11.050 ^b	33.05 \pm 11.041 ^b	45.00 \pm 0.009 ^b
C	D + Glibenclamide	51.20 \pm 27.000 ^{bc}	22.38 \pm 7.600 ^{bc}	28.82 \pm 19.400 ^{bc}
D	D + Extract (Seed)	19.40 \pm 11.000 ^{bc}	6.68 \pm 12.850 ^{bc}	12.72 \pm 23.850 ^{bc}
E	D + Extract (Leaf)	44.25 \pm 23.850 ^{bc}	18.02 \pm 0.089 ^{bc}	26.23 \pm 23.761 ^{bc}
F	D + Extract (Stem)	56.16 \pm 11.950 ^{bc}	25.96 \pm 6.800 ^{bc}	30.20 \pm 5.150 ^{bc}

D= Diabetic, DB= Direct bilirubin, TB= Total bilirubin, IDB= Indirect Bilirubin

Values are expressed as mean \pm SEM, n=3.

^aValues are significantly lower when compared to normal control ($p < 0.05$)

^bValues are significantly higher when compared to normal control ($p < 0.05$)

^cValues are significantly lower when compared to diabetic control ($p < 0.05$)

The body produces cholesterol, which is also derived from some food sources, and it is essential for several physiological processes (54). It contributes to hormone biosynthesis, bile acid synthesis, and cell membrane composition (55). Low-Density Lipoprotein (LDL), sometimes known as "bad" cholesterol, and High-Density Lipoprotein (HDL), also known as "good" cholesterol, are both included in total cholesterol (56). Although cholesterol is essential for overall health, high amounts can be harmful since they can cause arteries to constrict or become blocked (57). A significant rise in the risk of cardiovascular disease (CVD) is unfortunately seen in those with diabetes who are more likely to have unhealthily high cholesterol levels (58). By effectively managing cholesterol levels, individuals can lower their chances of developing CVD and premature death (59). Table 5 of this study demonstrates a considerable rise in HDL levels and, on the other hand, a significant fall in LDL levels. The *L. usititatissimum* aqueous extract and the control rats were equivalent. The fact that *L. usititatissimum* was able to considerably raise HDL levels and lower LDL levels in streptozotocin-induced diabetic rats suggests that it has hypolipidemic capabilities, in addition, research have shown that linoleic acid can improve insulin sensitive (60). The most common type of fat in the body, triglycerides, have normal levels that vary depending on age and gender. Atherosclerosis, or the buildup of fatty deposits within artery walls, is associated with elevated triglyceride levels, low HDL cholesterol, or high LDL cholesterol (61). Having high triglyceride levels increases the risk of heart attack, peripheral artery disease (PAD), and stroke. It is worth mentioning that elevated triglycerides do not directly cause diabetes. Rather, they indicate an impairment in the body's ability to convert food into energy (62).

Table 5: Effect of Aqueous extract of *L. usititatissimum* on Lipid profile in STZ induced Diabetic rats.

Group	Treatment	Total Cholesterol (mmol/L)	Triglyceride (mmol/L)	HDL (mmol/L)	LDL (mmol/L)
A	Normal Control	2.30 ± 0.100	0.95 ± 0.050	1.08 ± 0.015	0.78±0.105
B	Diabetic Control	4.80 ± 0.400 ^b	1.90 ± 0.500 ^b	1.00 ± 0.335 ^a	2.81±0.190 ^b
C	D + Glibenclamide	3.25 ± 0.450 ^{bc}	1.70 ± 0.100 ^b	2.16 ± 0.165 ^b	1.73±0.065 ^{bc}
D	D + Extract (Seed)	3.25 ± 0.450 ^b	1.35 ± 0.050 ^{bc}	0.80 ± 0.115 ^{ac}	1.76±0.045 ^{bc}
E	D + Extract (Leaf)	3.10 ± 0.300 ^{bc}	1.35 ± 0.050 ^{bc}	1.03 ± 1.185 ^{be}	1.48±0.160 ^{bc}
F	D + Extract (Stem)	3.55 ± 0.550 ^{bd}	0.90 ± 0.100 ^{ac}	1.00 ± 0.150 ^{ae}	1.85 ± 0.525 ^{bc}

HDL - High Density Lipoprotein, LDL – Low Density Lipoprotein

Values are expressed as mean ± SEM, n=3

^aValues are significantly lower when compared to normal control (p < 0.05)

^bValues are significantly higher when compared to normal control (p < 0.05)

^cValues are significantly lower when compared to diabetic control (p < 0.05)

^dValues are significantly higher when compared to diabetic control (p < 0.05)

^eValues are almost equal to compared to normal control (p > 0.05)

Individuals with type 2 diabetes mellitus (T2DM) have a higher likelihood of experiencing abnormalities in liver function tests compared to those without T2DM (63). Supporting this, Mandal et al. (2018) reported elevated levels of alanine aminotransferase (ALT), aspartate aminotransferase (AST), and alkaline phosphatase (ALP) in a diabetic population. Table 6 illustrates that the serum levels of ALT, AST, and ALP were significantly reduced (p ≤ 0.05) in groups treated with the aqueous extract of *L. usitatissimum* when compared to the diabetic control group. Similarly, decreased levels of ALT, AST, and ALP were observed in diabetic rats treated with Glibenclamide. The release of these enzymes into the

bloodstream is indicative of the hepatotoxic effect of diabetes (65). Therefore, the *L. usitatissimum* aqueous extract significantly lowered the levels of ALT, AST, and ALP in the serum, suggesting its hepatoprotective properties.

Table 6: Effect of aqueous extract of *L. usitatissimum* on some Serum Enzymes in STZ induced Diabetic rats.

Group	Treatment	ALT(U/L)	AST (U/L)	ALP(U/L)
A	Normal Control	68.50 ± 4.500	56.50 ± 3.500	95.50 ± 2.000
B	Diabetic Control	258.83 ± 95.500 ^b	202.50±70.500 ^b	300.50 ± 80.500 ^b
C	D + Glibenclamide	125.00 ± 23.000 ^{bc}	99.50 ± 21.500 ^{bc}	149.00 ± 19.000 ^{bc}
D	D + Extract (Seed)	113.00 ± 10.000 ^{bc}	87.50±17.500 ^{bc}	140.50 ± 17.500 ^{bc}
E	D + Extract (Leaf)	181.50 ± 86.500 ^{bc}	128.50±58.500 ^{bc}	196.50 ± 92.500 ^{bc}
F	D +Extract (Stem)	162.00 ± 54.500 ^{bc}	116.50±45.500 ^{bc}	184.00 ± 49.000 ^{bc}

AST – Aspartate transaminase, ALT – Alanine Aminotransferase, ALP – Alkaline Phosphatase

Values are expressed as mean ± SEM, n=3

^aValues are significantly lower when compared to normal control (p < 0.05)

^bValues are significantly higher when compared to normal control (p < 0.05)

^cValues are significantly lower when compared to diabetic control (p < 0.05)

^dValues are significantly higher when compared to diabetic control (p < 0.05)

In diabetes, markers such as urea, uric acid, and creatinine tend to increase due to kidney impairment, which can be caused by diabetes. This can result in the leakage of these biomarkers into the urine (66). Urea which is a by-product of protein metabolism and is typically eliminated by the kidneys. In diabetic

rats, the levels of urea, creatinine, and uric acid were significantly higher compared to normal rats, suggesting that diabetes can lead to kidney dysfunction. However, when diabetic rats were treated with different parts of an aqueous extract from *L. usititatissimum*, there was a significant reduction in the levels of urea, creatinine, and uric acid compared to untreated diabetic rats. This suggests that the extract has a protective effect on the kidneys. These findings are consistent with the studies conducted by Dinko and Edward (2015) and Dirk et al. (2015). Diabetes mellitus profoundly affects the kidneys and is considered the main culprit behind diabetic nephropathy, (69). In addition to oxidative stress and the presence of advanced glycation end-products, disrupted lipid metabolism and the build up of lipids in the kidneys have been proposed as factors that contribute to the development of diabetic nephropathy. Several research studies have reported the observation of lipid accumulations in the kidneys of both diabetic patients and animal models, suggesting that these deposits may play a significant role in the progression of diabetic kidney disease (70).

Table 7: Effect of aqueous extract of *L. usititatissimum* on serum urea, serum creatinine and serum uric acid concentrations in STZ induced Diabetic rats.

Group	Treatment	Urea (mmol/L)	Creatinine (μmol/L)	Uric Acid (μmol/L)
A	Normal Control	312.50 \pm 54.500	13.50 \pm 2.000	1.00 \pm 0.200
B	Diabetic Control	548.00 \pm 60.000 ^b	42.00 \pm 9.000 ^a	3.20 \pm 0.400 ^b
C	D + Glibenclamide	446.50 \pm 31.500 ^{bc}	33.00 \pm 8.000 ^{ab}	2.60 \pm 0.600 ^{bd}
D	D + Extract (Seed)	375.50 \pm 3.500 ^{bc}	20.50 \pm 4.500 ^{bc}	1.80 \pm 0.400 ^{bc}
E	D + Extract (Leaf)	436.50 \pm 17.500 ^{bc}	28.50 \pm 2.500 ^{bc}	2.55 \pm 0.250 ^{bd}

F D + Extract (Stem) 505.00±101.500^{bd} 27.00 ± 1.00^{bc} 2.55 ± 0.500^{bd}

Values are expressed as mean ± SEM, n=3

^aValues are significantly lower when compared to normal control (p < 0.05)

^bValues are significantly higher when compared to normal control (p < 0.05)

^cValues are significantly lower when compared to diabetic control (p < 0.05)

^dValues are significantly higher when compared to diabetic control (p < 0.05)

The increased volume and metabolites excretion via the kidneys, usually in excess of normal thresholds give rise to imbalance in homeostasis with respect to electrolytes (71). Usually measured electrolytes are Na⁺, K⁺, Cl⁻ and HCO₃⁻. These four ions in plasma exert the greatest influence on water balance and acid-base relationship (72). From table 8, when compared to the normal and diabetic control, Treatment with *L. usititatissimum* aqueous extract showed that there was a significant decrease in serum concentration of Na⁺, Cl⁻ and HCO₃⁻. The analysis plant extracts administration has impact on the electrolytes and exhibited properties capable of boosting the buffering function of the body system (73).

Table 8: Effect of aqueous extract of *L. usititatissimum* on some serum electrolytes Concentrations in STZ induced Diabetic rats.

Group	Treatment	Na ⁺ (mmol/L)	K ⁺ (mmol/L)	Cl ⁻ (mmol/L)	HCO ₃ ⁻ (mmol/L)
A	Normal Control	148.00 ± 0.00	4.50 ± 0.250	103.50 ± 0.159	28.00 ± 1.000
B	Diabetic Control	145.00 ± 1.000 ^a	4.85 ± 0.100 ^e	104.50 ± 1.500 ^b	25.00 ± 1.000 ^a
C	D + Glibenclamide	134.27 ± 2.000 ^{ac}	5.40 ± 0.500 ^{bd}	96.00 ± 4.000 ^{ac}	23.00 ± 2.000 ^{ac}
D	D + Extract (Seed)	134.00 ± 0.000 ^{ac}	4.90 ± 0.200 ^{ac}	98.50 ± 0.500 ^{ac}	24.50 ± 1.500 ^{ac}

E	D + Extract (Leaf)	136.00 ± 4.000 ^{ac}	5.00 ± 0.200 ^{bd}	99.00 ± 3.000 ^{ac}	25.50 ± 1.500 ^{be}
F	D + Extract (Stem)	137.00 ± 7.000 ^{ac}	5.55 ± 0.350 ^{bd}	100.50 ± 6.500 ^{ac}	22.00 ± 1.000 ^{ac}

Values are expressed as mean ± SEM, n=3

^aValues are significantly lower when compared to normal control (p < 0.05)

^bValues are significantly higher when compared to normal control (p < 0.05)

^cValues are significantly lower when compared to diabetic control (p < 0.05)

^dValues are significantly higher when compared to diabetic control (p < 0.05)

^eValues are significantly almost equal when compared to normal control (p < 0.05)

3.4 Analysis of haematological studies

Assessing the hematological parameters of Red Blood Cells (RBCs) and White Blood Cells (WBCs) can provide valuable information about the adverse effects on the blood components in animals, (74). Additionally, these parameters can help elucidate the blood-related functions of chemical compounds found in plant extracts (75). In individuals with diabetes, various factors like advanced glycation end products (AGEs), oxidative stress, angiotensin II, and cytokines can stimulate WBCs. This activation of WBCs in diabetic individuals leads to the release of cytokines and transcription factors that significantly contribute to inflammation, (76). When evaluating the effects of *L. usitatissimum* aqueous extract derived from the leaf, stem, and seed (as presented in Table 9), a noteworthy reduction in WBC count was observed across the experimental groups.

Table 9: Effects of Aqueous extracts of *L. usitatissimum* on White blood count and its parameters Concentrations in STZ induced Diabetic rats.

Treatment	WBC	NEU(%)	LYM(%)	MON(%)	EOS(%)	BAS(%)
-----------	-----	--------	--------	--------	--------	--------

A	Normal Control	6.35±0.158	10.86±0.318	65.36±3.351	0.53±0.088	0.36±0.033	0.00±0.000
B	Diabetic Control	19.99±1.669 ^{bd}	65.70±13.884 ^b	45.93±14.856 ^a	6.13±0.348 ^a	2.73±0.202 ^b	1.36±0.441 ^b
C	D + Glib	7.03±2.116 ^{ac}	30.63±1.746 ^{bc}	60.90±0.907 ^{bd}	3.13±0.866 ^{ad}	0.53±0.120	0.76±0.202
D	D + Extract (Seed)	7.51±1.370 ^b	37.76±1.281 ^{bc}	61.23±1.501 ^{bd}	0.63±0.088 ^{ac}	0.56±0.033 ^{bc}	0.52±0.000 ^{ac}
E	D + Extract (Leaf)	12.88±0.904 ^{bd}	51.26±1.648 ^{bc}	30.93±1.411 ^{ac}	1.10±0.057 ^{ac}	1.30±0.321 ^{bc}	1.31±0.600 ^{bd}
F	D + Extract (Stem)	13.94±1.280 ^a	49.20±0.550 ^{bc}	48.83±0.977 ^{ad}	2.73±2.783	0.93±0.371 ^{bc}	1.10±0.057 ^{bd}

WBC- White blood cells, NEU- Neutrophil, LYM- Lymphocyte, MON- Monocytes, EOS- Eosinophil, BAS- Basophile

Values are expressed as mean ± SEM, n = 3.

^aValues are significantly lower when compared with normal control (p < 0.05)

^bValues are significantly higher when compared with normal control (p < 0.05)

^cValues are significantly lower when compared with diabetic control (p < 0.05)

^dValues are significantly higher when compared with diabetic control (p < 0.05)

Erythrocytes, being a crucial element of the bloodstream, serve as a highly responsive indicator of an individual's overall health condition (77). The dangers that diabetic patients face include hyperglycemia, hyperosmolarity, oxidative stress, inflammation, and problems with lipid metabolism. These hazards can all have an impact on their red blood cells (erythrocytes). These elements influence the aggregation, deformability, and fluidity of the erythrocyte membrane. In the end, these modifications disturb microcirculation and aid in the emergence of diabetic problems (78) Table 10 shows that administration of the extracts had a positive change in the haematological parameters suggesting that it may have contributed to the ameliorative effect that is observed from the treatment group.

Table 10: Effects of Aqueous extracts of *L. usititatissimum* on Red Blood Count and its Parameters Concentrations in STZ induced Diabetic rats.

Group	Treatment	RBC	HGB	HCT(PCV)	PLT	PCT
A	Normal Control	8.01±0.500 ^{bd}	15.78±1.019 ^{bd}	47.50±0.458	279.67±7.446 ^{ac}	0.21±0.014
B	Diabetic Control	3.48±0.274 ^b	6.70±0.550 ^b	39.83±1.328 ^b	623.67±110.900 ^b	0.45±0.087 ^e
C	D + Glibenclamide	7.45±0.183 ^{bc}	14.80±0.450 ^{bd}	45.93±0.845 ^{bc}	306.00±5.568 ^{ac}	0.23±0.015 ^{ac}
D	D + Extract (Seed)	7.25±0.119 ^{bc}	13.83±0.120 ^{bc}	44.93±0.338 ^{bc}	373.67±41.289 ^{ac}	0.35±0.040 ^{ac}
E	D + Extract (Leaf)	6.96±0.136	12.10±0.152	43.60±2.635 ^{bd}	549.67±1.453 ^{ac}	0.40±0.045 ^{ac}
F	D + Extract (Stem)	7.03±0.063 ^{bc}	13.53±0.202 ^{bc}	41.50±0.556 ^{cc}	511.33±14.847	0.42±0.036 ^{ac}

RBC-Red Blood Cells, HGB-Haemoglobin, HCT-Haematocrit, PLT-Platelets and PCT-Plateletcrit

Values are expressed as mean ± SEM, n = 3.

^aValues are significantly lower when compared with normal control (p < 0.05)

^bValues are significantly higher when compared with normal control (p < 0.05)

^cValues are significantly lower when compared with diabetic control (p < 0.05)

^dValues are significantly higher when compared with diabetic control (p < 0.05)

^eValue is equal to normal control (p < 0.05)

3.5 Conclusion

On the basis of the current research carried out, it can conclude that the plant *L. usititatissimum* possesses hypoglycaemic, hypolipidemic activity, hypocholesterolaemia and hepatoprotective effect in diabetic rats. The traditional use of the plant to manage diabetes is supported by this laboratory finding. The progressive decrease in the blood glucose levels of rats and improvement

of biochemical parameters shown by the treatment with the aqueous extract of *Linum usitatissimum* may be due to the presence of phytochemicals found in abundance in the plant.

References

1. Maiti R., Jana D., Das UK., Ghosh D., Antidiabetic effect of aqueous extract of seed of *Tamarindus indica* in streptozocin-induced diabetic rats. *J.Ethnopharmacology*, (2004). 92: 85 – 91.
2. American Diabetes Association. Standards of medical care in diabetes—2012. *Diabetes care*, 35(Supplement 1), S11-S63.
3. Agarwal, M. M., Dhatt, G. S., Zayed, R., & Bali, N. Gestational diabetes: relevance of diagnostic criteria and preventive strategies for Type 2 diabetes mellitus. *Archives of gynecology and obstetrics*, (2007). 276(3), 237-243.
4. American Diabetes Association. Standards of medical care in diabetes. *Diabetes (2014) Care*;37(Suppl. 1):S14–S80
5. Hussain, A., Kausar, T., Jamil, M. A., Noreen, S., Iftikhar, K., Rafique, A., ... & Ali, A. In Vitro role of pumpkin parts as pharma-foods: antihyperglycemic and antihyperlipidemic activities of pumpkin peel, flesh, and seed powders, in alloxan-induced diabetic rats. *International Journal of Food Science*, 2022.
6. Giugliano, D., Ceriello, A., & Esposito, K. Glucose metabolism and hyperglycemia. *The American Journal of Clinical Nutrition*, (2008). 87(1), 217S-222S. <https://doi.org/10.1093/ajcn/87.1.217S>
7. Hutchins, A. M., Brown, B. D., Cunnane, S. C., Domitrovich, S. G., Adams, E. R., & Bobowiec, C. E. Daily flaxseed consumption improves glycemic control in obese men and women with pre-diabetes: A randomized study. *Nutrition Research*, (2013). 33(5), 367-375. <https://doi.org/10.1016/j.nutres.2013.02.012>
8. Parikh, M.; Netticadan, T.; Pierce, G.N. Flaxseed: Its bioactive components and their cardiovascular benefits. *Am. J. Physiol. Heart Circ. Physiol.* **2018**, 314, H146–H159. [\[Google Scholar\]](#) [\[CrossRef\]](#) [\[PubMed\]](#)

9. Parikh, M., Maddaford, T. G., Austria, J. A., Aliani, M., Netticadan, T., & Pierce, G. N. Dietary Flaxseed as a Strategy for Improving Human Health. *Nutrients*, (2019). *11*(5), 1171. <https://doi.org/10.3390/nu11051171>
10. Austria, J.A.; Aliani, M.; Malcolmson, L.J.; Dibrov, E.; Blackwood, E.P.; Maddaford, T.G.; Guzman, R.; Pierce, G.N. Daily food choices over one year when patient diets are supplemented with milled flaxseed. *J. Funct. Foods* **2016**, *26*, 772–780. [[Google Scholar](#)] [[CrossRef](#)]
11. Dodin, S.; Lemay, A.; Jacques, H.; Légaré, F.; Forest, J.C.; Mâsse, B. The effects of flaxseed dietary supplement on lipid profile, bone mineral density, and symptoms in menopausal women; a randomized, double-blind, wheat germ placebo-controlled clinical trial. *J. Clin. Endocrinol. Metab.* **2005**, *90*, 1390–1397. [[Google Scholar](#)] [[CrossRef](#)]
12. Rodriguez-Leyva, D.; Weighell, W.; Edel, A.L.; La Vallee, R.; Dibrov, E.; Pinneker, R.; Maddaford, T.G.; Ramjiawan, B.; Aliani, M.; Guzman, R.; et al. Potent anti-hypertensive action of dietary flaxseed in hypertensive patients. *Hypertension* **2013**, *62*, 1081–1089. [[Google Scholar](#)] [[CrossRef](#)]
13. Edel, A.E.; Aliani, M.; Pierce, G.N. Stability of bioactives in flaxseed and flaxseed-fortified foods. *Food Res. Int.* **2015**, *77*, 140–155. [[Google Scholar](#)] [[CrossRef](#)]
14. Aliani, M.; Ryland, D.; Pierce, G.N. Effect of flax addition on the flavor profile of muffins and snack bars. *Food Res. Int.* **2011**, *44*, 2489–2496. [[Google Scholar](#)] [[CrossRef](#)]
15. Egua, M. O.. Pharmacognostic Study of a Plant Seed Extract. In *Pharmacognosy-Medicinal Plants*. (2019) IntechOpen.
16. Okwu, D., and Josiah, C. Evaluation of the chemical composition of two Nigerian medicinal plants. *Afr. J. Biotechnol.* 2006. *5*(4), 357-361.
17. Shobowale, O. O., Ogbulie, N. J., Itoandon, E. E., Oregun, M. O., & Olatope, S. O. A. Phytochemical and antimicrobial evaluation of aqueous and organic extracts of *Calotropis procera* ait leaf and latex. *Nigerian Food Journal*, (2013). *31*(1), 77-82.

18. Sakthivel, K. M., & Guruvayoorappan, C. Biophytumsensitivum: Ancient medicine, modern targets. *Journal of advanced pharmaceutical technology & research*, (2012). 3(2), 83.
19. Atawodi, S. E., Ameh, D. A., Ibrahim, S., Andrew, J. N., Nzelibe, H. C., Onyike, E. O., ... & Sallau, A. B. Indigenous knowledge system for treatment of trypanosomiasis in Kaduna state of Nigeria. *Journal of Ethnopharmacology*, (2002). 79(2), 279-282.
20. Amin, T., & Thakur, M. A comparative study on proximate composition, phytochemical screening, antioxidant and antimicrobial activities of *Linum usitatissimum* L.(flaxseeds). *Int. J. Curr. Microbiol. App. Sci*, (2014). 3(4), 465-481.
21. Hussain, A., Kausar, T., Jamil, M. A., Noreen, S., Iftikhar, K., Rafique, A., ... & Ali, A. In Vitro role of pumpkin parts as pharma-foods: antihyperglycemic and antihyperlipidemic activities of pumpkin peel, flesh, and seed powders, in alloxan-induced diabetic rats. *International Journal of Food Science*, 2022.
22. Chinedu, E., Arome, D., & Ameh, F. S. A New Method for Determining Acute Toxicity in Animal Models. *Toxicology International*, (2013). 20(3), 224-226. <https://doi.org/10.4103/0971-6580.121674>
23. Rafiu A. A and Luka C. D Evaluation of the antidiabetic property of aqueous extract of ipomoea batata leaf on hyperglycemia, hyperlipidemia, blood electrolytes and enzymatic antioxidants of streptozotocin induced diabetic rats. *J Res Diabetes Metab*. (2018). 4(1): 021-026.
24. Clarkson, J. M., Martin, J. E., & F McKeegan, D. E. A review of methods used to kill laboratory rodents: Issues and opportunities. *Laboratory Animals*, (2022). 56(5), 419-436. <https://doi.org/10.1177/00236772221097472>
25. Bhargavi P., Josthna C., Naidu V., changes in serum Biochemical Parameters and lipid profile in Normal and STZ induced diabetic rats with the administration of ethanolic extract of *Polyalthlacerasoides* Stem Bark. *Int. Res. J. Pharm*. (2015), 6(2).
26. Sharma, S. Diabetes Mellitus: An Overview. *Journal of Advanced Scientific Research*, (2018). 9, 1.
27. OECD, 2001. Guidelines for testing chemical, Acute Oral toxicities up and down produce. OECD Reports 425, pp:1 - 26.

28. Nwogo Ajuka Obasi, Kalu Mong Kalu, Uchechukwu Okorie and Glory Otuchristian, Hypoglycemic effects of Aqueous and Methanolic Leaf Extracts of *Vitex doniana* on Alloxan Induced Diabetic Albino Rats. *Journal of Medical Sciences*, 2013. 13: 700-707. DOI: [10.3923/jms.2013.700.707](https://doi.org/10.3923/jms.2013.700.707)
29. Krishna, D.; Nagaraja Kumari, K.; Nagarjuna, B. V. R., Effect of enzymes and liver tonic supplementation on performance of quails fed processed karanj (*Pongamia glabra* Vent) cake. *Int. J. Agric. Sci. Vet. Med.*, 2013. 1 (3): 39-46
30. Middleton E., Kandaswami C., Jr., Theoharides T.C. The effects of plant flavonoids on mammalian cells: Implications for inflammation, heart disease, and cancer. *Pharmacol. Rev.* 2000;52:673–751. [[PubMed](#)] [[Google Scholar](#)]
31. Hossain M.K., Choi H.Y., Hwang J.S., Dayem A.A., Kim J.H., Kim Y.B., Cho S.G. Antiviral activity of 3,4'-dihydroxyflavone on influenza a virus. *J. Microbiol.* 2014;52:521–526. doi: 10.1007/s12275-014-4212-z. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
32. Vinayagam R., Xu B. Antidiabetic properties of dietary flavonoids: A cellular mechanism review. *Nutr. Metab. (Lond.)* 2015;12:60. doi: 10.1186/s12986-015-0057-7. [[PMC free article](#)] [[PubMed](#)]
33. Graf B.A., Milbury P.E., Blumberg J.B. Flavonols, flavones, flavanones, and human health: Epidemiological evidence. *J. Med. Food.* 2005;8:281–290. doi: 10.1089/jmf.2005.8.281. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
34. AL-Ishaq, R. K., Abotaleb, M., Kubatka, P., Kajo, K., & Büsselberg, D. Flavonoids and Their Anti-Diabetic Effects: Cellular Mechanisms and Effects to Improve Blood Sugar Levels. (2019) *Biomolecules*, 9(9). <https://doi.org/10.3390/biom9090430>
35. Edeoga H. O., Okwu d. E., Mbaebie B. O. Phytochemical constituents of some Nigerian medicina plants. *African Journal of Biotechnology* (2005). vol. 4 (7), PP 685-688
36. Hussain, A., Kausar, T., Sehar, S., Sarwar, A., Quddoos, M. Y., Aslam, J., ... & Nisar, R. A review on biochemical constituents of pumpkin and their role as pharma foods; a key strategy to improve health in post COVID 19 period. *Food Production, Processing and Nutrition*, (2023). 5(1), 1-14.

37. Luka C.D, Istifanus G, George M, Philip CJ., The effect of aqueous extract of citrus sinensis peel on some Biochemical parameters in Normal and Alloxan- induced Diabetic Wister Rats. *Am. J.ofphytomed. Clin. Ther.* (2017). Vol. 5No. 2:17.
38. M. S. Deuschländer, N. Lall, M. Van de Venter, and A. A. Hussein, "Hypoglycemic evaluation of a new triterpene and other compounds isolated from *Euclea undulata* Thunb. var. *myrtina* (Ebenaceae) root bark," *Journal of Ethnopharmacology*, vol. 133, no. 3, pp. 1091–1095, 2011. View at: [Publisher Site](#) | [Google Scholar](#)
39. Z. Liu, Y. Lv, Y. Zhang et al., "Matrine□type Alkaloids inhibit advanced glycation end products induced reactive oxygen species□mediated apoptosis of aortic endothelial cells in vivo and in vitro by targeting MKK3 and p38MAPK signaling," *Journal of the American Heart Association*, vol. 6, no. 12, Article ID e007441, 2017. View at: [Publisher Site](#) | [Google Scholar](#)
40. Bikash Adhikari, "Roles of Alkaloids from Medicinal Plants in the Management of Diabetes Mellitus", *Journal of Chemistry*, vol. 2021, Article ID 2691525, 10 pages, 2021. <https://doi.org/10.1155/2021/2691525>
41. S. Tiong, C. Looi, H. Hazni et al., "Antidiabetic and antioxidant properties of alkaloids from *Catharanthus roseus* (L.) G. Don," *Molecules*, vol. 18, no. 8, pp. 9770–9784, 2013. View at: [Publisher Site](#) | [Google Scholar](#)
42. Laddha, A. P., & Kulkarni, Y. A.). Tannins and vascular complications of Diabetes: An update. *Phytomedicine*, (2019) 56, 229-245. <https://doi.org/10.1016/j.phymed.2018.10.026>
43. Omar, N., Nazariah Ismail, C. A., & Long, I. Tannins in the Treatment of Diabetic Neuropathic Pain: Research Progress and Future Challenges. *Frontiers in Pharmacology*, (2021). 12. <https://doi.org/10.3389/fphar.2021.805854>
44. Oishi Y, Sakamoto T, Udagawa H *et al.* Inhibition of increases in blood glucose and serum neutral fat by *Momordica charantia* saponin fraction. *Biosci. Biotechnol. Biochem.* 71(3), 735–740 (2007).
45. Sood, A., Kaur, P., and Gupta, R. Phytochemical screening and antimicrobial assay of various seeds extract of Cucurbitaceae family. *Int. J. Appl. Biol. Pharmac. Technol.* 2012. 3(3), 401-409. S

46. Okwu, D., and Josiah, C. Evaluation of the chemical composition of two Nigerian medicinal plants. *Afr. J. Biotechnol.* 2006 5(4), 357-361. O
47. Okonkwo, P.O. and Okoye, Z.S. C. Hypoglycaemic Effects of the Aqueous Extract of Newbouldia Leaves Root in Rats, *Int. J. Biol. Chem.* (2009). 12: 42-48.
48. Luka C.D, Istifanus G, George M, Philip CJ,. The effect of aqueous extract of citrus sinensis peel on some Biochemical parameters in Normal and Alloxan- induced Diabetic Wister Rats. *Am. J.ofphhytomed. Clin. Ther.* (2017) Vol. 5No. 2:17.
49. Murray, R.K., D.K Granner, P.A. Mayes and V.W. Rodwell, Harper's Biochemistry. 25th Edn., Appleton and Lange, Philadelphia, (2000). pp: 179-203, 621-630
50. Jefferson LS, Liao WS, Peavy DE, Miller TB, Appel MC, Taylor JM. Diabetes-induced alterations in liver protein synthesis. Changes in the relative abundance of mRNAs for albumin and other plasma proteins. *J Biol Chem.* 1983 Jan 25;258(2):1369-75. PMID: 6822503.
51. Omonkhua A.A., Adebayo E.A., Saliu J.A., Ogunwa T.H., Adeyelu T.T.. Liver function of Streptozotocin- Induced Diabetic Rats Orally Administered Aqueous Root-BarkExtracts of *Tetrapleura tetraptera* (Taub). *Nigerian Journal of Basic and Applied Science* (Sep. – Dec., 2014), 22(3&4): 99-106
52. Wang J, Li Y, Han X, Hu H, Wang F, Li X, Yang K, Yuan J, Yao P, Miao X, Wei S, Wang Y, Cheng W, Liang Y, Zhang X, Guo H, Yang H, Yuan J, Koh WP, Hu FB, Wu T, Pan A, He M. Serum bilirubin levels and risk of type 2 diabetes: results from two independent cohorts in middle-aged and elderly Chinese. *Sci Rep.* 2017 Feb 6;7:41338. doi: 10.1038/srep41338. PMID: 28164994; PMCID: PMC5292699.
53. Luka, C.D. A biochemical Study of the effect of extract of goat's rue (*Galega officinalis*) in alloxan-induced diabetic rats. (2013). pp 158-159
54. Omonkhua, A. A., Onoagbe, I. O., Fajimeye, I. A., Adekola, M. B., &Imoru, Z. A. Long-term anti-diabetic and anti-hyperlipidaemic effects of aqueous stem bark extract of *Irvingiagabonensis* in streptozotocin-induced diabetic rats. *Biokemistri*, (2014). 26(1), 1-8.
55. Renuka, N., Sood, A., Prasanna, R., & Ahluwalia, A. S. Phycoremediation of wastewaters: a synergistic approach using microalgae for bioremediation and biomass

- generation. *International Journal of Environmental Science and Technology*, (2015). 12(4), 1443-1460.
56. Saxena, S., & Katare, C. Evaluation of flaxseed formulation as a potential therapeutic agent in mitigation of dyslipidemia. *Biomed J*, (2014). 37(6), 386-90.
57. Risérus, U., Willett, W. C., & Hu, F. B. Dietary fats and prevention of type 2 diabetes. *Progress in lipid research*, (2009). 48(1), 44-51.
58. M. Makni, H. Fetoui, N. K. Gargouri, E. M. Garoui, and N. Zeghal, "Antidiabetic effect of flax and pumpkin seed mixture powder: effect on hyperlipidemia and antioxidant status in alloxan diabetic rats," *Journal of Diabetes and its Complications*, vol. 25, no. 5, article S1056872710000978, pp. 339–345, 2011.
59. Alasalvar C, Amaral J. S., & Shahidi F. Functional lipid characteristics of Turkish Tombul hazelnut (*Corylus avellana* L.). *J Agric Food Chem* (2006). 54, 10177–10183
60. Alvarez, F., Bustos, M., Villanueva, E., Ramírez, F., León, M., Duggirala, R., Macías, T., Cuevas, S., Jalife, A., Serratos, F., & Carlos, J. (2017). Effects of Conjugated Linoleic Acid and Metformin on Insulin Sensitivity in Obese Children: Randomized Clinical Trial. *The Journal of Clinical Endocrinology & Metabolism*, 102(1), 132-140. <https://doi.org/10.1210/jc.2016-2701>
61. Parhofer, K. G. Interaction between Glucose and Lipid Metabolism: More than Diabetic Dyslipidemia. *Diabetes & Metabolism Journal*, (2015). 39(5), 353-362. <https://doi.org/10.4093/dmj.2015.39.5.353>
62. Massing, M. W., Sueta, C. A., Chowdhury, M., Biggs, D. P., & Simpson, R. J. Lipid management among coronary artery disease patients with diabetes mellitus or advanced age. *American Journal of Cardiology*, (2001). 87(5), 646-649.
63. Mathur S, Mehta DK, Kapoor S, Yadav S. Liver function in type-2 diabetes mellitus patients. *Int J Sci Stud*. 2016;3:43 [[Google Scholar](#)]
64. Mandal A, Bhattarai B, Kafle P, Khalid M, Jonnadula SK, Lamicchane J, Kanth R, Gayam V. Elevated Liver Enzymes in Patients with Type 2 Diabetes Mellitus and Non-alcoholic Fatty Liver Disease. *Cureus*. 2018 Nov 23;10(11):e3626. doi: 10.7759/cureus.3626. PMID: 30697502; PMCID: PMC6347442.
65. Dodin, S.; Lemay, A.; Jacques, H.; Légaré, F.; Forest, J.C.; Mâsse, B. The effects of flaxseed dietary supplement on lipid profile, bone mineral density, and symptoms in

menopausal women; a randomized, double-blind, wheat germ placebo-controlled clinical trial. *J. Clin. Endocrinol. Metab.* **2005**, *90*, 1390–1397. [[Google Scholar](#)] [[CrossRef](#)]

66. Dinko S. and Edward D.F. hyperuricemia: a biomarker of renal hemodynamic impairment. Hypertension research Laboratory, Ochsner clinic, New Orleans, La., USA. *Ccardiorenal Med.* *5*:175-182. (2015)
67. Dirk JV, Luis MR, Alan SM, Kevin D., Biomarkers of renal injury and function: diagnostic, prognostic and therapeutic implications in heart failure. *European heart journal*, (2015). Vol. *37*:2577- 2585.
68. Adesokan, A. A., Oyewole, O. I., & Turay, B. M. S. Kidney and liver function parameters in alloxan-induced diabetic rats treated with aloe barbadensis juice extract. *Sierra leone journal of biomedical research*, *1*(1), 33-37. (2009).
69. Molitoris, B. A., Levin, A., Warnock, D. G., Joannidis, M., Mehta, R. L., Kellum, J. A., ... & Shah, S. V. Improving outcomes of acute kidney injury: report of an initiative. *Nature clinical practice Nephrology*, (2007). *3*(8), 439-442.
70. M. M. Yassin, H. I. Altibi, and A. E. El Shanti, "Clinical and biochemical features of type 2 diabetic patients in Gaza Governorate, Gaza Strip," *West African Journal of Medicine*, vol. 30, no. 1, pp. 51–56, 2011. View at: [Publisher Site](#) | [Google Scholar](#)
71. Luka, C.D. and Mohammed, A. Evaluation of the antidiabetic property of aqueous extract of *Mangifera indica* leaf on normal and alloxan-induced diabetic rats. *J. Nat. Prod. Plant Resour.*, (2012). *2* (2):239-243.
72. Gludovatz, B., Hohenwarter, A., Catoor, D., Chang, E. H., George, E. P., & Ritchie, R. O. A fracture-resistant high-entropy alloy for cryogenic applications. *Science*, (2014). *345*(6201), 1153-1158.
73. Rafiu A. A and Luka C. D. Evaluation of the antidiabetic property of aqueous extract of *ipomoea batata* leaf on hyperglycemia, hyperlipidemia, blood electrolytes and enzymatic antioxidants of streptozotocin induced diabetic rats. *J Res Diabetes Metab.* (2018). *4*(1): 021-026.
74. Ashafa, O.T., M.T. Yakubu, D.S. Grierson and A.J. Afolayan). Toxicological Evaluation of the Aqueous Extract of *Felicia muricata* Thunb. Leaves in Wistar Rats. *African Journal of Biotechnology* (2018). *8*:949-54.

75. Lara, O., Tong, X., Zborowski, M., & Chalmers, J. J. Enrichment of rare cancer cells through depletion of normal cells using density and flow-through, immunomagnetic cell separation. *Experimental hematology*, (2004). 32(10), 891-904.
76. Adane, T., Asrie, F., Getaneh, Z., & Getawa, S. White blood cells and platelet profiles of diabetic patients at University of Gondar specialized referral hospital: A comparative cross-sectional study. *Journal of Clinical Laboratory Analysis*, (2021). 35(6). <https://doi.org/10.1002/jcla.23808>
77. Wang, Y., Yang, P., Yan, Z., Liu, Z., Ma, Q., Zhang, Z., Wang, Y., & Su, Y. The Relationship between Erythrocytes and Diabetes Mellitus. *Journal of Diabetes Research*, (2021). <https://doi.org/10.1155/2021/6656062>
78. Belete Biadgo, Mulugeta Melku, Solomon Mekonnen Abebe, Molla Abebe. Hematological Indices And Their Correlation With Fasting Blood Glucose Level And Anthropometric Measurements In Type 2 Diabetes Mellitus Patients In Gondar, Northwest Ethiopia. (2016). *J Diabetes, Metabolic*