

Original Research Article

Response of Sulphur and Zinc on Yield and Economics of Maize. (*Zea mays* L.)

ABSTRACT

A field experiment was conducted during *Rabi* season of 2022 at Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) India. To study the Response of Sulphur and Zinc on yield and economics of *Rabi* Maize. The treatment ~~consist~~consists of Sulphur 20, 30, 40 kg/ha and Zinc 15, 20, 25 kg/ha. The soil of experimental plot was sandy loamy intexture, nearly neutral in soil reaction (pH 7.8), low in organic carbon (0.35%) available Nitrogen (163.42 kg/ha), available Phosphorus (21.96 kg/ha) and available Potassium (256.48 kg/ha). Results revealed, that the higher grain yield (6.42 t/ha) and higher stover yield (14.36 t/ha) were significantly influenced with application of Sulphur 40 kg/ha + Zinc 25 kg/ha. Higher Gross returns (INR 1,30,268.40/ha), Net returns (INR 90,007.00/ha) and maximum Benefit cost ratio (2.23) were also recorded in treatment-9 (Sulphur 40 kg/ha + Zinc 25 kg/ha).

Comment [dr1]: Additions of sulphur?

Comment [dr2]: Why those quantities

Keywords: Maize, *Rabi*, Sulphur, zinc, yield attributes and economics

INTRODUCTION

Maize (*Zea mays* L.) is one of the versatile crops grown throughout the tropical as well as temperate regions of the world. It belongs to family Poaceae or Graminae. Maize is one of the most important cereal crops in the world due to its high yielding, ease of processing, readily digested and less cost of cultivation. (Jaliya *et al.* 2008). Maize is used as a staple human food, fodder for livestock, poultry feed, for fermentation and many industrial purposes. Maize has immense potential in the tropics and yield of up to 7500 kg/ha can be obtained if the crop is managed properly. Unfortunately, yields are below 5000 kg/ha (FAO 2007).

Comment [dr3]: Please insert a more recent reference

In India, maize is grown in an area of 9.86 million hectares with production of 31.51 million tonnes and productivity of 3195 kg/ha. In Uttar Pradesh the area, production and

productivity of maize are 0.77 million hectares, 1.80 million tonnes and 2331 kg/ha respectively. (DA&FC- Directorate of economics and statistics 2020-2021). Among Indian states Madhya Pradesh and Karnataka has highest area under maize (15%) each followed by Maharashtra (10%), Rajasthan (9%) and Uttar Pradesh (8%). Andhra Pradesh is having the highest productivity and records as high as 12 t/ha. (IIMR- Indian Institute of Maize Research). Approximately in India, 47% is used as poultry feed. Of the rest of produce 13% is used as livestock feed and food purpose each. 12% for industrial purpose 14% in starch industry, 7% as processed food and 6% for export and other purpose.

Comment [dr4]: Please consult referentiation normative

Comment [dr5]: Referentiation normative

Comment [dr6]: This phrase redaction needs to be reviewed is not clear

The role of secondary nutrients in general and particularly sulphur has role in increasing the production of maize. Sulphur plays a vital role in the primary metabolism of higher plants and ~~involve is involved~~ in the synthesis of secondary metabolites. It not only influences yield but also improves crop quality owing to its influence on protein metabolism and ~~oi~~ lipids synthesis. It is involved in the synthesis of essential amino acids, like cysteine, cystine and methionine. (Singh, 2001).

Comment [dr7]: Review redaction

Comment [dr8]: Referentiation normative

Comment [dr9]: What is IAA?

Zinc, is an essential micronutrient, it is crucial for the growth and development of the Maize crop. In addition to controlling the plant growth hormone Indole Acetic Acid, Zinc plays a critical function in the metabolism of proteins and carbohydrates (IAA). It is a crucial part of dehydrogenase and proteinase and encourages the development of starch, seed maturity, and production. (Vinay Singh and Mamtapandey (2018)). Zinc deficiency is a common phenomenon in cereals, particular in coarse treatment, soil semi-arid regions. Balanced fertilization is the key to achieve higher productivity and nutrient use efficiency (Vinay Singh et al., 2015).

Comment [dr10]: This is not well referenced

Keeping these points in view, the present investigation entitled “Effect of Sulphur and Zinc on Yield and Economics of Rabi Maize (*Zea mays* L.)” was conducted during Rabi 2022-2023, at Crop Research Farm, SHUATS, Prayagraj (U.P.).

Comment [dr11]: The study objectives are normally stated at the end of the introduction, along with the research question or working hypotheses

Comment [dr12]: This statement has already been used at the beginning of the work

MATERIALS AND METHODS

A field experiment was conducted during Rabi season of 2021-22 at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, SHUATS, Prayagraj (U.P.) India. The soil of the experimental plot was sandy loamy in texture, nearly neutral in soil reaction (pH 7.8), low in organic carbon (0.35%). The experimental treatments consist of Sulphur 20 kg/ha + zinc 15 kg/ha, Sulphur 20 kg/ha + zinc 20 kg/ha, Sulphur 20 kg/ha + zinc 25 kg/ha, Sulphur 30 kg/ha + zinc

15 kg/ha, Sulphur 30 kg/ha + zinc 20 kg/ha, Sulphur 30 kg/ha + zinc 25 kg/ha, Sulphur 40kg/ha + zinc 15 kg/ha, Sulphur 40 kg/ha + zinc 20 kg/ha, Sulphur 40 kg/ha + zinc 25 kg/ha. The experiment was laid out in Randomized Block Design, with 9 treatments replicated thrice. The observations were recorded for Number of cobs per plant, Number of Grains/cob, seed Index (g), Grain yield (t/ha) Stover yield (t/ha) and harvest Index (%). The data were subjected to statistical analysis by analysis of variance method (Gomez and Gomez, 1976).

Formatted: Right: 0.11", Space Before: 0 pt

Comment [dr13]: The experimental treatment should be expressed in a table for a graphical support, you have it at the end of the manuscript, but as you don't mention it is very difficult to infer that there is a table

RESULT AND

DISCUSSION YIELD

ATTRIBUTES

Number of cobs per plant

Treatment 40 kg/ha sulphur + 25 kg/ha zinc resulted in significantly highest no. of cobs per plant (2.33). However, treatment 30 kg/ha sulphur + 20 kg/ha zinc, 30 kg/ha sulphur + 25 kg/ha zinc, 40 kg/ha sulphur + 20 kg/ha zinc were found to be statistically at par with 40 kg/ha sulphur + 25 kg/ha zinc.

The experimental data indicated that, the yield contributing characters namely cob/plant (1.34), number of rows/cob (14.19), number of seeds/cob (405), cob weight (97.88), cob length (14.82 cm) were significantly higher by seed treatment of Zn at 4g/kg with sulphur 20 kg/ha. Ali *et al*, 2013 found that application of sulphur at 25 and 35 kg/ha gave significant increase in no. of days of tasseling and silking.

Comment [dr14]: The methods section must consider how the results were estimated... how many counts were made, how the results were obtained

Number of grains per cob

Significant effect was observed by the statistical analysis of number of grains/cob. Treatment (9) 40 kg/ha sulphur + 25 kg/ha zinc recorded significant and highest number of grains/cob (533.8). However, treatment (6) 30 kg/ha sulphur + 25 kg/ha zinc and treatment (8) 40 kg/ha sulphur + 20 kg/ha zinc were found to be statistically at par with 40 kg/ha sulphur + 25 kg/ha zinc.

Comment [dr15]: The statistical report must be completed including the values obtained by ANOVA??

Singh *et al* 2021 revealed that application of Zinc 30 kg/ha resulted in higher No. of grains per cob (415.0). Sinha *et al* 1995 started that application of sulphur gave significant increase in number of rows/cob, cobs and grains weight of maize.

Comment [dr16]: This is an important result, however, without any visual help is very hard to interpret.... Please for all the results insert graphics

Grain yield (t/ha)

The grain yield showed increasing trend with the application of sulphur and zinc in maize. The highest grain yield was obtained with the treatment 40 kg/ha sulphur + 25 kg/ha zinc (6.42 t/ha). Treatment with 40 kg/ha sulphur + 20 kg/ha (6.20) zinc were found to be statistically at par with 40 kg/ha sulphur + 25 kg/ha zinc.

Yield and yield attributes were significantly affected by sulphur at different levels. Maize crop fertilized with 150 kg N/ha along with 45 kg S/ha significantly resulted in seeds/cob (30.65), test weight (20.90), grain yield (4.87). **Alam et al., (2003)**.

Stover yield (t/ha)

The stover yield of maize was also influenced by the application of sulphur and zinc. Highest stover yield (14.36 t/ha) was recorded highest in treatment (9) with 40 kg/ha sulphur + 25 kg/ha zinc. However no par values are observed. The lowest stover yield (12.96) was observed with treatment (1) with 20 kg/ha sulphur + zinc 15 kg/ha

Mehta 2005 reported that application of sulphur at 60 kg/ha gave highest uptake by maize grains and stover from his study. **Subhradip et al. 2022** stated that better photosynthetic mobilization was observed due to application of Zinc Sulphate at 60 kg/ha and 30 kg/ha respectively.

Harvest index (%)

The data showed significant difference in with 40 kg/ha sulphur + 25 kg/ha zinc (30.9) harvest index. However, treatment (6) with 30 kg/ha sulphur + 25 kg/ha zinc and treatment (8) 40 kg/ha sulphur + 20 kg/ha zinc were found to be statistically at par with 40 kg/ha sulphur + 25 kg/ha zinc.

The highest stover yield, biological yield and maximum harvest index was recorded under application zinc at 30 kg/ha. (**Chauhan et al., 2014**). Significant increase in the grain yield (4606 kg/ha) and stover yield (7115 kg/ha) with application of 40 kg S/ha compared to control. (**Choudhary et al., 2013**).

ECONOMIC ANALYSIS

Gross Returns

Observations regarding the economics of treatments are given in table 2.

Highest gross return (1,30,268.40 INR/ha) were obtained in treatment-9

(Sulphur 40kg/ha+ zinc25kg/ha)ascomparedto othertreatments.

NetReturns

Netreturn(90,007.00 INR/ha)were higherintreatment-9(Sulphur40kg/ha+zinc25kg/ha)ascomparedto othertreatments.

BenefitCostRation

Maximum Benefit Cost ratio (2.23) was found to be highest in treatment-9 with (Sulphur 40kg/ha+ zinc25kg/ha)ascomparedto othertreatments.

CONCLUSION

It is concluded that with the application of Sulphur 40 kg/ha in combination with the zinc 25 kg/ha(Treatment-9), has achieved maximum yield and economics. Significantly higher number of cobs per plant, number of grains/cob, grain yield, stover yield and harvest Index wererecordedandproveneconomicallyviablewithapplicationofSulphur40kg/haalongwiththezinc25kg/ha(Treatment-9).Thesefindingsarebasedononeseasontherefore,further trials may berequired forfurtherconfirmation.

REFERENCES

- Anonymous(2011).AreaandproductionofpearlmilletinIndia.AgriculturalStatistics,Directorateof Economics&Statistics,Govt. ofIndia.
- Alam MM, Islam MN. (2003); Effect of sulphur and nitrogen on the yield and seed quality of maize. *Journal of Biosciences*3:643-654.
- Ali A, Iqbal Z., Hassan SW, Yasin M, Khaliq T, Ahmad S 2013. Effect of nitrogen and sulphur on phenology, growth and yield parameters of maize crop. *Science International*;25(2): 363-366.
- Chauhan, T.M., Ali, J., Singh, S.P., Singh, S.B., 2014. Effect of nitrogen and zinc nutrition on yield, quality and uptake of nutrients by wheat. *Annals Plant Soil Research*16(2), 98-101.
- Choudhary R, Singh D, Singh P, Dadarwal RS, Chaudhari R. Impact of nitrogen and sulphur fertilization on yield, quality and uptake of nutrient by maize in southern Rajasthan. *Annals of Plant and Soil Research* 2013;15(2):118-121.
- Jagdeep Singh, Ram Partap, Amritpal Singh, Naresh kumar and Krity, (2021). Effect of Nitrogen and Zinc on Growth and Yield of Maize (*Zea mays* L.) *International Journal of Bio-resource and Stress Management* 2021, 12(3):179-185.

Comment [dr17]: The way meaning the HOW was the economic analysis made, the models or equations employed must be explicitly stated at the methodology section

Mehta YK, Shaktawat MS, Singhi SM. Influence of sulphur, phosphorous and farmyard manure on yield attributes and yield of maize (*Zea mays* L.) in southern Rajasthan conditions. *Indian Journal of Agronomy* 2005; **50**(3):203-205

Singh, M. V., Importance of sulphur in balanced fertilizer use in India. *Fertilizer News.*, **46**:31-35(2001).

Subhradip Bhattacharjee, V.M.Bhale, Pramod kumar, Rakesh Kumar. (2022). Effect of Potassium and Zinc Nutrition on Growth and Yield of Short Duration Maize (*Zea mays* L.) *Indian Journal of Agricultural Research* **56** (1): 22-27

UNDER PEER REVIEW

Table-1. Effect of Sulphur and zinc on yield attributes of Rabi Maize.

S. No.	Treatment combinations	Number of cobs/plant	Number of Grains/cob	Grain yield (t/ha)	Stover yield (t/ha)	Harvest Index (%)
1.	Sulphur 20 kg/ha + Zinc 15 kg/ha	1.83	417.3	4.93	12.96	27.5
2.	Sulphur 20 kg/ha + Zinc 20 kg/ha	1.87	428.5	4.97	13.13	27.5
3.	Sulphur 20 kg/ha + Zinc 25 kg/ha	2.00	483.8	5.58	13.79	28.8
4.	Sulphur 30 kg/ha + Zinc 15 kg/ha	1.93	460.8	5.23	13.30	28.2
5.	Sulphur 30 kg/ha + Zinc 20 kg/ha	2.10	489.2	5.83	13.95	29.5
6.	Sulphur 30 kg/ha + Zinc 25 kg/ha	2.17	509.6	6.10	14.12	30.2
7.	Sulphur 40 kg/ha + Zinc 15 kg/ha	1.97	471.0	5.33	13.59	28.2
8.	Sulphur 40 kg/ha + Zinc 20 kg/ha	2.23	517.8	6.20	14.22	30.4
9.	Sulphur 40 kg/ha + Zinc 25 kg/ha	2.33	533.8	6.42	14.36	30.9
	F test	S	S	S	S	S
	SEm (±)	0.10	8.20	0.10	0.04	0.37
	CD (p=0.05)	0.30	24.58	0.31	0.13	1.12

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Table-2. Economic analysis of different treatment combinations of *Rabi* Maize.

S. No.	Treatment combinations	Cost of cultivation	Gross returns	Net returns	B:Cratio
1.	Sulphur 20kg/ha+Zinc 15 kg/ha	38,774.25	1,006,146.60	61,840.35	1.59
2.	Sulphur 20kg/ha+Zinc 20 kg/ha	39,304.50	1,014,474.40	62,142.90	1.58
3.	Sulphur 20kg/ha+Zinc 25 kg/ha	39,834.75	1,13,616.60	73,781.85	1.85
4.	Sulphur 30kg/ha+Zinc 15 kg/ha	38,988.25	1,06,602.60	67,614.35	1.73
5.	Sulphur 30kg/ha+Zinc 20 kg/ha	39,518.50	1,18,569.60	79,051.10	2.00
6.	Sulphur 30kg/ha+Zinc 25 kg/ha	40,048.75	1,23,918.00	83,869.25	2.09
7.	Sulphur 40kg/ha+Zinc 15 kg/ha	39,200.25	1,08,651.60	69,451.35	1.77
8.	Sulphur 40kg/ha+Zinc 20 kg/ha	39,730.50	1,25,910.00	86,179.5	2.16
9.	Sulphur 40kg/ha+Zinc 25 kg/ha	40,260.75	1,30,268.40	90,007.00	2.23

Formatted: Spanish (Colombia)

Formatted: Font: 11 pt, Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)

Formatted: Spanish (Colombia)