

# Effect of sulphur on growth and yield of rice (*Oryza sativa* L.) varieties

## ABSTRACT:

The field experiment was conducted during *kharij* 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). The soil of experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.36 %), available N (171.48 kg/ha), available P (15.2 kg/ha) and available K (232.5 kg/ha). The experiment was laid out in Randomized Block Design with nine treatments, replicated thrice. The treatments consist of T1: MTU – 7029 + Sulphur 10Kg/ha, T2: MTU – 7029 + Sulphur 20Kg/ha, T3: MTU – 7029 + Sulphur 30Kg/ha, T4: BPT – 5204 + Sulphur 10Kg/ha, T5: BPT – 5204 + Sulphur 20Kg/ha, T6: BPT – 5204 + Sulphur 30Kg/ha, T7: ARIZE – 6444 + Sulphur 10Kg/ha, T8: ARIZE – 6444 + Sulphur 20Kg/ha, T9: ARIZE – 6444 + Sulphur 30kg/ha are used. The results showed that application of BPT – 5204 + Sulphur 30kg/ha, recorded significantly higher plant height (111.20 cm), number of tillers/plant (11.45), plant dry weight (53.07 g/plant), Crop Growth rate (16.07 g/m<sup>2</sup>/day) panicles/plant (17.00), number of grains per hill, panicle (100.69), Test weight (21.51 g), panicle length (20.20 cm), grain yield (5.62 t/ha), Straw yield (9.68 t/ha), Harvest index (36.72 %) were recorded with the treatment of BPT – 5204 + Sulphur 30kg/ha. The higher gross returns (Rs. 1,72,050.00/ha), net return (Rs. 1,12,790.00/ha) and benefit cost ratio (1.90) was obtained in the treatment of BPT – 5204 + Sulphur 30Kg/ha.

**Keywords:** Nitrogen, Phosphorous, Potassium, Rice.

## Introduction:

Rice is the seed of the grass species *Oryza sativa* (Asian rice) or less commonly *O. glaberrima* (African rice). The name wild rice is usually used for species of the genera *Zizania* and *Porteresia*, both wild and domesticated, although the term may also be used for primitive or uncultivated varieties of *Oryza*. As a cereal grain, domesticated rice is the most widely consumed staple food for over half of the world's human population, in Asia and Africa. It is the agricultural commodity with the third-highest worldwide production, after sugarcane and maize. Since sizable portions of sugarcane and maize crops are used for purposes other than human consumption, rice is the most important food crop with regard to human nutrition and caloric intake, providing more than one-fifth of the calories consumed worldwide by humans. There are many varieties of rice and culinary preferences tend to vary

regionally. Sulphur is one of the essential elements required for normal plant growth and plays an important role as a component of many plant processes. This is because plant metabolism is dependent on sulphur and a deficiency of this nutrient leads to impaired basal metabolism. Not only does it reduce yields, it also reduces product quality. If the supply of S is inadequate, applying high levels of other nutrients (N, P, and K) may not increase yields due to imbalances in plant N/S and P/S ratios. In addition to, a sufficient and balanced supply of nutrients promotes the correct development of plants and has a positive effect on yield. Therefore, under such circumstances, the application of S fertilizer is imminent to increase the yield of aerobic rice. Various sulfur fertilizers are available in India, but gypsum is the most common. Because it's cheap and readily available. Another inexpensive sulfur source, phosphogypsum, a by-product of the calcium superphosphate industry, has gained attention and market acceptance as a sulfur source. Sulfate leaching is expected under aerobic rice, so the use of slow-acting S carriers may be helpful (Nayaket *al.*, 2013).

### **Material and Methods:**

The field experiment was conducted during *khair*/2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). The experiment was laid out in Randomized Block Design with nine treatments including control each replicated thrice on the basis of one year experimentation. The treatments consist of T1: MTU – 7029 + Sulphur 10Kg/ha, T2: MTU – 7029 + Sulphur 20Kg/ha, T3: MTU – 7029 + Sulphur 30Kg/ha, T4: BPT – 5204 + Sulphur 10Kg/ha, T5: BPT – 5204 + Sulphur 20Kg/ha, T6: BPT – 5204 + Sulphur 30Kg/ha, T7: ARIZE – 6444 + Sulphur 10Kg/ha, T8: ARIZE – 6444 + Sulphur 20Kg/ha, T9: ARIZE – 6444 + Sulphur 30Kg/ha are used. The observations recorded are plant height, No. of tillers/hill, plant dry weight, no of panicles/hill, no of grains/hill, Grain yield. The collected data were subjected to statistical analysis by analysis of variance methods (Gomez and Gomez, 1984)

### **Results and Discussion:**

#### **Growth parameters:**

##### **Plant height:**

The perusal of data indicates that plant height measured at (i.e., 60 DAS), At 60 DAS, there was significant difference among the treatments. However, highest plant height (53.80 cm) was recorded with the application of BPT - 5204 + sulphur 30kg/ha, whereas minimum plant height (47.10 cm) was recorded with the treatment ARIZE – 6444 + Sulphur 10Kg/ha and BPT - 5204 + sulphur 20kg/ha (53.40 cm), MTU – 7029 + sulphur 30kg/ha (52.10 cm) were statistically at par with T6. There was significant difference among the treatments. Similar findings were

reported by **Anil kumaret al (2012)** ,**Ram et al. (2018)**, **Rajesh et al. (2018)** increase in plant height might be owing to the positive role of S in plant metabolic activity, which may have led to the increased photosynthesis and thereby plant height. These results clearly indicated that sulphur has a definite positive impact on plant height.

#### **No. of tillers/plant:**

However, **highest** number of tillers (7.90) was recorded with the application of BPT – 5204 + Sulphur 30Kg/ha, whereas minimum number of tillers (6.14) was recorded with the treatment ARIZE – 6444 + Sulphur 10Kg/ha and BPT - 5204 + sulphur 20kg/ha (7.66) was statistically at par with T6, there was significant difference among the treatments. Similar findings were reported by **Anil Kumar et al (2012)**, **Dinesh et al. (2018)**, **Ram et al. (2018)**, **Rajesh et al. (2018)** tillering is the product of expanding **auxillary** buds and closely associated with the nutritional conditions of the mother culm during its early growth period which gets improved by the **application** of sulphur.

#### **Plant Dry weight**

However, highest dry weight (14.00 gm) was recorded with the application of BPT - 5204 + sulphur 30kg/ha, whereas minimum dry weight (12.30 gm) was recorded with the treatment ARIZE – 6444 + Sulphur 10Kg/ha and BPT - 5204 + sulphur 20kg/ha (13.87 gm) was statistically at par with T6. Similar findings were reported by **Ram et al. (2018)**, **Anil Kumaret al (2012)**, **Rajesh et al. (2018)** increases in dry weight which is quite obvious that continued and balanced supply of nutrients right from early stage of growth resulted in **vigorous** plant growth and its results to better dry weight.

#### **Yield parameters:**

##### **No of panicles/hill:**

Significantly maximum **n**umber of **p**anicles/hill (17.00) was recorded with the treatment of application of BPT - 5204 + sulphur 30kg/ha and BPT - 5204 + sulphur 20kg/ha (17.00) over all the treatments, minimum was recorded in treatments ARIZE – 6444 + Sulphur 10Kg/ha (12.00) and MTU – 7029 + sulphur 30kg/ha (16.50) which was found to be statistically at par with T6. Similar findings were reported by **Bassiouniet al (2016)**, **Rahmanet al (2007)**, **Rajesh et al. (2018)** increases in panicles that owing to improving soil properties by reducing **ph** of saline soil drainage improvement, encouraging aggregates formation and nutrient availability reflecting on plant growth and salinity tolerance of rice and it improves the higher assimilation and translocation of carbohydrates to panicle induced by sulphur. And the sulphur also could stimulate synthesis of chloroplast protein resulting in greater photosynthetic efficiency which in

turn resulted in increased yield

#### **No. of grains/panicle:**

Significantly maximum number of grains/hill (100.69) was recorded with the treatment of application of BPT - 5204 + sulphur 30kg/ha over all the treatments, minimum was recorded in treatments MTU - 7029 + Sulphur 10Kg/ha (92.76) and BPT - 5204 + Sulphur 20Kg/ha (100.48), MTU - 7029 + Sulphur 30Kg/ha (99.35) was statistically at par with T6. Similar findings were reported by Anil Kumar *et al* (2012), Jiten Singh *et al* (2018), Rajesh *et al.* (2018) tillering is the product of expanding axillary buds and closely associated with the nutritional conditions of the mother culm during its early growth period which gets improved by the application of sulphur.

#### **Grain yield:**

Significantly maximum grain yield (5.62 t/ha) was recorded with the treatment, application of BPT - 5204 + sulphur 30kg/ha over all the treatments, minimum was recorded in treatments ARIZE - 6444 + Sulphur 10Kg/ha (4.50 t/ha) and BPT - 5204 + sulphur 20kg/ha (5.37 t/ha) which were found to be statistically at par with T6. Similar findings were reported by Ram *et al.* (2018), Jyothish *et al.* (2019) application of sulphur to soil increases the availability of S in soil which might have helped the crop to achieve better growth as it is an evident that S application significantly and positively increased the values of yield attributes, which might have increased the grain yield significantly.

#### **Conclusion:**

It is concluded that application of BPT - 5204 + Sulphur 50Kg/ha (Treatment 6) was recorded significantly higher grain yield (5.82 t/ha), and benefit cost ratio (2.92) as compared to other treatments.



**Table 1: Influence of varieties and sulphur on growth and Yield attributes and their combination on growth and yield attributes of Rice**

Treatments	Plant height	Number of tillers/plants	Dry weight	Number of panicle/hills	Number of grains/panicle	Grain yield (kg/ha)
T1	47.70	6.34	12.52	13.10	92.76	4.59
T2	51.60	7.30	13.21	16.00	98.78	4.89
T3	52.10	7.54	13.40	16.50	99.35	4.90
T4	48.60	6.54	12.61	14.50	96.39	4.96
T5	53.40	7.66	13.87	17.00	100.48	5.37
T6	<b>53.80</b>	<b>7.90</b>	<b>14.00</b>	17.00	100.69	5.62
T7	47.10	6.14	12.30	12.00	95.71	4.50
T8	49.40	7.02	13.08	14.80	97.25	4.96
<b>T9</b>	49.90	7.15	13.08	15.30	98.46	4.98
F- Test	S	S	S	S	S	S
SEm	0.79	0.12	0.20	0.2	1.53	0.08
CD(p=0.05)	2.39	0.35	0.61	0.59	2.58	0.26

## References:

- Anil Kumar Singh, Mani B., M. K. Meena & Ashutosh Upadhyaya (2012) Effect of Sulphur and Zinc on Rice Performance and Nutrient Dynamics in Plants and Soil of Indo Gangetic Plains; *Journal of Agricultural Science*; Vol. 4(11).
- Asha, R, Kumar, D, Singh N, Anjali A, (2014). Effect of sulphur on growth productivity and economics of aerobic rice (*Oryza sativa* L.) *Indian journal of Agronomy* 59(3):404-409
- Chandel, R.S., Sudhakar, P.C. and Singh, K. (2002). Effect of sulphur application on growth and yield of rice in rice–mustard crop-rotation sequence. *Crop Research* 24(2): 261–65
- Dewal, G.S. and Pareek, R.G. (2004). Effect of phosphorus, sulphur and zinc on growth yield and nutrient uptake of wheat (*Triticum aestivum* L.). *Indian Journal of Agronomy* 49(3): 160–62.
- Chandra, J.T, Sen, P, Kumar, A.D. (2019). Effect of sulphur on yield and yield attributes of rice and subsequent its residual effect on mustard and green gram crops. *Indian Agriculturist* 63:1-9
- Kumar, R, Verma, K.K, Ashok, K, Kumar, P, Yadav, K (2018). Effect of levels and sources of sulphur on growth and yield economics and quality of Rice (*Oryza sativa* L.) under partially reclaimed sodic soil. *The Pharma innovation journal* 7(8):41:44
- Nayak, A.K., Lal, B., Shahid, M., Panda, B.B., Tripathi, R., Raja, R. and Mohapatra, T. (2013). Fertilizer best management practices in rice for higher productivity. *Indian Journal of Fertilisers* 9(4): 54–66.
- Rahman, M. T., Jahiruddin, M., Humayan, M. R., Alam, M. J., & Khan, A. A. (2008). Effect of Sulphur and Zinc on Growth, Yield and Nutrient Uptake of Boro Rice (*Cv. BrriDhan 29*). *J. Soil. Nature.*, 2, 10-15.
- Singh, A. K., Meena, M. K., & Bharati, R. C. (2011). Sulphur and Zinc Nutrient Management in rice-lentil cropping system. International Conference on “Life Science Research for Rural and Agricultural Development” (pp 66-67). 27-29 December, 2011, CPRS Patna (Bihar).
- Singh, A.K., Mani B., K., Meena, M.K., and Upadhyaya, A., (2012). Effect of sulphur and zinc on rice performance and nutrient dynamics in plants and soils of Indo gangetic plains. *J. Agril. Sci.*, 4(11): 162.

Samaraweera, D.N. (2009.) Studies on forms and transformation of sulphur and response of rice to sulphur application in rice–rice cropping sequence. Ph.D. thesis, submitted to *University of Agricultural Sciences, Dharwad*.