

Original Research Article

Influence of Biofertilizers and Phosphorus on growth, yield and economics of field pea (*Pisum Sativum* L.)

ABSTRACT

The field experiment was conducted during *Rabi* season, 2022 at Crop Research Farm, Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P) to study the "Influence of Biofertilizers and Phosphorous on growth and Yield of Field Pea ". The results showed that treatment 9 [PSB + Rhizobium along with (70 kg/ha) Phosphorus] recorded significantly higher plant height (54.75 cm), more number of branches/plant (5.15), maximum plant dry weight (8.99 g/plant), maximum pod length (8.05), maximum number of pods/plant (24.97), higher number of grain/pod (7.24), maximum test weight (44.74 g), higher seed yield (16.25 q/ha), higher stover yield (37.11 q/ha) compare to other treatment. The maximum gross return (141133.00 INR/ha), maximum net return (101262.45 INR/ha) and highest B:C ratio (2.53) was recorded in treatment 9 application of [PSB + Rhizobium along with (70 kg/ha) Phosphorus] as compared to other treatments.

Keywords: Field pea, Biofertilizers, Phosphorus, Growth, Yield and Economics.

Introduction

Field pea is herbaceous, annual in habit and self-pollinated vegetable crop. The crop is grown for its green pods and seeds. The immature green seeds are consumed fresh, canned or in dehydrated jars and is leading frozen vegetable food. It is one of the most important vegetables in the world and ranks among top ten vegetable crops. India is second largest producer of pea in the world and accounts for 21% of the world production. Punjab is fifth largest producer of pea in the country and accounts for 6.7% of India's production. It is second important vegetable crop of Punjab and is grown on an area of 31.3 thousand hectare with annual production of 315.87 thousand tonnes. (Singh *et al.* 2019)

Pulses occupy a unique position in agriculture and are rich in protein, the protein content of field pea ranges from 15.5-39.7%. Fresh peas contain (per 100g) 44 calories, 75.6% water, 6.2g protein, 0.4 g fat, 16.9 carbohydrate, 2.4 g crude fiber, 0.9 g ash, 32 mg Ca, 102 mg P, 1.2 mg Fe, 6 mg Na and 350 mg K. (**Gowher *et al.*, 2013**) they maintain soil fertility through biological nitrogen fixation in soil and thus play a vital role in furthering sustainable agriculture. Pea belongs to the family Leguminosae. The field pea is a cool-season legume crop that is grown on over 25 million acres worldwide. The field pea is a type of pea sometimes called *P. sativum* subsp. *Arvense* (L.) *Asch.* It is also known as dun (grey-brown) pea, and it is one of the oldest domesticated crops, cultivated for at least 7,000 years. Field peas are now grown in many countries for both human consumption and stock feed. There are several cultivars and colors including blue, dun (brown), maple and white. The field pea is a cool-season legume crop that is grown all over worldwide. The total area of pulses was 28.83 million/ha. The total production of pulses 25.72 million tonnes and yield was 892 kg/ha. The production of Field pea in India 2020 - 21 was 7.92 lakh/ha. Uttar Pradesh having the highest production of field pea in 2020 - 21 by producing 4.953 lakh/ha. As of 49% of total production in India. (**GOI. 2021**)

Fertilizer being vital agriculture input to increase the production but the main drawbacks in the use and manufacture of chemical fertilizer viz., energy crises and inavailability of indigenous material like neptha, sulphur etc. at the national level and hazardous effect of chemical fertilizer on our health and environment. All these thing have led to research of alternative renewable source of nutrients to the crop through fertilizer of biological origin (bio-fertilizer). (**Ahmad *et al.* 2010**). In early growth stage, hairy vetch plants sometimes show very poor plant growth and exhibit nutritional disorder that the leaves and stems turn to red color. From the results we investigated, the poor plant growth and the nutritional disorder is caused by nitrogen deficiency. For that reason, hairy vetch cannot easily utilize soil nitrogen under low temperature conditions. On the other hand, it is considered that nitrogen deficiency is related to the poor nodule formation and low nitrogen fixation activity of the rhizobia in the nodules. (**Siddiqui and Debbarma 2022**)

Bio-fertilizers are known to play an important role in increasing availability of nitrogen and phosphorus besides improving biological fixation of atmospheric nitrogen and enhance

phosphorus availability to crop. Therefore, introduction of efficient strains of Rhizobium and PSB in soil, which is poor in nitrogen, may help in boosting up production and consequently more nitrogen fixation. Being a legume crop, major portion of N requirement of the crop is met through biological nitrogen fixation. Therefore, the present study was designed to assess the influence of phosphorus and bio-fertilizers on the growth and yield of field pea. **(Bhat *et al.* 2013)**

Rhizobium belongs to family Rhizobiaceae and is symbiotic in nature. Rhizobium has ability to fix atmospheric nitrogen in symbiotic association with legumes and certain non-legumes like Parasponia. It is useful for legumes like field pea. Rhizobium incorporated in pea rhizosphere through seed treatment probably induced more amount of nitrogen fixation in nodules of pea and solubilisation of fixed nitrogen from non-available to exchangeable pool which imparts more vegetative growth. **(Singh *et al.* 2020)**

Phosphate-solubilizing bacteria may also improve P availability and crop growth by promoting biological nitrogen fixation, through releasing growth promoters such as indoleacetic acid, gibberellic, and cytokinin's. PSB inoculation has been found to improve the yield and P nutrition of crops such as rice, maize, and other cereals. Thus, PSB can be an efficient, environmentally friendly and economically beneficial substitute for expensive P fertilizers. However, the potential of PSB in soils of a calcareous nature and with an alkaline reaction has not been well documented. **(Krishnaraj *et al.* 2014).**

Phosphorus is known to play an important role in growth and development of the crop and have direct relation with root proliferation, straw strength, grain formation, crop maturation and crop quality. The requirement of P, which is essential for root growth and nodulation, has to be largely fulfilled through inorganic fertilizers. **(Bhat *et al.* 2013)**. Keeping in view the above fact, the experiment was conducted to find out “Influence of Biofertilizers and Phosphorus on growth, yield and economics of field pea (*Pisum Sativum* L.)”

MATERIALS AND METHODS

This experiment was laid out during the *Rabi* season of 2022 at Crop Research Farm, Department of Agronomy, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj

(U.P.). The soil of the field constituting a part of central gangetic alluvium is neutral and deep. The soil of the experimental field was sandy loam in texture, nearly neutral in soil reaction (pH 8.0), low level of organic carbon (0.62%), available N (225 kg/ha), P (38.2 kg/ha) and K (240.7 kg/ha). The treatment consist of 3 different levels of Biofertilizers (PSB, Rhizobium and PSB + Rhizobium) with combination of 3 different levels of Phosphorus (50, 60 and 70 kg/ha). The experiment was laid out in RBD with 10 treatments each replicated thrice. The treatment combination are T₁ – PSB + 50 kg/ha P₂O₅, T₂ – PSB + 60 kg/ha P₂O₅, T₃ – PSB+ 70 kg/ha P₂O₅, T₄ – Rhizobium + 50 kg/ha P₂O₅, T₅ – Rhizobium + 60 kg/ha P₂O₅, T₆ – Rhizobium + 70 kg/ha P₂O₅, T₇ – PSB + Rhizobium + 50 kg/ha P₂O₅, T₈ – PSB + Rhizobium + 60 kg/ha P₂O₅, T₉ – PSB + Rhizobium + 70 kg/ha P₂O₅, T₁₀ - Control. The Data recorded on different aspects of crop, viz., growth parameters, yield attributes were subjected to statistically analysis by analysis of variance method. (Gomez and Gomez, 1976)

RESULT AND DISCUSSION:

Growth parameters

Plant height (cm)

The data revealed that a significantly higher plant height (54.75cm) was recorded in treatment 9 (PSB+ Rhizobium +70 kg/ha P₂O₅) as compared to rest of the treatments. However, treatment-8 (PSB+ Rhizobium +60 kg/ha P₂O₅) was found to be statistically at par with treatment 9 (PSB+ Rhizobium +70 kg/ha P₂O₅) in (Table 1). Significant and higher plant height was recorded with the application of PSB along with Rhizobium which may be due to seed inoculation which improved soil N and P status hence also subsequently boosted N and P uptake, increasing plant growth particularly plant height. Similar result was also reported by **Singh and Yadav. (2008)**. Further significant and higher plant height was recorded with the application of 70kg/ha P₂O₅ might be due to higher phosphorus levels which may have cause plants to grow taller, which could have a positive effect of phosphorus on root multiplication, nodulation, and the speeding up the height of plant. Similar results were also reported by **Ade et al. (2018)**

Number of branches/ plant

The data revealed that a significantly maximum number of branches/plant (5.15) was recorded in treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) as compare to rest of treatments. However, the treatment 8 (PSB + Rhizobium + 60 kg/ha P₂O₅) was found to be statistically at par with treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅). in (Table 1). Significant and higher plant height was recorded with the application of PSB along with Rhizobium which may be a result of PSB's enhancement of phosphorus availability to the plant and its ability to mitigate the negative effects of excessive nitrogen in the soil which may help to improve number of branches. Similar results were also reported by **Rather et al. (2010)**. Further significant and maximum number of branches/plant was recorded with the application of 70kg/ha P₂O₅ because phosphorus boosts Rhizobium activity, which improves N fixation in the root nodules as well as promotes better plant growth and development which lead to the enhanced cell division in addition to growth brought on by higher phosphorus levels which showed the production of more apical bud primordia and ultimately, more branches. Similar results were also reported by **Bhat et al. (2013)**

Plant dry weight

The data revealed that significantly maximum dry weight (8.99) was recorded in treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) as compare to rest of treatments. However, the treatment 8 (PSB + Rhizobium + 60 kg/ha P₂O₅) was found to be statistically at par with treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) in (Table 1). Significant and maximum dry weight was recorded with the application of PSB along with Rhizobium, this could be a result of symbiotic and non-symbiotic nitrogen fixation, which enhances the availability of phosphorus to the plant through PSB and reducing the negative effects of excessive nitrogen in the soil. Similar results were also reported by **Rather *et.al* (2010)**. Further significant and maximum dry weight was recorded with the application of 70kg/ha P₂O₅ which could be attributable to the favourable effect of phosphorus application on root growth and phosphorus availability which results in more root surface for bacterial infection and improved biological nitrogen fixation which helps in better dry matter production. Similar results were also reported by **Jayshree and Umesha (2021)**.

Crop growth rate (g/m²/day)

The data revealed that significantly higher crop growth rate (6.54) was recorded in treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) as compare to rest of treatments. However, the treatment 8 (PSB + Rhizobium + 60 kg/ha P₂O₅) was found to be statistically at par with treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) in Table 1. Significant and higher crop growth rate was recorded with the application of PSB along with Rhizobium which might be due to better accumulation of dry matter throughout the plant's vegetative and reproductive phase that enhances the physiological and metabolic activity and growth by assimilating the available nutrients at exponential rate and ease more photosynthesis, which results into higher crop growth rate. Similar results were also reported by **Jayshree and Umesha (2021)**.

Yield attributes

Pods length (cm)

The data revealed that significantly and higher pod length (8.05) was recorded in treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) as compare to rest of treatments. However, the treatment 8(PSB + Rhizobium + 60 kg/ha P₂O₅) was found to be statistically at par with treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) in Table 2. Significant and higher pod length was recorded with application of Rhizobium along with PSB which might be brought on by more nitrogen being available that promotes more vegetative growth, higher vegetative growth may be caused by inoculation in the pea rhizosphere through seed treatment, which likely increase the amount of nitrogen fixation in the pea nodules and cause the fixed nitrogen to be released from the exchangeable pool causing increased vegetative growth particularly pod length. Similar results were also reported by **Singh *et al.* (2020)**

Number of pods/plant

The data revealed that significantly and maximum number of pods/ plant (24.97) was recorded in treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) as compare to rest of treatments. However, the treatment 8(PSB + Rhizobium + 60 kg/ha P₂O₅) was found to be statistically at par with treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) in Table 2. Significant and maximum number of pods were with application of Rhizobium along with PSB may be because rhizobium introduced into the pea rhizosphere by seed treatment most likely increase nitrogen fixation in pea nodules and solubilization of fixed nitrogen from the non-available to the exchangeable pool results in increased number of pods/plant. Similar results were also reported by **Singh *et al.* (2019)**. Further significant and maximum number of pods/plant was recorded with the application of 70kg/ha P₂O₅ which may be because of the essential role of phosphorus in photosynthesis, fast energy transfer may have enhanced photosynthetic efficiency and consequently photosynthesis availability which further resulted in an increase in overall biomass production and plant part translocation. Similar results were also reported by **Hangsing *et al.* (2020)**

Number of grains/pod

The data revealed Significant and maximum number of grains/pod (7.24) was recorded in treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) as compare to rest of the treatments. However, the treatment 8(PSB + Rhizobium + 60 kg/ha P₂O₅) was found to be statistically at par with treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) in Table 2. Significant and maximum number of grains/pod was recorded with the application of 70kg/ha P₂O₅ which can be due to high phosphorus availability resulting in increased photosynthetic activity because phosphorus is a major constituent of ATP, and ATP is used in the dark reactions of photosynthesis which also increases the production of carbohydrates, sugars, starch, amino acids, and proteins, resulting in increase number of pods/ plant and seed yield, which eventually plays a role in increasing biological yield. Similar result were reported by **Kumar *et al.* (2022)**.

Test weight (g)

The data revealed significantly and higher test weight (44.74) was recorded in treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) as compare to rest of treatments. However, the treatment 8(PSB + Rhizobium + 60 kg/ha P₂O₅) was found to be statistically at par with treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) in Table 2. Significant and higher test weight was recorded with the application of 70kg/ha P₂O₅ due to phosphorus, which was brought about by increased photosynthesis, respiration, energy storage, cell division, and elongation, all of which ultimately improved the weight of seeds. Similar results were also reported by **Dey *et al.* (2021)**.

Grain yield (q/ha)

The data revealed Significant and higher grain yield (16.25) was recorded in treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) as compare to rest of treatments. However, the treatment 8(PSB + Rhizobium + 60 kg/ha P₂O₅) was found to be statistically at par with treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) in Table 2. Significant and higher grain yield was recorded with application of Rhizobium along with PSB which may be due to the combined effect of biofertilizers seed inoculation might have improved yield attributes specially number of

Pods/plant and number of grain/pod than Rhizobium or PSB inoculation alone. Similar results were also reported by **Abisha and Singh (2022)**. Further, significant and maximum grain yield was recorded with the application of 70kg/ha P₂O₅ which may be caused by improved root proliferation, higher development of roots, increased nutrient availability and uptake energy conversion along with plant metabolic activities. Similar results were also reported by **Yadav et al. (2017)**.

Stover yield (q/ha)

The data revealed significant and higher stover yield (37.11) was recorded in treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) as compared to rest of treatments. However, the treatment 8 (PSB + Rhizobium + 60 kg/ha P₂O₅) was found to be statistically at par with treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) in Table 2. Significant and higher stover yield was recorded with application of Rhizobium along with PSB which may be determined by the genetically influenced yield traits rather than the external environmental influences. Similar results were also reported by **Siddiqui and Debarma (2022)**. Further, significant and higher stover yield was recorded with the application of 70kg/ha P₂O₅ which may have enhanced fixation of nitrogen along with nodulation and greater amount of nitrogen fixation will boost crop yield. Similar results were also reported by **Singh et al. (2011)**.

Harvest Index (%)

At harvest, there was non-significant difference between the treatments and maximum harvest index (30.45 %) was observed with the applications of PSB + Rhizobium + 70 kg/ha P₂O₅, whereas the lowest value (31.06) was observed in treatment 10 (Control) in Table 2. Significant and higher harvest index was recorded with application of Rhizobium along with PSB that might be due to increased and balanced availability of both N and P in dual inoculation which could be attributable to seed inoculation with bio-fertilizers, because of the increased nitrogenase activity and accessible P status of the soil, the synergistic effect of Rhizobium and PSB may have increased growth, yield characteristics, and eventually higher harvest index values observed with

dual inoculation indicated increased photosynthate production and translocation to the sink. Similar reports were also recorded by **Bhat *et al.* (2013)**.

Economic

The result showed that, maximum Gross return (141133.00 INR/ha), maximum net return (101262.45 INR/ha) and highest B:C ratio (2.53) was recorded maximum in treatment 9 (PSB + Rhizobium + 70 kg/ha P₂O₅) in (Table 3). Higher benefit cost ratio was recorded with PSB along with rhizobium which might be due to higher grain and stover yield with combine doses of biofertilizer inoculation with seeds. Further, higher gross return and net return was with application of phosphorus (70 kg/ha) which might be due to phosphorus, by means of energy transfer plays a fundamental role in photosynthesis, increasing photosynthetic efficiency and thus the availability of photosynthates which helps to obtain higher benefit cost ratio. Similar report was also recorded by **Nsiah *et al.* (2018)**

CONCLUSION:

Based on the above findings it is concluded that in field pea, application of PSB along with Rhizobium and 70 kg/ha P₂O₅ (Treatment 9) was observed with higher seed yield and benefit cost ratio.

REFERENCE

1. **A.H. Mir, S.B. Lal, M. Salmani, M. Abid and I. Khan (2013)**. Growth, yield and nutrient content of blackgram (*Vigna mungo*) as influenced by levels of phosphorus, sulphur and PSB. *SAARC J. Agri.*, **11**(1): 1-6.
2. **Abisha P. and Shikha Singh (2022)**. Effect of biofertilizer and phosphorus on growth, yield components and yield of chickpea. *International Journal of Plant & Soil Science*, **34**(20): 326-331.
3. **Aluri Manoj Kumar, C. Umesha and Gorla Venkata Raju (2022)**. Effect of phosphorus and boron levels on growth and yield of chickpea. *International Journal of Plant & Soil Science*, **34**(21): 266-271.
4. **Farhin Siddiqui and Victor Debbarma (2022)**. Effect of biofertilizers and organic manures on growth and yield of field pea. *The Pharma Innovation Journal* **11**(11): 315-318.
5. **Gomez, K.A and Gomez, A.A. (1976)**. Three more factor experiment in: Statistical procedure for agricultural Research 2nd edition.P.139-141.
6. **Jakir Hussain, Poonam Kashyap, AK Prusty, Debashis Dutta, Shiv Shankar Sharma, AS Panwar and Sunil Kumar (2019)**. Effect of phosphorous fertilisation on growth, yield and quality of pea(*Pisum sativum*). *Indian journal of Agricultural Sciences*, **89**(8): 1303-7
7. **Jayshree and Umesha C. (2021)**. Effect of biofertilizers and phosphorus on growth parameters and yield of cowpea in sandy loam soil of Prayagraj. *Environment Conservation Journal*, **22**(1&2): 137-141.
8. **Konda Saikishore, Vikram Singh and Dhananjay Tiwari (2020)**. Effect of levels of phosphorus and zinc on growth and yield of Blackgram(*Vigna mungo* L.). *International Journal of Current Microbiology and Applied Sciences*, **9**(10): 1933-1937.

9. **N. Hangsing, L. Tzudir, A.P. Singh (2020)**. Effect of spacing and levels of phosphorus on growth and yield of green gram under rainfed condition of Nagaland. *Agricultural Science Digest*, (): D-5022 [1-5].
10. **Peter Fernandes and Satish A. Bhalerao (2015)**. Effect of Biofertilizer on the growth and Biochemical parameters of Mungbean (*Vigna radiata* L.). *International Journal of Advanced Research in Biological Sciences*, **2**(4): 127-130.
11. **R.S Singh and M.K Yadav (2008)**. Effect of phosphorus and biofertilizers on growth, yield and nutrient uptake of long duration pigeonpea under rainfed condition. *Journal of Food Legumes*, **21**(1): 46-48.
12. **Rakesh Mandloi, Sukhlal Waskel, Sunil Kumar Jatav, K.N. Tambi and D.R Agashe (2020)**. Varietal and Phosphorus Response on Pod Yield of Pea (*Pisum sativum* L.). *International Journal of Current Microbiology and Applied Sciences*, **10**: 213-221.
13. **Ripudaman Singh, Sanjiv Kumar, Hemant Kumar, Mukesh Kumar, Ashok Kumar and Dharmendra Kumar (2017)**. Effect of irrigation and integrated nutrient management on growth and yield of chickpea. *Plant Archivers*, **17**(2): 1319-1323.
14. **S.Adjei-Nsiah, B.U.Alabi, J.K.Ahiakpa and F.Kanampiu (2018)**. Response of Grain Legumes to Phosphorous Application in the Guinea Savanna Agro-Ecological Zones of Ghana. *Agron J.*, **110**(3): 1089-1096.
15. **Sarita Rani, Parveen Kumar and Anil Kumar Yadav (2017)**. Effect of biofertilizers in conjunction with chemical fertilizers on growth behaviour and profitability of field pea (*Pisum sativum* L.) grown in western plains of Haryana. *Chemical Science Review and Letters*, **6**(22), 801-805.
16. **Shabeer Ahmad Rather, Mir Aashiq Hussain and Narinder Lal Sharma (2010)**. Effect of biofertilizers on growth, yield and economics of field pea. *International Journal of Agricultural Sciences*, **1**: 65-66.

17. **Subbarao and Joy Dawson (2022)**. Effect of phosphorus and zinc on growth and yield of field pea (*Pisum sativum*). *The Pharma Innovation Journal*, **11**(5):254-256.
18. **Suman Kumar Dey, Vikram Singh and Dhananjay Tiwari (2021)**. Effect of phosphorus and sulphur on the growth and yield of summer mungbean. *The Pharma Innovation Journal*, **10**(8): 1442-1445.
19. **Talwinder Singh, Harish Chandra Raturi and S.P. Uniyal (2021)**. Effect of biofertilizer and mulch on yield and quality of pea (*Pisum sativum* L.). *Indian Journal of Agricultural Research*, 10.18805/IJ Are.A-5696 [1-6].
20. **Talwinder Singh, Harish Chandra Raturi, Dilip Singh Kachwaya, Sandeep Kumar Singh and Arinderpal Kaur (2019)**. Effect of biofertilizer and mulch on yield and quality of pea. *Journal of Pharmacognosy and Phytochemistry*, SPI 205-208.
21. **Tauseef A. Bhat, Meenakshi Gupta, S. Sheraz Mahdi, Manzoor A. Ganai, Hilal A. Bhat, J.A. Bhat, Iqbal A. Wani and M. Hussain Dar (2013)**. Growth, yield and economics of field pea as influenced by phosphorus and biofertilizers under subtropical conditions of Jammu. *Journal of Pure and Applied Microbiology*, Vol. **7**(1), p. Proof.
22. **U.K Ade, A.S. Dambale and D.B. Jadhav (2018)**. Effect of phosphorus and biofertilizer on growth, yield and economics of pigeonpea under rainfed condition. *International Journal of Current Microbiology and Applied Sciences*, **6**: 1408-1416.

Table 1: Influence of Biofertilizers and phosphorus on the growth attributes of field Pea.

S.no	Treatments combination	Plant height(cm) (80DAS)	Number of branches/plant (80DAS)	Plant Dry weight (g/plant) (80DAS)	CGR (g/m ² /day) (60-80)	RGR (g/g/day) (60-80)
1.	PSB inoculation + 50 kg/ha P ₂ O ₅	42.77	3.27	5.26	3.42	0.0248
2.	PSB inoculation + 60 kg/ha P ₂ O ₅	45.13	3.42	5.59	3.80	0.0262
3.	PSB inoculation + 70 kg/ha P ₂ O ₅	46.32	3.66	5.86	3.97	0.0261
4.	Rhizobium inoculation + 50 kg/ha P ₂ O ₅	48.16	4.04	6.16	4.05	0.0251
5.	Rhizobium inoculation + 60 kg/ha P ₂ O ₅	49.54	4.36	6.62	4.27	0.0245
6.	Rhizobium inoculation + 70 kg/ha P ₂ O ₅	50.80	4.54	7.29	4.91	0.0258
7.	PSB inoculation + Rhizobium inoculation + 50 kg/ha P ₂ O ₅	51.90	4.64	8.09	5.93	0.0290
8.	PSB inoculation + Rhizobium inoculation + 60 kg/ha P ₂ O ₅	53.37	4.86	8.72	6.75	0.0319
9.	PSB inoculation + Rhizobium inoculation + 70 kg/ha P ₂ O ₅	54.75	5.15	8.99	6.54	0.0285
10.	Control	39.15	3.02	5.01	3.19	0.0161
	F-test	S	S	S	NS	NS
	SEm (±)	2.06	0.36	0.39	0.43	0.0019
	CD (P=0.05)	6.15	1.08	1.67	-	-

Table 2: Influence of biofertilizers and phosphorous on yield and yield attribute of field pea.

S.no	Treatments combination	Pod length (cm)	Number of pod/ plant	Number of grain/pod	Test weight (g)	Seed yield (q/ha)	Stover yield (q/ha)	Harvest index (%)
1.	PSB inoculation + 50 kg/ha P□O□	5.29	17.24	3.73	36.36	12.37	27.69	30.89
2.	PSB inoculation + 60 kg/ha P□O□	5.52	17.80	4.03	37.32	12.78	29.28	30.39
3.	PSB inoculation + 70 kg/ha P□O□	5.78	18.82	4.44	38.17	13.22	31.18	29.78
4.	Rhizobium inoculation + 50 kg/ha P□O□	6.17	19.98	5.08	39.53	13.62	32.57	29.49
5.	Rhizobium inoculation + 60 kg/ha P□O□	6.51	21.36	5.47	40.65	14.00	33.49	29.48
6.	Rhizobium inoculation + 70 kg/ha P□O□	6.90	22.51	6.03	41.91	14.46	34.35	29.63
7.	PSB inoculation + Rhizobium inoculation + 50 kg/ha P□O□	7.33	23.43	6.35	42.65	15.15	35.06	30.18
8.	PSB inoculation + Rhizobium inoculation + 60 kg/ha P□O□	7.66	23.94	6.78	43.56	15.69	36.23	30.22
9.	PSB inoculation + Rhizobium inoculation + 70 kg/ha P□O□	8.05	24.97	7.24	44.74	16.25	37.11	30.45
10.	Control	5.06	16.22	3.35	33.95	10.58	23.48	31.06
	F-test	S	S	S	S	S	S	NS

	SEm (\pm)	0.39	1.11	0.36	1.87	0.78	1.64	0.66
	CD (P=0.05)	1.16	3.29	1.07	5.55	2.31	4.89	-

Table 3: Influence of biofertilizers and phosphorus on economics of field pea.

Sr. No.	Treatment combination	Economic			
		Cost of cultivation (INR/ha)	Gross return (INR/ha)	Net return (INR/ha)	B:C
1.	PSB inoculation + 50 kg/ha P ₂ O ₅	38969.55	107267.00	68297.45	1.75
2.	PSB inoculation + 60 kg/ha P ₂ O ₅	39532.05	111024.00	71491.95	1.80
3.	PSB inoculation + 70 kg/ha P ₂ O ₅	40094.55	115114.00	75019.45	1.87
4.	Rhizobium inoculation + 50 kg/ha P ₂ O ₅	38521.55	118731.00	80209.45	2.08
5.	Rhizobium inoculation + 60 kg/ha P ₂ O ₅	39084.05	122047.00	82962.95	2.12
6.	Rhizobium inoculation + 70 kg/ha P ₂ O ₅	39646.55	125985.00	86338.45	2.17
7.	PSB + Rhizobium inoculation + 50 kg/ha P ₂ O ₅	38745.55	131718.00	92972.45	2.39
8.	PSB + Rhizobium inoculation + 60 kg/ha P ₂ O ₅	39308.05	136389.00	97080.95	2.46
9.	PSB + Rhizobium inoculation + 70 kg/ha P ₂ O ₅	39870.55	141133.00	101262.45	2.53
10.	Control	35517.05	91684.00	56166.95	1.58