

Original Research Article

Evaluation of the Fertility Indices of Female Mice Exposed to Dimethoate and Beetroot Extract (*Beta vulgaris*)

Abstract

Aim: The aim of this study was to evaluate of the fertility indices of female mice exposed to Dimethoate and Beetroot Extract (*Beta vulgaris*).

Experimental Design: A completely randomized experimental design using standard methods for analysis.

Location and Duration of Study: This study was carried out in the Animal house, Department of Animal and Environmental Biology of Rivers State University, Nkpolu-Oroworukwo Port Harcourt, Nigeria. GPS 4°47'50"N 6°58'49"E. This study lasted for 21 days.

Methodology: Thirty (30) sexually matured female mice were randomly divided into 6 groups (A-F) with 5 mice each. They were exposed for a period of 21 days to Dimethoate and 100% extract of *Beta vulgaris*. Group A was orally administered distilled water and rodent pellet only and acted as control. Group B was administered 20mg/kg/day of Dimethoate only. Group C was administered 5mg/kg/day of Dimethoate + 250mg/kg/day of 100% fresh extract of *Beta vulgaris*. Group D was administered 10mg/kg/day of Dimethoate + 250mg/kg/day of 100% fresh extract of *Beta vulgaris*. Group E was administered 15mg/kg/day of Dimethoate + 250mg/kg/day of 100% fresh extract of *Beta vulgaris*. Group F was administered 20mg/kg/day of Dimethoate + 250mg/kg/day of 100% fresh extract of *Beta vulgaris*. At the end of the 21 days exposure the final weight of the mice in each group was taken and blood collected by ocular puncture for hormonal profiling. Hormones such as Prolactin, Progesterone, estradiol, Follicle Stimulating Hormone (FSH) and luteinizing hormone (LH) were analyzed based on the manufacturer's instruction using Radox Monza assay kit. The females were then allowed to mate with fifteen unexposed males in ratio 2:1. At the end of 24 days and following successful parturition essential data were collected from the F₁ animals.

Result: Results showed significant increase in body weight for groups that were co-administered *Beta vulgaris* as compared to Group B that was administered Dimethoate only. Litter size gradually decrease significantly from seven in the control to two in group B which received Dimethoate only. Reproductive hormones significantly increased in groups coadministered *Beta vulgaris* and dimethoate compared with group B administered dimethoate only.

It is thus deduced that B. vulgaris might increase fertility in experimental animals, while Dimethoate showed significant impact on the reproduction in female experimental animals by reducing the fertility indices, such as reproduction hormones, reduction in fetal implantation leading to low number of offspring (Litter size).

Keywords: Beetroot, Dimethoate, Fertility indices, Hormones, Parturition,

1.0 Introduction

Pesticides constitute a major group of harmful chemicals released into the environment. A suspected mechanism of the disruption of female reproductive function is the induction of hormonal imbalance reported to be brought about by pesticides.[1]

The increased use of pesticides all over the world is alarming and this has drawn the attention of researchers due to the adverse effects they have on non-target organisms [2].

Pesticides are used in agriculture to enhance food production by eradicating unwanted pests and controlling disease vectors. Organophosphorus compounds are the [groups of pesticides widely used in agriculture \[3\]](#). Dimethoate is [groups of pesticides widely used in agriculture \[3\]](#). Dimethoates are an organophosphate insecticide with numerous uses on both agricultural and non-agricultural site [4,5,5]. In human, the main groups at risk of exposure to higher doses of dimethoate are producers and workers at pesticides factories and farm owners. Indeed, majority of population are exposed to lower doses of dimethoate through food contamination, drinking water or by application of household insecticides containing [Dimethoate](#)[[Dimethoate](#)]6]. Several physiological and behavioral dysfunctions occur in animals after exposure to low doses of organophosphate [pesticide,pesticide](#); hence it is possible that dimethoate could affect humans and wildlife in their natural habitat. According to [7] dimethoate has been reported to cause developmental toxicity as well as reproductive failures in organisms upon repeated exposures. Developmental and Reproductive toxicity of this pesticide on adult rodents of both sexes has been demonstrated including decrease number of implantation and live fetuses, incidences of resorptions and decrease fetal body weights, Irregularities of estrous cycle and altered levels of serum

gonadotrophins in females while impairment of fertility, suppressed libido low semen quality and altered testosterone levels and testicular degeneration have been reported in males [1,8,9].

Dimethoate at 28mg/kg was reported to be associated with a significant decreased of sperm count, motility and viability, and significantly increased percentage of morphologically abnormal spermatozoa compared with the controls [9].

The primary site of action of organophosphate pesticides in mammals is the central and peripheral nervous system and it's by inhibiting acetylcholinesterase, a biochemical event that result in accumulation of endogenous acetylcholine at the nerve ending [10].

Beta vulgaris is a plant in the Chenopodiaceous family. It is the well-known in its numerous cultivated varieties, the most well-known is the purple root vegetable known as beetroot or table garden beet.

Beetroot can be eaten raw, used for juice extraction baked or boiled. Beets have used in traditional medicine for hundreds of years to treat constipation, gut and joint pain, dandruff [11].

It was shown by modern pharmacology that red beets extract exhibit anti-anemic, anti-ischemic, anti-inflammatory, antioxidant, ~~anti~~and anti-carcinogenic, antihypertensive and hypoglycemic activities [12]. Not only does the beetroot have nutritional value, its leaf also does but they are rarely eaten. However, the uniqueness of red beets does not reside in their nutritional value.

Beta vulgaris has been ranked among the most potent antioxidant vegetables [13]. As a known antioxidant vegetables and fruits nutrient are important to maintain human health and neutralize a negative effect of oxidative stress in human. Oxidative stress is indicated in ~~many disease~~many diseases including atherosclerosis neuro degenerative disease, Myocardial infarction, cancer as well as ageing [14].

Beside considerable antioxidant and anti-inflammatory activity, *Beta vulgaris* phytochemicals provide an anti-proliferative effect in breast, liver, colon and bladder cancer cell lines through the induction of both intrinsic and extrinsic apoptotic pathway [15]. However, it is important to note that Red beet belongs to oxalate-accumulating plants. Oxalic acid is one of the ~~compound~~compounds present in beetroot at high concentration and may have some side effects. As a result of this, regular consumption of red beet is contraindicated for people with urolithiasis.

Pesticides are made for the eradication of pest as the name implies which are the target species but still affect non-target species such as humans and animals. In developing nations such as Nigeria, women play an important role in agriculture irrespective of their reproductive status.

During pregnancy, the fetus can be exposed to pesticides during farming thereby endangering the health of the unborn child. Since the use of pesticides cannot completely be avoided, this study was undertaken to evaluate the role of Beetroot as a potential antioxidant in mitigating the effect of this pesticides in human.

2.0 Materials and methods

2.1 Experimental Location

The experiment was carried out in the animal house of the Department of Animal and Environmental Biology, Rivers State University, Port Harcourt. (Coordinates 4°48'14"N 6°59'12"E)

2.2 Experimental Animals

Thirty sexually matured female mice (mean weight 23.0 ± 2.24 g) were used in the study. The mice were housed individually in plastic cages under standard conditions (12 hL:12hD; 12hD room temperature $26 \pm 2^\circ\text{C}$ relative humidity 54%) and was acclimatized for 14 days before commencement of experimentation. The animals had free access to feed (rodent pellet) and cool clean water. All experiments were conducted according to the institutional animal care protocols at the Rivers State University Port Harcourt, Nigeria and follow guidelines for the ethical treatment of experimental animals.

2.3 Chemical

Dimethoate an organophosphate insecticide was obtained from a reputable agrochemical outlet in Port Harcourt, Rivers State.

2.4 Plant Extract

Beetroot which belongs to the family of Chenopodiaceae was bought from mile1 market in Port Harcourt and washed thoroughly to remove dirt. 100% of Beetroot extract was extracted using a juice extractor (model Bender ST-002).

2.5 Experimental Design and Procedure

Thirty (30) adult mice were randomly divided into 6 groups (A-F) with 5 mice each. they were exposed for a period of 21 days to Dimethoate and 100% extract of *Beta vulgaris*.

Group A was orally administered distilled water and rodent pellet only and acted as the control. Group B was administered 20mg/kg/day of Dimethoate only. Group C was administered 5mg/kg/day of Dimethoate + 250mg/kg/day of 100% fresh extract of *Beta vulgaris*. Group D was administered 10mg/kg/day of Dimethoate + 250mg/kg/day of 100% fresh extract of *Beta vulgaris*. Group E was administered 15mg/kg/day of Dimethoate + 250mg/kg/day of 100% fresh extract of *Beta vulgaris*. Group F was administered 20mg/kg/day of Dimethoate + 250mg/kg/day of 100% fresh extract of *Beta vulgaris*. At the end of the 21 days exposure the final weight of the mice in each group was taken.

All animals were observed daily for behavioral changes and mortality as well as food and water intake throughout the experimental duration. Animals were weighed and the weight recorded daily to the nearest 0.01gram. For hormonal profile analysis, blood was collected in a plain venipuncture tube by ocular puncture. The blood was allowed to clot and centrifuged at 3000g to separate the serum from the cells. Samples were stored at 4°C till analysis. Hormones such as Prolactin, Progesterone, estradiol, Follicle Stimulating Hormone (FSH) and luteinizing hormone (LH) were analyzed based on the manufacturer's instruction using Randox Monza assay kit

2.6 Mating Procedures

At the end of the twenty-one (21) days administration, fifteen (15) males that were not previously exposed to the pesticide were introduced to the 30 females of the same age and species. They were paired in the ratio 1: 2 for mating purposes. Twenty-four hours after introduction of males to the females, vaginal swap was taken and examined under the microscope to ensure that mating had taken place. The gestation length was fixed at 21 – 24days. The gestation length, body weight, liter size were determined and pups were checked for deformities. At the end of 24 days and following successful parturition essential data were collected from the F₁ animals.

Statistical Analysis: Data were subjected to analysis of variance using the SPSS 20 software for determination of Mean and Standard deviation. Mean separation where significant difference was observed was carried out using Tukey's pairwise comparison test.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

3.0 RESULTS

3.1 Visual observation

There were no visible changes in the physical appearances of mice during the period of study. However, animals showed signs of restlessness, itching of paws and increased feeding pattern especially in the group that received dimethoate only without coadministration of *Beta vulgaris*.

3.2 Effect of *Beta vulgaris* on the body weight of Swiss mice

The effect of oral administration of Dimethoate and *Beta vulgaris* on bodyweight in Swiss mice is shown in Figure 1A-F.

Fig 1A shows the effect of oral administration of Dimethoate and *Beta vulgaris* on the bodyweight of animals in the group A for 21 days. The initial body weight increased from 26.34 ± 2.56 g to 28.84 ± 2.34 g in week 1 and gradually increased to 30.52 ± 1.99 g in week 2. A steady increase was also observed in week 3 with 31.18 ± 2.04 g. By the week 3, a statistically significant increase equivalent of 20% weight gain was observed.

In the group B animals administered Dimethoate only the body increased steadily from 26.65 ± 2.04 g in week 1 to 27.85 approximately 4.80% increase in wk 2 followed by 29.31 ± 1.92 g in week 2 (5.90%) and 31.89 ± 1.21 g (-1.21 g (9.20%) in week 3, respectively. This was approximately 0.5% reduction in weight gain considered statistically non-significant compared to the control group.

Fig 2c shows the effect of oral administration of Dimethoate and *B. vulgaris* on the body weight of animals in the group C exposed to 5mg/kg/bw/day and 250mg/kg/bw/day of 100% fresh extract of *B.vulgaris*. The initial body weight of 26.53 ± 2.35 g increased to 29.69 ± 2.12 g in week 1 and 32.63 ± 2.04 g, 33.89 ± 1.09 g in week 2 and week 3 respectively. There was significant ($p < 0.05$) increase 27.74% in body weight gain with administration of *B. vulgaris*.

Group D animals also showed a gradual body weight increase from initial weight of 26.63 ± 1.81 g to 28.86 ± 1.56 g in week 1, 29.91 ± 2.11 g, 30.35 ± 1.23 g in week 2 and week 3, respectively indicating 13.97% weight gain.

Group E and Group F also showed a gradual increase in body weight throughout the duration of experiment. However, the increase recorded in all groups was not statistically different from group A that was not exposed to any treatment. Feed intake was also not adversely affected by any of the regime of treatments.




Fig 1a. Body weight of Group A animals




Fig 1b. Body weight of Group B animals





Fig 1c. Body weight of Group C animals




Fig 1d. Body weight of Group D animals




Fig 1e. Body weight of Group E animals

Fig 1f. Body weight of Group F animals

3.3 Effect of Varying Concentrations of Dimethoate and Coadministration of *Beta vulgaris* on Hormonal Profile of the Female Swiss Mice.

The concentrations of the hormone Prolactin was not significantly different from the control animal in groups A, C-F. However, Prolactin level was significantly ($p < 0.01$) elevated in group B (Fig. 2 A, 1.40ng/ml) which received 20mg/kg/bw/day of Dimethoate only.

Fig. 2B shows the effect of coadministration of Dimethoate and *Beta vulgaris* in Swiss mice Estradiol. The level of Estradiol was significantly (< 0.05) reduced in group B as compared to the control. Other treatment groups C-F coadministered 250mg/kg/bw/day had elevated Estradiol from 51pg/ml in C treated with 5mg/kgbw/day along with 250mg/kgbw/day to D administered 10mg/kgbw/day and coadministered 250mg/kgbw/day followed by 52 and 50mg/kgbw/day respectively.

Comparison of the concentration of Progesterone level between the control group and the treatment groups showed significantly (< 0.05) reduction (7.23 vs 12.67ng/ml) in group B that did not receive extract of *Beta vulgaris* and control, respectively. There was increased concentration in the D, E, E, E, F groups that received 250mg/kgbw/day of *B. vulgaris* extract (Fig. 2C)

Influence of varying concentrations of Dimethoate and coadministration of *B. vulgaris* on the concentration of Luteinizing hormone is shown in Fig. 2D. There was gradual elevation of hormone from

(1.42 iu/L) in control group to 2.35, 1.64, 2.30, 1.70, 1.70 i.u/L) for groups C-F, respectively. The lowest concentration was recorded in group B which received 20mg/kgbw/day of Dimethoate only.

The impact of Dimethoate and coadministration of *B. vulgaris* on concentration of Follicle Stimulating Hormone is shown in Fig. 2E. The control group had a concentration of 0.30 iu/ml. The value of the hormone increased significantly ($p < 0.05$) to 0.62 iu/ml and thereafter gradually decreased non-significantly from groups D, E, F respectively. The lowest value of the hormone significant ($p < 0.05$) was observed from group B that was administered 20mg/kgbw/day only for the duration of the study.




Fig.2A: Effect of Dimethoate and *Beta vulgaris* on Prolactin in female Swiss mice




Fig.2B: Effect of Dimethoate and *Beta vulgaris* on Estradiol in female Swiss mice




Fig.2C: Effect of Dimethoate and *Beta vulgaris* on Progesterone in female Swiss mice




Fig.2D: Effect of Dimethoate and *Beta vulgaris* on Luteinizing hormone in female Swiss mice

Fig.3.3B: Effect of Dimethoate and *Beta vulgaris* on estradiol in female Swiss mice




Fig.2E: Effect of Dimethoate and *Beta vulgaris* on Follicle stimulating hormone in female Swiss mice

3.4 Impact of Dimethoate and *Beta vulgaris* on Litter Size of Female Swiss Mice.

Litter size gradually decrease non significantly from seven in the control and in group C which received 5mg/kgbw/day of Dimethoate along with 250mg/kgbw/day of 100% extract of *B.vulgaris*. Groups D,E,F had litter size of six, five, five respectively. The lowest litter size of two pups was obtained from group B which received 20mg/kgbw/day of Dimethoate only as treatment for the duration of the experiment




Fig.3: Effect of Varying Concentrations Dimethoate and *Beta vulgaris* on Litter Size in Female

4.0 Discussion

The lack of significant reduction in bodyweight gain compared to the control animals of exposed and control animals indicate that Dimethoate is a systemic toxicant.

Endocrine toxicity of pesticides has been well documented and many studies have reported their adverse effect on the reproductive axis of both males [16,17,18,19] females [20]. This study evaluating the impact of *Beta vulgaris* on the reproduction of female mice exposed to various concentrations of Dimethoate showed that Dimethoate influenced the Hypothalamus-Pituitary-Ovarian axis as shown in the concentrations of the reproductive hormones.(Fig.2A-E).The elevated concentration of Prolactin observed in group B that received Dimethoate only, also reflected in the significant ($p<0.01$) reduction of the litter size from seven pups in the control to two pups in group B. All other groups that received Beetroot in addition to varying concentrations of Dimethoate (5,10,15,20mg/kg/bw/day) for the duration of the experiment did show significant increase in the reproductive hormones(Progesterone, Estradiol, Follicle Stimulating Hormone, Luteinizing hormone).These results indicate that *B.vulgaris* exhibited potential anti-oxidant activity as observed in the increase in the Litter size from seven in the control ,seven in group C that received 5mg/kg/bw/day of *B.vulgaris* and 250mg/kg/bw/day, six pups in group D that received 10mg/kg/bw/day and coadministered 250mg/kg/bw/day. The remaining treatment groups E and F received 15 and 20mg/kg/bw/day respectively produced five pups each. The least Litter size was obtained from group B.

Information from the phytochemical components of *B.vulgaris* indicates that the extract has the potential to increase the concentration of estradiol, testosterone because of the presence of boron. The result agrees with the work of [20].The concentrations of Luteinizing hormone and Follicle stimulating (FSH) also increase significantly ($p<0.05$) above levels in the control animals and significantly ($p<0.01$) above the group B animals. This study also showed significant ($p<0.05$) increase in concentration of estrogen, progesterone in the groups that received 250mg/kgbw/day of *B.vulgaris*. It is worthy of note that elevated estrogen, progesterone plays an important role in the physiology, anatomy and perhaps molecular balance of the uterine milieu for fetal implantation and growth.

Conclusion

It is thus deduced that *B. vulgaris* might increase fertility in experimental animals, as well as, enhance the development of the brain in such animals. Dimethoate on the other hand showed significant impact on the reproduction in female experimental animals by reducing the fertility indices, such as reproduction hormones, reduction in fetal implantation leading to low number of offspring (Litter size), inhibition of fertilization and ovulation and embryonic [growth, growth](#) increase in fetal mortality rate and delayed parturition.

Recommendation

Beta vulgaris contains vitamins, minerals and element and has potential as an aphrodisia, the high content of vitamin C, Folic acid and the element boron confer on it a high anti-oxidant activity. It is recommended as refreshing juice for both males and females as it enhances production of reproductive hormones.

References

1. Farag, A.T., Goda, N.F., Shaaban, N.A. and Mansee, A.H. (2007). Effects of oral exposure of synthetic pyrethoid, cypermethrin on the behaviour of F1-progeny in mice. *Reproductive Toxicology*, 23: 560-567.
2. Obulor-Godwin, A.O., Orlu, E.E., Daka, E.R., Orumbo, I.F. & Ebere, N. (2018). Cypermethrin-Induced anti-fertility of the ameliorating effect of lycopene supplementation in female rats. *Journal of Agriculture and Veterinary Science*, 11(3), [57, 57-63](#).
3. Bagchi, M.K., Tsai, M.J., O'Malley, B.W. and Tsai, S.Y. (1992). Analysis of the mechanism of steroid hormone receptor-dependent gene activation in cell-free systems. *Endocrine Review*, 13: 525–535.
4. Health Canada (2011). Proposed Re-Evaluation Decision. Dimethoate. Pest Management Regulatory Agency (PMRA); Health Canada, PRVD2011-12, Ottawa, Ontario.
5. Health Canada (2015). Re-Evaluation Decision Document. Dimethoate. Pest Management Regulatory Agency (PMRA); Health Canada, RVD2015-04, Ottawa, Ontario.
6. Prakasam, A., Sethupathy, S. & Lalitha, S. (2001). Plasma and RBCs antioxidant status in occupational male pesticide sprayers. *Clinica Chimica Acta*, 310(2): 107-112,
7. Farag, A.T., Goda, N.F., Mansee, A.H. and Shaaban, N.A. (2006). Effects of permethrin given before mating on the behavior of F1-generation in mice. *Neurotoxicology*, 27: 421–428.
8. Kaur, R.P., Gupta, V., Christopher, A.F. and Bansal, P. (2015). Potential pathways of pesticide action on erectile function – A contributory factor in male infertility. *Asian Pacific Journal of Reproduction*, 4(4): 322-330.

9. Ben-Abdallah, N. M., Slomianka, L., Vyssotski, A. L., & Lipp, H. P. (2010). Early age-related changes in adult hippocampal neurogenesis in C57 mice. *Neurobiology of aging*, 31(1): 151–1
10. Gore, A.C. (2001). Environmental toxicant effects on neuroendocrine function. *Endocrine-Basingstoke*, 14(2): 235-246.
11. Hamed, S. and Honarvar, M. (2018) Beta vulgaris—A Mini-Review of Traditional Uses in Iran. *Phyto-Chemistry and Pharmacology Current Drug Discovery Technologies*, 15, 1.
12. Costa, D.A., Stahl Hermes, V., de Oliveira Rios, A. and Hickmann Flores, S. (2017). Minimally Processed Beetroot Waste as an Alternative Source to Obtain Functional Ingredients. *Journal of Food Science and Technology*, 54: 2050-2058.
13. Vodnar, D.C., Călinoiu, L.F., Dulf, F.V., Ștefănescu, B.E., Crișan, G. and Socaciu, C. (2017). Identification of the Bioactive Compounds and Antioxidant, Antimutagenic and Antimicrobial Activities of Thermally Processed Agro-Industrial Waste. *Food Chemistry*, 231, 131-140.
14. Whaley-Connell, A., Mc Cullough, P.A. and Sowers, J.R. (2011). The Role of Oxidative Stress in the Metabolic Syndrome. *Reviews in Cardiovascular Medicine*, 12, 21-29.
15. Ninfali, P., Antonini, E., Frati, A., and Scarpa, E.-S. (2017) C-Glycosyl Flavonoids from *Beta vulgaris Cicla* and Betalains from *Beta vulgaris rubra*: Antioxidant, Anticancer and Antiinflammatory Activities—A Review. *Phytother. Res.*, 31: 871– 884.
16. Orlu, E. E. (2014). Deltamethrin-induced Alterations in sperm morphology and spermatogenesis impairment in Adult Sprague-Dawley Rats. *Research Journal of Applied Sciences, Engineering and Technology*, 7 (11), 2324-2331.
17. Obulor, A.O. & Orlu, E.E. (2018b). Evaluation of the therapeutic effect of lycopene on Cypermethrin-Induced reproductive toxicity in rats. *Journal of Environmental Sciences, Toxicology and Food Technology*, 12(3), 10-16.
18. Iboroma, M., Orlu, E.E., Ebere, N. & Obulor, A.O. (2018). Androgenic and antioxidant activity of *Stelleria media* on rat following sub-chronic exposure to Dichlorvos. *Journal of Pharmacy and Biological Sciences*, 13(6), 38-46.
19. Titus, L.K., Orlu, E.E. & Obulor, A.O. (2019). Evaluation of the therapeutic role of Citrullus lanatus and Annona muricata fruit extracts on Cyhalothrin-induced toxicity. *Journal of Advances in Biology and Biotechnology*, 22(4):1-10.
20. El Gamal, A.A., Al-Said, M. M. R., Al-Sohaibani, M., Al-Massarani, S.M., Ahmad, A., Ahmad, A., Hefnawy, M., Al-Yahya M., Basoudan, O.A., Rafatllah, S (2014). Beetroot extract ameliorates Gentamicin-induced Nephrotoxicity associated oxidative stress, inflammation and apoptosis in Rodent models. *Mediators of inflammation*, 2014, 1-12