

## Original Research Article

# Effect of Storage on Dietary fibre rich, Multigrain Cookies

### ABSTRACT

This study focused on the effect of value addition of multigrain cookies with ripe banana peel powder and date powder during storage. Physical and **physiochemical** analysis of value added multigrain cookies were determined as per AOAC and evaluate the effect of these parameters during storage. **In this study firstly optimize the formulation of multigrain flour i.e. wheat, oats, finger millet and barley flour by blending these flour in various ratios, then on the basis of sensory evaluation 70:30 ratio (wheat flour and other flour) was found acceptable.** Incorporation of banana peel powder and date powder in ratio of 2.5 to 10 % with shortening, powdered sugar, milk, vanilla essence and baking powder resulted in nutritionally rich cookies compared to control in the manner of protein, crude fiber and carbohydrate. Data obtained from analysis clearly indicated that significant higher score was observed for T<sub>5</sub> cookies containing 60:30:5:5 ratio of wheat flour: multigrain flour: banana peel powder: date powder. The effect of storage time was sustainably significant on cookies quality parameters, indicating that **cookies were acceptable up to 1-2 months** stored in polyethylene bags without any preservative under ambient conditions.

Comment [L1]: confirmation

Comment [L2]: Correct it according to grammar rules

Comment [L3]: Not suitable with the result and discussion

**Keywords:** Banana peel, Cookies, Dates powder, Dietary fibre, Storage study

### 1. INTRODUCTION

Wheat (*Triticum* spp.) is the second most important winter cereal in India after rice. Bread wheat contributes approximately 95% to total production while another 4% comes from *durum* wheat and *Dicoccumsharein* wheat production remains only 1%. India has witnessed a significant increase in total food grain production to the tune of 233.88 million tonnes with a major contribution of wheat with 80.58 million tonnes (34.5%) during 2008-09 [1]. The wheat flour is major ingredient in chapatti, bread **and bakery products such as cakes, cookies, crackers, doughnuts, sweet rolls, biscuits etc.** Wheat flour is basic ingredient in bakery products due to **their inherited property to form dough and retain gases.**

Comment [L4]: Give the exactly reason

Oats (*Avena sativa*) are excellent sources of different dietary fibre components of linkage (1-3),

(1-4)- $\beta$ -D glucan arabinoxylans and cellulose [2]. Soluble fibre of oats has been reported to reduce elevated blood cholesterol, triglyceride, and glucose levels. Oats also good sources of insoluble fibre functions as a water holding capacity agent and can reduce intestinal transit time when present in adequate amounts in foods [3].

Finger millet (*Eleusine coracana*) also known as *Ragi* is mainly used in India and Africa. *Ragi* is important millet crop in dry hill area of India. It is rich in protein, calcium, phosphorus, iron, fiber and vitamin content [4]. The calcium content is higher than all cereals and iodine content is considered to be highest among all the food grains. Ragi has best quality protein along with the presence of essential amino acids, vitamin A, vitamin B and phosphorus [5].

Banana is one of the most common fruit crops grown in almost all tropical countries, including India. As industrial by-products, peels represent about 30-40 % of fruit weight, this resulted in 200 tons of waste from banana peels. The banana peels waste is normally disposed in municipal landfills, which contribute to the existing environmental problems [6]. The peel is rich in fiber, proteins, essential amino acids, polyunsaturated fatty acids, potassium, and phenolic compounds [7,8]. The fruit fibre has a better quality than other fibre sources due to its high total and soluble fibre content, water and oil holding capacities, and colonic ferment ability, as well as a lower phytic acid and caloric value content. Besides using whole grains, the nutritional enhancement of product could be done by replacing the purified sugar with natural traditionally used sweet products like date (*Phoenix dactylifera*) powder [9]. Various investigations justified the use of date powder as a sweetener in dairy products, however investigation related to its utility in bakery product is merely considered.

."Cookie" is chemically leavened product, also known as "biscuit". Generally the term biscuit is used in the European countries and cookies in the USA. They differ from other baked products like bread and cakes because of having low moisture content, ensure comparatively free from microbial spoilage and confer a long shelf life of the product [10]. Aim of this study was to examine the **physiochemical** properties of value added multigrain cookies and assess the storage stability of cookies.

Comment [L5]: space

## 2. MATERIALS AND METHODS

The experiment was conducted in Department of Processing and Food Engineering, Vaugh Institute of Agricultural Engineering and Technology, SHUATS, Prayagraj, India.

### 2.1 Procurement of raw materials

Good quality of raw materials wheat, **Ragi**, Barley, Oats, sugar, shortening, date, banana fruit etc. required to prepare cookies was purchased from local market of Allahabad. Care was taken the flour was creamy white coloured and free from bean fragment. All necessary hygiene and sanitation was maintained during processing.

Comment [L6]: Confirmation language

Vanilla essence, Ether, Sodium bicarbonate, Ammonium bicarbonate was used for preparation and analysis purpose. All chemicals used were of analytical grade.

## 2.2 Methods

### 2.2.1 Formulation of multigrain flour

Firstly same amount of various flour (**Ragi**, Barley and Oats) were mixed in a bowl as shown in Table 1. Then replace the wheat flour with the multigrain flour in the ratio of 100:0, 80:20, 70:30 and 60:40. Multigrain flour was mixed with the other ingredients, after that dough was kneaded and rested for 10 min. The dough were rolled and chopped with the cookie cutter and baked at 175°C for 15 min. Sensory evaluation will be carried out for superior sample and the best sample will further incorporated with banana peel powder and date powder.

Comment [L7]: confirmation

### 2.2.2 Preparation of value added multigrain Cookies

On the basis of sensory evaluation 70:30 ratio (wheat flour and other flour) was found acceptable. Banana peels powder and date powder in ratio of 2.5 to 10 % with shortening, powdered sugar, milk, vanilla essence and baking powder was incorporated and formulation of value added cookies are shown in Table 1 and 2.

**Table 1. Formulation of Multigrain flour in the cookies**

Sample	Wheat Flour	Multigrain Flour
T <sub>0</sub>	100	0
T <sub>1</sub>	80	20
T <sub>2</sub>	70	30
T <sub>3</sub>	60	40

**Table 2. Formulation of value added cookies**

Sample	Wheat flour	Multigrain flour	Banana peel powder	Date powder
T <sub>4</sub>	65	30	2.5	2.5
T <sub>5</sub>	60	30	5	5
T <sub>6</sub>	50	30	10	10

## 2.3 Physical evaluation of cookies

### **2.3.1 Weight, diameter and thickness**

The weight, diameter and thickness of cookies were calculated as per AACC [11] methods.

### **2.3.2 Spread ratio (D/T)**

Spread ratio of cookies was calculated by AACC [11] method. It was estimate as ratio of diameter to thickness of cookies.

$$\text{Spread ratio} = \frac{\text{Diameter of cookie}}{\text{Thickness of cookie}} \quad \dots \text{Equation 1}$$

### **2.3.3 Volume of cookies (cm<sup>3</sup>)**

Volume of cookies is defined as area of the cookie multiplied by thickness.

$$\text{Volume} = \pi r^2 \times T \quad \dots \text{Equation 2}$$

### **2.3.4 Density of cookies**

After calculating volume, density of cookies is obtained by ratio of weight of volume.

$$\text{Density (g/cm}^3\text{)} = \frac{\text{Mass of sample}}{\text{Volume of sample}} \quad \dots \text{Equation 3}$$

## **2.4 Evaluation of physicochemical composition**

Prepared cookies were evaluated for moisture, crude fat, crude protein, crude fibre, ash content and carbohydrate using standard methods [12].

## **2.5 Statistical Analysis**

Experiments were conducted in triplicate. Data were expressed as the means of these values  $\pm$  the standard deviations (SD). Analysis of variance (ANOVA)- one way ANOVA was used to assess data.


## **3. RESULT AND DISCUSSION**

The value added cookies were stored in low density polyethylene pouches at room temperature and stored for 30 days.

### **3.1 Physical properties of value added multigrain cookies during storage**


**Comment [L8]:** Choose which is better using pictures or tables because of boring expectations.

There was slight decrease in spread ratio of cookies during storage and it decreased from 2.39 to 3.1 at the end of 30 days. Fig.1 shows decrease in spread ratio with increased banana peel powder and date powder, showed that starch polymer molecules are tightly bound with granules and swelling is limited in the cookies with wheat flour when heated. On cooling the starch formed a rigid gel with capacity characteristics of large molecular aggregates and there was a slight decrease in spread ratio during the storage because of the degradation of the starch in cookies during the storage. Similar result was obtained by [13] where spread ratio decreased significantly with increased in proportion of soya flour and wheat sorghum composite biscuit.




**Fig.1 Study of changes in spread ratio of cookies during storage**

The sample shows slight decrease in volume during storage. The decrease in the volume of the banana peel flour and dates powder are due to decrease in the mass of the cookies prepared, the mass decreased due to the loss of protein, fat and other nutrient content of the sample. There was also a decrease in the volume during incorporation of banana peel powder and date powder is due to decrease in spread ratio because they are interrelated to each other as spread ratio is decreasing the volume is also decreasing. Likewise the same result was obtained by Rufeng *et al* [14] in defatted soy flour.



**Fig.2 Study of changes in volume (cm<sup>3</sup>) of cookies during storage**

The decrease in the density is due to the increase in the banana peel powder and date powder because these two contains fat retaining capacity during baking and it is decreasing during the storage because fat, protein and vitamins present in the cookies loss during the storage. Likewise same result was obtained by Rufeng *et al* [14] in defatted soy flour.



**Fig.3 Study of Density of cookies during storage**

### 3.2 Physicochemical properties of value added multigrain cookies during storage

The moisture content of cookies increased with increase in concentration of banana peel powder and date powder and also increased from 2.30 to 3.60 % during storage due to high

water binding capacity of ripe banana peel powder and date powder as well as baking powder which retain higher moisture content in ultimate products the moisture content is also increasing during storage due to the permeability of packaging material. The result for moisture content of cookies was similar with the result obtained by researcher who incorporate potato flour in preparation of the cookies and found increased moisture content of the potato flour based biscuit [15].

**Table 3. Effect on moisture content (%) of cookies during storage**

SAMPLE	0 DAY	10 DAYS	20 DAYS	30 DAYS
T <sub>0</sub>	2.30± 0.83	2.65± 1.66	2.90± 0.94	3.15± 0.48
T <sub>4</sub>	2.49± 0.66	2.85± 0.32	3.10± 0.45	3.60± 0.75
T <sub>5</sub>	2.53± 1.05	2.75± 0.58	3.30± 0.89	3.55± 0.67
T <sub>6</sub>	2.65± 0.87	2.80± 1.12	3.25± 0.62	3.50± 0.49


Table 4 shows the effect of different treatment and storage period on percent fat content of T<sub>0</sub>, T<sub>4</sub>, T<sub>5</sub> and T<sub>6</sub>. The data clearly indicate that there was slight decrease (from 23.76 to 22.18 %) in fat content of all prepared samples. The decrease in fat content in cookies may be attributed to the development of rancidity. The fat deterioration during storage may be the activity of lipase enzyme which split off the fat into free fatty acids and glycogen in the presence of catalyst like moisture, light and heat and there is increase in the fat content during storage is due to fat retaining capacity of banana peel powder and date powder during baking. An increase in the fat content of cookies by [16] was reported and explained to be largely due to the incorporation of defatted mustard flour.

**Table 4. Effect on Fat content (%) of cookies during storage**

SAMPLE	0 DAY	10 DAYS	20 DAYS	30 DAYS
T <sub>0</sub>	22.35± 0.22	22.30± 1.21	22.23± 0.34	22.18± 0.52
T <sub>4</sub>	23.52± 0.56	23.45± 0.79	23.39± 1.08	23.33± 0.84
T <sub>5</sub>	23.61± 0.25	23.59± 0.86	23.25± 0.69	22.25± 0.75
T <sub>6</sub>	23.76± 0.96	23.65± 0.53	23.57± 0.78	23.50± 0.72


Protein content justifies the suitability of banana peel powder and date powder in making nutritionally enhanced cookies. The protein content of cookies found to increase (8.10 to 11.30 %) linearly with increase with the banana peel powder and date powder because banana peel powder and date powder contains good amount of protein. There was a significant decrease in protein content from 7.70 to 11.30 % during storage at the end of 30 days; it is due to

denaturation of protein during storage and processing, non-enzymatic reaction may cause foods deterioration and reduce the shelf life [17]. The result obtained about the protein content of cookies is similar with Padmaja & Jisha [18], incorporating legumes in cassava based composite flour.



**Fig. 4 Effect on protein content (%) of cookies during storage**

Crude fiber content justifies the suitability of banana peel powder and date powder in making nutritionally enhanced dietary fiber rich multigrain cookies. The crude fiber content of cookies found to increase linearly with increase with the amount of banana peel powder and date powder because banana peel powder and date powder contains good amount of crude fiber as well as added multigrain also have a good amount of crude fiber. There was a slight decrease in crude fiber content during 30 days storage. Reactions during processing may affect the dietary fibre and formation of resistant starch is the reason [19]. It was not showing significant difference which proves sustainability of crude fiber content of dietary fiber rich multigrain cookies.



**Fig. 5 Effect on crude fibre content (%) of cookies during storage**

Carbohydrate content of cookies found to decrease linearly with increase with the amount of banana peel powder and date powder because banana peel powder and date powder contains good amount of crude fiber as well as added multigrain also have a good amount of crude fiber, which replaces the carbohydrate which results in decreasing level of carbohydrate as increased level of banana peel powder and date powder. There was a slight decrease in carbohydrate content during 30 days storage; it was not showing significant difference which proves sustainability of dietary fiber rich multigrain cookies.

**Table 5. Effect on carbohydrate content (%) of cookies during storage**

SAMPLE	0 DAY	10 DAYS	20 DAYS	30 DAYS
T <sub>0</sub>	61.50± 0.97	61.47± 0.78	61.47± 0.73	61.20± 1.42
T <sub>4</sub>	42.53± 0.65	42.46± 1.12	42.36± 0.94	42.27± 1.07
T <sub>5</sub>	35.37± 1.34	35.16± 0.64	35.14± 1.05	34.99± 1.18
T <sub>6</sub>	30.25± 1.62	30.11± 0.32	30.19± 0.82	30.10± 0.67

**Comment [L9]:** Write the completeness of the all tables so that it is easy to understand

### CONCLUSION

In the present investigation, it is concluded that the cookies prepared with the incorporation of banana peel powder, date powder and multigrain i.e. (barley, finger millet and oats) with wheat flour, at different concentration levels, which were analyzed for their physicochemical, cooking qualities and sensory acceptability among those seven samples T<sub>5</sub> was found to be more acceptable with respect to mentioned quality parameters. The cookies made from this combination could be stored for 30 days in ambient condition without major quality deterioration.

**Comment [L10]:** Too long sentence

## REFERENCES

Comment [L11]: Write Reference number according to the number in the text

Singh SS. Wheat production in India and future prospects. 8<sup>th</sup> International Wheat Conference, St. Petersburg, Russia: June 1- 4, 2010.

Skendi A, Biliaderis CG, Lazaridou A, and Lzydorczyk, MS. Structure and rheological properties of water soluble  $\beta$ -glucan from oat cultivars of *Avena sativa* and *Avena bysantina*. Journal of Cereal Science. 2003; 38: 15-31.

Anderson J W. Fiber and health: an overview. American Journal of Gastroenterology. 1986; 81: 892-897.

Desai A D, Kulkarni S S, Sahoo A K, Ranveer R C, and Dandge P B. Effect of supplementation of malted ragi flour on nutritional and sensorial quality characteristic of cake. Advance Journal of Food Science and Technology. 2010; 2(1):67-71

Glew R S, Chuang LT, Roberts J L, and Glew R H. Amino acid, fatty acid and mineral content of black finger millet, cultivated on the Jos Plateau of Nigeria, Food Chemistry. 2008; 2:115-18

Schieber A, and Stintzing R C. By-products of plant food processing as a source of functional compounds — recent developments – Review. Trends in Food Science and Technology. 2001; 12(11): 401–413.

Padam B S, Tin H S, Chye F Y, and Abdullah M I. Banana by-products: an under- utilized renewable food biomass with great potential. Journal of Food Science and Technology. 2014; 51(12): 3527-3545.

Gomez-Montano F J, Bolado-Garcia V E, and Blasco-Lopez G. Compositional and antioxidant analysis of peels from different banana varieties (*Musa spp.*) for their possible use in developing enriched flours, *acta Universe*. 2019; 29: e2260.

Habib H M, and Ibrahim W H. Nutritional quality evaluation of eighteen date pit varieties. International Journal of Food Science and Nutrition. 2008; 60(01): 99-111.

Wade P. Biscuit, cookies and crackers: The principles of the craft. Vol. I. Elsevier Applied Sci., London; 1988.

AACC. American Association of Cereal Chemists 7th Edition; 1976.

A.O.A.C. Official methods of analysis of Association of Official Analytical Chemists International 7th ed. Gaithersburg, Method 991.3, Total Dietary Fibre, Enzymatic-Gravimetric Method; 2000.

Mc Watters K H. Cookie baking properties of defatted peanut, soybean and field pea flour, Cereal Chemistry. 1978; 55(6): 853-863.

Rufeng N P, Rao H K, and Laxmidevi N. Preparation and evaluation of high protein biscuit containing whey protein concentrate. J Food sci. Technol. 1995; 44 (5): 532-535.

Mishra R A, and Kalpana R C. Factors in hard flours responsible for reduced cookie spread, Cereal Chem. 2003; 74:330-336.

Tyagi S K, Manikantan M R, Oberoi H S, Kaur G. Effect of mustered flour incorporation on nutritional, textural and organoleptic characteristics of biscuits. J. Food Engineering. 2006; 80: 1043-1050.

Singh R P. Scientific principles of shelf life evaluation of protein biscuits. Journal of Food Science and Technology. 2000; 31: 117-121.

Padmaja T U, and Jisha R. Water absorption and solubility indices of extruded African bread fruit blends. Journal of Food Technology. 2005; 4(1): 64-69.

Nadarajah S, and Mahendran T. Influence of storage conditions on the quality characteristics of wheat-defatted coconut flour biscuits packed in metalized polypropylene. International Journal of Engineering Research and Technology. 2015; 4 (7): 948-951.