

POLYCYCLIC AROMATIC HYDROCARBONS CONCENTRATIONS IN SMOKED AND FRESH CLARIAS GARIEPINUS FROM GOVERNMENT ASSISTED CLUSTERED FISH FARM, IKORODU, LAGOS STATE.

ABSTRACT

Aims: Evaluation of polycyclic aromatic hydrocarbons (PAHs) concentrations was conducted on fresh and smoked *Clarias gariepinus*

Study Design: Fifty numbers of fresh and smoked *Clarias gariepinus* collected from selected clustered farmers and separated into five groups based on their weight range of 200.5g and 655.2g for the analysis.

Place and Duration of Study: The sampled fish were collected from government assisted clustered fish farm estate. The analyses for PAHs were carried out at the central laboratory of Nigeria Institute for Oceanography and Marine Research, Lagos.

Methodology: The laboratory procedures and analyses for the detection of sixteen PAHs in the muscles of fresh and smoked on each five grouped of *C. gariepinus* were carried out using Gas Chromatography (Model HP 6890A).

Results: The significant levels of PHAs concentration were determined using a two-way analysis of variance (SPSS version 25). There is no significant difference in the means concentrations of PAHs in both fresh and smoked fish muscles but significant difference occurs ($p=.05$) in the grouped smoked fish. However, Indeno (1, 2, 3-c, d) Pyrene mean concentrations was higher in both fresh and smoked fish.

Conclusion: PAHs concentrations from both fresh and smoked *C. gariepinus* muscles are below FAO/WHO threshold limit. It is therefore safe for human consumption but gas or electric powered smoking kiln is hereby recommended to prevent further accumulation of PAHs in smoked fish which could be carcinogenic to human consumers.

Keywords: Polycyclic Aromatic Hydrocarbons, *Clarias gariepinus* Evaluation

1. INTRODUCTION

Food toxicity from the harmful substances and the determination of lethal level concentrations to consumers is the mainstream of food safety and standards regulatory agencies all over the world. The toxic substances in food in most cases are usually chemical compounds which are either elemental or complex harmful materials or rather organics toxicants such as polycyclic aromatic hydrocarbons (PAHs)

According to [1] PAHs are a group of organic compounds consisting of two or more fused aromatic rings and a group of lipids soluble compounds that are carcinogenic in nature and their presence have been found in all our bionetworks [2]. The formation of PAHs are known by researchers to be through the natural processes of carbonization and subsequent separations into small amount of persistent organic pollutants that usually occur in our entire biospheres [3], [4], [5] formed from distance and settled elsewhere or formed as a native of a particular place (allochthonous and autochthonous) [6], and it could also be a by-products from the incomplete combustion or burning and organic matter such as tobacco smoke, engine exhausts, petroleum distillates, and coal-derived products, with combustion sources predominating during industrial processing or from natural processes of volcanic eruptions and, various anthropogenic activities [7].

PAHs are dispersed extensively into the atmosphere and this could form another complex compounds with sulfur dioxide, nitrogen oxides, and ozone pollutants under processes known as Photo-Degradation [8]. There are sixteen (16) known PAHs and their sources are but not limited to; coal tar, petroleum fraction, diesel exhaust fumes, spruce needle, tree leaves, grass and plants, vehicle exhaust, processing of organic material, emissions from petroleum refining, incomplete combustion of organic matter, agricultural burning and hazardous waste sites, tobacco smoke, coal tar, soot and certain food products, especially smoked and barbecued foods [9] and [8].




Fig 1: Chemical structure of 16 PAHs (Source: [10])

[11] Classified PAHs as ninth among other chemical compounds causing health issues in humans. The health risk factor associated with PAHs could serve as precursor for incurable diseases such as cancer attacks on consumers of such contaminated food items.

In Nigeria, smoked or barbecued fish is one of the consumer's preferences of eating fish and data shows that out of 30,000 Tons of various freshwater and brackish water fish species produced in Nigeria, about 61% of it are smoked [12]. Fish smoking is one of the prehistoric technologies of fish processing and preservation that has been assisting fishermen, fish farmers and fish mongers for ages. Smoked fish commands high market price due to its aesthetic taste and the availability for consumption during period of fresh fish scarcity and most times, some species of fish are considered as a "status food" when smoked. The traditional fish smoking techniques involve subjecting the entire or gutted fish into smoldering wood or charcoal that comes into direct contact with the fish and this process gives opportunity for the accumulation of PAHs on fish with its attendant carcinogenic effect on consumers [13]. The rate of accumulation of PAHs on smoked fish depends on the techniques, combustion temperature and intensity of smoking used during smoking [14].

Clarias gariepinus, is a cultivable fish species and of great economic importance in Nigeria. The species of fish is widely accepted and most times, it is at the center of meal in some households in Nigeria. It is a

preferred source of animal protein due to its distinctive taste, flavor and tenderization [15]. Smoking of *C. gariepinus* is one of the popular methods of processing and preserving the fish as well as improving the taste qualities of the fish [16].

Government assisted clustered fish farm estate in Ikorodu occupies about 34 hectares of land with 250 numbers of prospective fish farmers that form clustered groups with a production capacity of over 12 tons of both fresh and smoked fish per annum, majorly *Clarias gariepinus*. This annual tonnage of fish production can support in meeting the protein need of about one-third of Lagos population annually. Hence, the need to conduct toxicological study on the fish produced considering the number of people that their health will be at risk if the fish is not evaluated to determine the level toxicants in them and make recommendations to appropriate authority.

2. MATERIALS AND METHODS

Clarias gariepinus with weight range between 200.5g and 655.2g were randomly taken from government assisted clustered fish farm estate, Ikorodu, Lagos State for the analyses of sixteen PAHs. Fifty (50) numbers of *C. gariepinus* were collected from selected clustered farmers and are separated into 5 groups based on their weight range (table. 1)

Table 1: Grouped wet weight of *C. gariepinus* for PAHs determination

GROUP	NUMBER OF FISH	WEIGHT RANGE (g)
1	10	200.5 – 220.1
2	10	315.2 - 325.5
3	10	410.3 – 455.5
4	10	515.4 – 550.3
5	10	620.2 – 655.2

Fish muscles were taken from each 50 fresh *C. gariepinus* and tagged before smoking accordingly to avoid mix-up. Likewise, fifty (50) samples of *C. gariepinus* muscles from the tagged fish were taken after smoking from the same set of fish.

The smoked *C. gariepinus* were subjected to the same smoking temperature between 66^oc and 77^oc for 6 hours in a 1000kg capacity charcoal-powered smoking kiln. From each smoked and fresh *C. gariepinus*, 10g of fish muscle was taken respectively at the ventral axis of the fish girth and the samples collected were conveyed to the laboratory using standard chains of custody involved in PAHs sampling and assessment. The analysis of PAHs concentrations in the muscles of fresh and smoked *C. gariepinus* was carried out at the central laboratory of Nigeria Institute for Oceanography and Marine Research, Lagos.

2.1 Investigated organics toxicant

Nephthalene, Acenaphthalene, Fluorene, Phenanthrene, Anthracene, Fluoranthene - pyrolytic, Pyrene, Benz(a) anthracene, Chrysene, Benzo(b) fluoranthene, Acenaphthene, Benzo [k] Fluoranthene, Benzo [a] pyrene, Benzo [g,h,f] Perylene, Indeno [1,2,3–c,d] and Dibanzo[a,h]Anthracene.

2.1.1 Reagents, Stock Solution Preparation, and Extraction Methods for Fish Muscle Samples

The reagents used in stock solution preparation are; Silica gel, 60-120 mesh, baked at 130^o C for 8 hours, Dichloromethane (hplc-grade), n- Hexane (hplc-grade), Acetone (hplc-grade) and Sodium sulfate, Anhydrous (ACS-grade). Fish samples were shade-dried, and separately homogenized with a blender. A 5g portion of each sample homogenate was separately saponified with 200 ml methanol/KOH (12% KOH in 95% methanol) solution in an ultrasonic bath at 60 °C, for 30 min. The extract was cooled and

filtered, through glass wool, into a separatory funnel. The filtrate was extracted twice with 25 ml hexane. The extract was washed with methanol/water (4:1) mixture, and then concentrated to 1ml with a rotary evaporator. The concentrate was fractionated through a silica gel column, first eluted with 10 ml hexane to collect the aliphatic hydrocarbon fraction, and then with 15 ml methylene chloride to collect the aromatic hydrocarbon fraction. The aromatic hydrocarbon fraction was dried over anhydrous sodium sulfate, and then concentrated to 1 ml with a rotary evaporator, and saved in GC vials for subsequent determination of PAHs.

2.1.1.1 Analysis protocol, GC Set-up conditions for PAHs and quantitation and detection limits

The extracts in the GC vials were transferred to a GC auto-sampler and analyzed by GC/FID following the procedures specified in the appropriate PAHs method. Gas Chromatography model HP 6890A with Column DB 5 MS (30m x 0.25mm x 0.5um) was used in the determination of PAHs. The oven was set to 40°C and hold for 2 min, then ramp 250°C at 25°C/min to 265°C, and at 5°C/min to 300°C. Then hold 4 min, at 25°C. Flame Ionization Detector (FID) was set at 350°C. The Carrier Gas, Hydrogen at 4 ml/min (constant flow Injector: 350°C). The Injection volume: 1 uL, splitless, (hold 2 min). 1µL of the aromatic hydrocarbon fraction was injected into the GC, set-up for the quantitation of PAHs, using the Agilent Chemstation Software. The lowest concentration of Polycyclic Aromatic Hydrocarbons was taken as detection limits that are extractable with good recoveries. The procedures in PAHs determination in smoked and fresh *C. garipenus* is followed with slight modifications by [17]

2.1.1.1.1. Applications and Statistical Analysis

Evaluation of polycyclic aromatic hydrocarbon concentrations in smoked and fresh *C. garipenus* from government assisted clustered fish farm estate and its dietary effect on human health. Statistical significance was assessed using a two-way analysis of variance (ANOVA) and the software used is SPSS version 20. The correlation coefficient and differences between groups were considered significant when $p = .05$ (two tailed)

3. RESULTS AND DISCUSSION

The results of levels of concentrations (µg/kg) of PAHs from GC analyses on fresh and smoked fish muscles are shown in table 2. It indicated that Dibenz (a,h)anthracene had the highest concentration in fresh fish from Group 3 (0.085µg/kg) while smoked fish had the highest concentration of Benzo (g, h, i) Perylene (0.085µg/kg) in group 1. Benzo (a) Pyrene had the lowest concentrations of 0.001µg/kg in both fresh and smoked fish from groups 5 and 1 respectively. Though, Napthalene, Anthracene, Chrysene, Benzo (b) Fluoranthene, Benzo (k)Fluoranthene, and Benzo (a) Pyrene concentrations from fresh and smoked fish in both Group 3 are below the detection level (BDL) while Acenaphthylene concentration was below detection level in Group 1 of fresh fish muscle but detected in all grouped smoked fish muscle. Furthermore, other listed PAHs in Table 2 and their detected concentrations are predominant in all the grouped wet and dried weights of fish. The mean concentrations of PAHs (Σ PAHs) present in smoked fish are higher than that in fresh fish, smoked fish muscles had mean total of 0.0014176µg/kg while fresh fish had mean total of 0.0009496µg/kg but Indeno (1, 2, 3-c, d) Pyrene had the highest mean PAHs concentrations in smoked fish while Dibenz (a, h) Anthracene had the highest mean PAHs concentrations in fresh fish as indicated in table 2. Nevertheless, the mean PAHs concentrations did not increase with increase in the weight of fresh and smoked fish but statistical analysis shows no significant differences within the grouped fresh and smoked fish respectively.

Table 2: showing the concentrations of Polycyclic Aromatic Hydrocarbons in fresh and smoked muscle of *Clarias gariepinus*

S/N	PAHs	PAHs Concentrations in Fresh Fish Muscle						PAHs Concentrations in Smoked Fish Muscle					
		Grp. 1 (µg/kg)	Grp. 2 (µg/kg)	Grp. 3 (µg/kg)	Grp. 4 (µg/kg)	Grp. 5 (µg/kg)	Mean (STD±)	Grp. 1 (µg/kg)	Grp. 2 (µg/kg)	Grp. 3 (µg/kg)	Grp. 4 (µg/kg)	Grp. 5 (µg/kg)	Mean (STD±)
1	Napthalene	0.004	0.004	BDL	0.004	0.004	0.003±0.000 ^b	BDL	0.006	BDL	BDL	0.004	0.002±0.001 ^d
2	Acenaphthylene	BDL	0.003	BDL	0.003	0.005	0.002±0.001 ^b	0.004	0.006	BDL	0.008	0.004	0.004±0.002 ^d
3	Acenaphthene	0.011	0.006	0.011	0.011	0.023	0.013±0.006 ^b	0.044	0.028	0.028	0.024	0.017	0.028±0.010 ^{a,b,c}
4	Fluorene	0.008	0.007	0.008	0.007	0.008	0.008±0.001 ^b	0.011	0.010	0.011	0.011	0.012	0.011±0.001 ^{b,c,d}
5	Anthracene	0.013	0.012	BDL	0.011	0.004	0.008±0.004 ^b	0.006	0.015	0.008	0.016	0.013	0.012±0.004 ^{b,c,d}
6	Phenanthrene	0.004	0.004	0.012	0.004	0.004	0.006±0.004 ^b	0.011	0.006	0.013	0.006	0.003	0.008±0.004 ^{c,d}
7	Fluoranthene	0.009	0.006	0.008	0.006	0.004	0.007±0.002 ^b	0.014	0.008	0.011	0.010	0.007	0.01±0.003 ^{b,c,d}
8	Pyrene	0.004	0.004	BDL	0.004	0.004	0.003±0.000 ^b	0.004	0.006	0.011	0.009	0.005	0.007±0.003 ^{b,c,d}
9	Benz (a) Anthracene	0.010	0.005	0.011	0.007	0.006	0.008±0.003 ^b	0.011	0.013	0.012	0.007	0.006	0.010±0.003 ^{b,c,d}
10	Chrysene	0.005	0.005	BDL	0.005	0.004	0.004±0.001 ^b	0.004	BDL	BDL	BDL	0.004	0.002±.000 ^d
11	Benzo (b)Fluoranthene	0.007	0.006	BDL	0.008	0.008	0.006±0.001 ^b	0.008	0.008	BDL	0.008	0.007	0.006±0.001 ^{c,d}
12	Benzo (k)Fluoranthene	0.007	0.005	BDL	0.006	0.006	0.005±0.001 ^b	0.007	BDL	BDL	BDL	0.005	0.002±0.001 ^d
13	Benzo (a) Pyrene	0.002	0.012	BDL	0.007	0.001	0.004±0.005 ^b	0.001	0.026	BDL	BDL	0.021	0.010±0.013 ^{b,c,d}
14	Dibenz (a, h) Anthracene	0.071	0.026	0.085	0.005	0.042	0.046±0.033 ^a	0.051	0.043	0.071	0.012	0.024	0.040±0.023 ^a
15	Indeno (1,2,3 – c,d)Pyrene	0.015	0.007	0.031	0.005	0.015	0.015±0.010 ^b	0.095	0.006	0.073	0.033	0.015	0.044±0.038 ^a
16	Benzo (g, h, i) Perylene	0.004	0.004	0.027	0.004	0.036	0.015±0.015 ^b	0.085	BDL	0.068	BDL	BDL	0.031±0.012 ^{a,b}
	Total PAHs (µg/kg)	0.174	0.116	0.193	0.097	0.174		0.356	0.181	0.306	0.144	0.147	
	Mean of PAHs (ΣPAHs)	0.011 ^a	0.008 ^a	0.012 ^a	0.007 ^a	0.011 ^a		0.024 ^a	0.014 ^{ab}	0.031 ^{ab}	0.013 ^b	0.010 ^b	
	STD ±	0.018	0.006	0.026	0.002	0.012		0.031	0.012	0.028	0.008	0.007	

* Grp = Group

**[18] Threshold limit =10µg/kg)

*** BDL = below detection level




Fig. 2 shows the combined PAHs concentrations in both fresh and smoked fish.

The overall concentrations of PAHs shows that PAHs in smoked fish is higher than that of fresh fish in Groups 1 to 4, but in group 5, the overall concentrations of PAHs in fresh fish is higher than that of smoked fish as indicated in the multiple bar chart above in Figure 3. The sixteen types of PAHs concentration between and within the grouped fresh fish in the experiment shows that there is no significant difference among the means concentrations while PAHs concentration between and within the grouped smoked fish also shows that there is no significant difference among the mean concentrations of sixteen types of PAHs but the concentration of each of PAHs type in grouped smoked fish showed that there is significant difference among the mean concentration ($p = .05$). Moreover, positive correlation in PAHs concentrations exists between fresh fish and smoked fish. The dependent variable for all the analyses in the experiment is the concentrations of Polycyclic Aromatic Hydrocarbons in both fresh and smoked fish muscles.




Fig. 3 Overall concentrations of PAHs in smoked and fresh fish

Dietary effect of food on human growth, performance and health has been the major concern of the stakeholders saddled with the responsibility of public health, food security and safety. The focus of food growers and food processing industries should be on food items that are consumed to be holistic and devoid of toxicants and food poisoning substances that can cause ill-health in human but with high nutritive values. However, the presence of toxicant in some food items has been identified to cause adverse health effect in human consumers with various clinical manifestations and symptoms [19].

Fresh *C. geriepinus* that was taken from rearing pond from fish farm estate had some levels of PAHs concentrations in their muscle which may probably be gotten from ingestion of fish feed because they were not exposed to direct carbonization during rearing. The contamination of food items could be from the raw ingredients before harvesting or during storage or processing for fish food [20]. Other route by which PAHs can bioaccumulate in the muscle of fresh fish is through PAHs contaminated water that enters into fish pond and permeate through their skin. This process is referred to as "atmospheric deposition" where water is polluted through atmospheric pollutants and such pollutants are detected in fish [21]. Most fish farmers from fish farm estate depends on deep bore-hole water for their source of water and overhead tanks for water storage and the atmospheric carbon can find their way into the storage tank. The levels PAHs concentrations from fresh fish muscle is significantly lower than that of USEPA threshold limit for fish possibly, because the fish are raised under the manageable production system of aquaculture.

The assessment of PAHs concentrations in smoked farmed *C. geriepinus* muscles are also below USEPA threshold limit but, comparatively it has elevated concentrations than that of fresh fish. The elevated concentrations in smoked fish muscle might be due to the charcoal used in smoking the fish. This observation agreed with the scientific publication of [13] on the effect of traditional smoking techniques on quality of smoked fish using charcoal or wood. Moreover, some PAHs elements that are in fresh fish but not detected in the same fish when smoked may perhaps be due to volatility properties of some PAHs types that are initially bioaccumulated in fresh fish but when subjected to heat during smoking, it becomes volatile [22] as cited by [19] on the volatility properties of PAHs under certain heat condition.

All the sixteen (16) PAHs tested were found in both fresh and smoked fish muscles of *C. geriepinus*, although some that were found in fresh fish may not occur in smoked fish and vice versa because farmed *C. geriepinus* are known to be rich in saturated fatty acids, monounsaturated fatty acids, and omega-6 polyunsaturated fatty acids [23] PAHs are described to be lipophilic in nature which gives a strong bounding between the fatty acids deposit and PAHs concentrations in fish [2] and [24]. It is the fatty acids content, properties of the PAHs and methods of rearing the fish and processing it that determines the level of PAHs concentrations in their muscles. However, the weight of the fish plays no significant roles in the concentration of PAHs in their muscles.

In addition, it is sufficient to mention the adverse effect of consuming smoked and fresh fish that have accumulated PAHs in their muscles to human health. The health risk involve in ingesting fish laden with PAHs that is above agency regulatory permissive limit ranges from carcinogenic to complex physiological health issues [14]. The presence of Pyrene and benzo (a) pyrene in fresh and smoked *C. geriepinus* muscles could be fatal to human health when consumed. These two types of PAHs have the ability to undergo numbers of phase transformation and DNA methylation in human when ingested in the food and the concentrations are above threshold limits and when this occur, it could lead to dangerous and unsolicited gene editing, and or genetic mutagen in human as reiterated by [24] on the effect of using different wood in smoking fish and PAHs concentrations.

4. CONCLUSION

PAHs concentrations from both fresh and smoked *C. geriepinus* muscles from the government assisted clustered fish farm are below USEPA threshold limit and therefore safe for human consumption but the long-term effect of bioaccumulation and subsequent bio-magnification of it in consumers over time should not be overlooked. The need to reduce the accumulation of PAHs on fish items during processing, preservation and storage is very important. It is therefore, recommended that periodic sensitization programs from The National Agency for Food and Drug Administration and Control in conjunction with, Standard Organization of Nigeria be embarked on to those in the fish production line from this teaming fish farmers from government assisted fish farm estate on the essentiality of quality assurance and quality control using standard operating procedures in fish production, processing, preservation, and storage.

Government should also endeavor to introduce to the clustered fish farmers the use of gas or electric powered smoking kiln instead of using wood/charcoal that will promote PAHs accumulations on the smoked fish. The food and safety regulatory agencies and extension workers from Ministry of Agriculture should also visit the clustered fish farmers on a regular basis to educate them and promote new technology in fish processing, preservation and storage that will improve the quality of fish produced. Above all, bush burning should be discouraged and the use of environment-friendly materials should be encouraged.

REFERENCES

- [1] Claire E. Bollinger, Monique McCallister, Ryan Clark, Raina Rhoades, Mark Maguire, Russell E. Savage, Yuqin Jiao, Aramandla Ramesh, Darryl B. Hood. Polycyclic Aromatic Hydrocarbons: Implications for Developmental, Molecular, and Behavioral Neurotoxicity, Editor(s): Ramesh C. Gupta, Handbook of Toxicology of Chemical Warfare Agents (Second Edition), Academic Press, Pages 249-265, ISBN 9780128001592; 2015Abdulazeez T. Lawal. Polycyclic aromatic hydrocarbons. A review, *Cogent Environmental Science*, 2017: 3:1, 1339841, DOI: 10.1080/23311843.2017.1339841
- [2] Harvey Ronald G. Polycyclic Aromatic Hydrocarbons. John Wiley & Sons, Inc., ISBN: 978-0-471-18608-3. Pages 682; 1997
- [3] Charis P. Marston, Cliff Pereira, Jennifer Ferguson, Kay Fischer, Olaf Hedstrom, Wan-Mohaiza Dashwood and William M. Baird. Effect of a complex environmental mixture from coal tar containing polycyclic aromatic hydrocarbons (PAH) on the tumor initiation, PAH-DNA binding and metabolic activation of carcinogenic PAH in mouse epidermis. *Carcinogenesis* vol.22 no. 7 pp.1077-1086. Oxford University Press; 2001.
- [4] Deshpande S .S . Handbook of Food Toxicology. New York: Marcel Dekker Inc; 2002.
- [5] Carlos Magno Oliveira Tadeu, Luciana Pena Mello Brandão, José Fernandes Bezerra-Neto and Diego Pujoni and F.A.R. Barbosa. Photodegradation of autochthonous and allochthonous dissolved organic matter in a natural tropical lake. *Limnologica - Ecology and Management of Inland Waters*. 87(4):125846. DOI: 10.1016/j.limno.2020.125846.
- [6] Chuangshou Wu, Changchen Ji, Benwei Shi, Yaping Wang, Jianhua Gao, Yang Yang, Jinbin Mu. The impact of climate change and human activities on streamflow and sediment load in the Pearl River basin. 2019., *International Journal of Sediment Research*, Volume 34, Issue 4, Pages 307-321
- [7] Patel AB, Shaikh S, Jain KR, Desai C, Madamwar D. Polycyclic Aromatic Hydrocarbons: Sources, Toxicity, and Remediation Approaches. *Front Microbiol*. 11:562813. doi: 10.3389/fmicb.2020.562813. PMID: 33224110; PMCID: PMC7674206.
- [8] Yang Q, Chen H, Li B. Polycyclic Aromatic Hydrocarbons (PAHs) in Indoor Dusts of

- Guizhou, Southwest of China: Status, Sources and Potential Human Health Risk. 2015, *PLoS ONE* 10(2): e0118141. doi:10.1371/journal.pone.0118141
- [9] Masato Honda and Nobuo Suzuki .Toxicities of Polycyclic Aromatic Hydrocarbons for Aquatic Animals. 2020, *Int. J. Environ. Res. Public Health*. 17, 1363;doi:10.3390/ijerph17041363
- [10] Ifegwu O C, and Anyakora C. Polycyclic Aromatic Hydrocarbons: Part I. Exposure. *Adv Clin Chem*. 2015;72:277-304. doi: 10.1016.. PMID: 26471085.
- [11] Uzomah, Abimbola, Anne-Katrine Lundebye, Marian Kjellevoid, Fubara A. Chuku, and Oluwafemi A. Stephen.A Review of Chemical Contaminants in Marine and Fresh Water Fish in Nigeria. . 2021, *Foods* 10, no. 9: 2013. <https://doi.org/10.3390/foods10092013>
- [12] Abou-Arab, N. & Abd-El-Samea, M. & Malhat, Farag & El-TaHER, S. Detection of some potential carcinogenic polycyclic aromatic hydrocarbons in smoked fish. 2014, *Global Journal of Agriculture and Food Safety Sciences*. 1. 346-356.
- [13] Yusuf, K. A., Ezechukwu, L. N., Fakoya, K. A., Akintola, S. L., Agboola, J. I., Omoleye, T. O. Influence of fish smoking methods on polycyclic aromatic hydrocarbons content and possible risks to human health. 2015, *African Journal of Food Science*, 9(3), 126-135.
- [14] Osibona, Adesola & Kusemiju, Kola & Akande, G.R.Proximate composition and fatty acids profile of the African Catfish *Clarias gariepinus* . 2009, *Acta Satech*. 3. 85-89.
- [15] Olayemi Folorunsho Foline, Adedayo Majekodunmi Rachael, Bamishaiye Eunice Iyabo and Awagu Emenike Fidelis. Proximate composition of catfish (*Clarias gariepinus*) smoked in Nigerian stored products research institute (NSPRI): Developed kiln. 2011, *International Journal of Fisheries and Aquaculture* Vol. 3(5), pp. 96-98,
- [16] Mohammadi, Abdorreza & ghasemzadeh-mohammadi, Vahid & Haratian, Parivash & Khaksar, R. & Chaichi, Maryam. Determination of polycyclic aromatic hydrocarbons in smoked fish samples by a new microextraction technique and method optimisation using response surface methodology. 2013, *Food chemistry*. 141. 2459-65. 10.1016/j.foodchem.2013.05.065.
- [17] USEPA (Environmental Protection Agency). Polycyclic aromatic hydrocarbons (PAHs) EPA fact sheet. Washington (DC): National Center for Environmental Assessment, Office of Research and Development. 2008.
- [18] Hussein I.A.S., Mona S.M.M. A review on polycyclic aromatic hydrocarbons: source, environmental impact, effect on human health and remediation. 2016, *Egyptian Journal of Petroleum*. 25: 107-123. [DOI: 10.1016/j.ejpe.2015.03.011]
- [19] Wang, Chunhui & Wu, Shaohua & ZHOU, Shenglu & SHI, Yaxing & SONG, Jing. Characteristics and Source Identification of Polycyclic Aromatic Hydrocarbons (PAHs) in Urban Soils: A Review. 2017, *Pedosphere*. 27. 17-26. 10.1016/S1002-0160(17)60293-5
- [20] Norton, Stephen & Wilson, Tiffany & Handley, Michael & Osterberg, Erich. Atmospheric deposition of cadmium in the northeastern USA. *Applied Geochemistry - APPL GEOCHEM*. 22. 1217-1222. 10.1016/j.apgeochem.2007.03.014. 2007.
- [21] Feng You and Tertry F Bidleman Influence of volatility on the collection of polycyclic aromatic hydrocarbon (PAH) vapor with polyurethane foam. 1984, *Environmental Science and Technology*. 18(5), 330-333.
- [22] Taiwo, O.E., Usman, K., Ogono, T.H. and Osoniyi, R.O Proximate and Lipid Profile Analysis of Cultured and Wild African Catfish, *Clarias gariepinus* [Burchell]. 2014, *Ife Journal of Science* vol. 16, no.1
- [23] Ezike, Christopher. Assessment of Polycyclic Aromatic Hydrocarbons (Pahs) in Hardwood, Palmwood and Softwood - Smoked Fish. 2018, *International Journal of Ecotoxicology and Ecobiology*. 2. 10.11648/j.ijee.20170204.17.