

Canola yield and water productivity as affected by irrigation, tillage and nitrogen levels in northwest India

Comment [AC1]: Suggested title: Effect of tillage and nitrogen levels on yield and productivity in *Brassica napus* in North East India.

ABSTRACT

Depleting ground water resources calls for the need of diversification to low water requiring crops and adoption of water saving techniques. To evaluate individual and interactive effects of deep tillage, irrigation and nitrogen rates on yield and water productivity of canola (*Brassica napus*). Treatments included combinations of two tillage systems viz; deep tillage (DT) and conventional tillage (CT), with three irrigation regimes viz; no (I_0), one (I_1) and two irrigations (I_2) in main plots and four nitrogen rates viz; 0 (N_0), 50 (N_{50}), 75 (N_{75}) and 100 (N_{100}) kg ha⁻¹ in sub plots with three replications. Maximum rooting depth was observed with irrigation and 100 kg N ha⁻¹. Root mass density in upper 60 cm soil depth was higher under irrigated plots whereas below 60 cm, it was higher under I_0 . Higher root density was recorded under DT and N_{100} plots. Dry matter accumulation significantly increased with irrigation, tillage and N application. Seed yield significantly increased under DT (10%) and I_2 (26.2% over I_0) treatment. Water productivity improved with DT and N_{100} . Oil yield and N uptake increased under DTI_2N_{100} . Higher nitrogen rates at low irrigation frequency resulted in yield similar to low nitrogen rates at higher irrigation frequency. While yield produced under DT with one post sowing irrigation was equivalent to that produced under CT with two irrigations. The results suggest the saving of irrigation water and yield optimisation with high N rate and deep tillage in canola.

Keywords: Irrigation, nitrogen rates, oilseed rape, root growth, tillage, water productivity

Comment [AC2]: The abstract must include the objective of the research, a brief description of the methodology and the results. Not only mention the results but their impact at a biological and agronomic level.

Introduction

Due to high productivity and profitability, rice-wheat (R-W) cropping system is dominant in the alluvial tract of Indo-gangetic plains. High water demand and low water productivity of conventional irrigated R-W system has led to the depletion of surface and ground waters. Hence, it calls for diversification to low water requiring crops and implementation of water saving techniques. Oilseed rape provides management options for irrigators seeking to reduce irrigation requirements and diversification and/or to reduce input costs due to low water requirements (25–35 cm, Rathore *et al* 2017).

Comment [AC3]: The keywords should be different from the title and more in line with the nature of the work. Authors are suggested to consult international thesauri for their selection.

India is the world's largest importer of edible oils (FAO 2019). Improvement in yield with the judicious utilization of available resources is required to meet the country's demand. Development, water use and yield of oilseed crops are inter-related. An encouraging effect of irrigating rapeseed mustard at critical stages has been observed (Konwar *et al.* 2019). However, frequent irrigation though sometimes is necessary for yield maximization but it usually lowers the water use efficiency because soil moisture is lost through evaporation from moist soil surface, thus increasing consumptive use (Shivran *et al.* 2018). An efficient irrigation water management in brassica can affect seed and oil content enormously and also the response to

Comment [AC4]: Try to include more recent references.

other applied inputs (Rathore *et al.* 2019). Therefore, there is a need to make proper irrigation scheduling which will provide irrigation at critical growth stages matching the crop evapotranspiration (Kar *et al.* 2005).

Formatted: Highlight

Canola is considered as nutrient exhaustive crop with higher nitrogen (N) requirement than cereals and removes a higher amount of nitrogen until flowering with the relatively lower amount taken during reproductive phase (Rathke *et al.* 2005). The seed yield in canola depicts positive response to increasing rates of fertilizer N (White *et al.* 2015; Ma and Herath 2016) as high as 180 kg N ha⁻¹ (Brandt *et al.* 2002). But excessive use of N fertilizers may promote vegetative growth in the plant at the expense of reproductive growth along with environmental pollution (Akmal *et al.* 2014) and lower nitrogen use efficiency (Chamoro *et al.* 2002). Water and nitrogen exhibit interaction effects on rapeseed-mustard growth and yield (Mandal *et al.* 2010) and their economic analysis is helpful to enhance water and N use efficiency. Thus, it calls for optimization of nitrogen application with an appropriate irrigation schedule to meet the crop necessities and higher yield. Coarse-textured soils show low water holding capacity, poor fertility and rapid development of mechanical resistance to roots leading to nutrient and water stress in growing crops. The degree and duration of nutrient and water stress can be reduced by synchronizing the active root zone with a soil zone containing nutrients and water- CITES.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Tillage is the practice for managing water content (Sharma *et al.* 1990) and control the hydrothermal regime of soil in the root zone (Moroizumi and Horino 2002) through its effects on the shape, size and continuity of soil pores. Deep tillage lowers the mechanical resistance, thus favouring root growth (Arora *et al.* 2011). Several studies have shown the benefits of deep tillage, irrigation and nitrogen fertilizer as individual factors and little information is available on the interactive effects of these three factors especially on canola crop. It was hypothesized that interaction of irrigation, tillage and nitrogen may lead to better resource use efficiency along with yield. Thus, the present study was carried out with objective to evaluate the individual as well as interactive effects of irrigation regimes, tillage and nitrogen rates on yield and water productivity of canola.

Formatted: Highlight

Comment [AC5]: ?

Formatted: Highlight

Formatted: Font: Italic, Highlight

Formatted: Highlight

Materials and methods

Site description

A multi-factor study was conducted at the Punjab Agricultural University; Ludhiana situated at 30°54' N latitude and 75°48' E longitude at a height of 247 m above mean sea level

Comment [AC6]: Complement the previous studies considering the international and national context, the species, where they were carried out and their main conclusions. Consult articles from no more than five years ago. Finish with the objective of the study plus its impact at a biological and agronomic level.

during 2017-19 *rabi* seasons. Important soil physical and chemical properties and weather information of the experimental site are given in Table 1 and Table 2, respectively. Total rainfall during the two growing seasons was 76.0mm (2017-18) and 171.6mm (2018-19). Pan evaporation during cropping season was less than the normal during both the years. Mean maximum air temperature varied between 18.5-35.8°C during different growing seasons as against the normal values of 18.2-34.4°C; while mean minimum temperature varied from 5.5-19.9°C compared to the normal value of 5.6-17.1°C, respectively. CITE.

Vegetal material

Comment [AC7]: Include the main characteristics of the plant material used.

Formatted: Font: Bold

Formatted: Font: Bold

Treatments

Combinations of irrigation regimes and tillage systems as main plots and nitrogen rates as subplots were evaluated in a factorial split-plot design with three replications during two year extensive field trials. Tillage included conventional tillage (CT) - two discs, two cultivators followed by planking operation and deep tillage (DT) - sub-soiling/chiseling ploughed up to 45 cm deep and 50 cm apart followed by CT. Irrigation regimes comprised of no post sowing irrigation (I_0), one irrigation (I_1); at 4 weeks after sowing (WAS) and two irrigations (I_2); one at 4 WAS and second in December end or first week of January. Four different N fertilizer rates viz., 0 (N_0), 50 (N_{50}), 75 (N_{75}) and 100 kg ha⁻¹ (N_{100}) were applied to canola crop. The gross plot size and the net plot size were 3.9 x 3.3 m² and 3.6 x 3.0 m², respectively. The experiment was conducted on the same location and treatments were imposed on same plots in both the years of study.

Crop management

After harvest of preceding maize crop, the field with DT plots were deep tilled (sub-soiled) with tractor drawn chiseler in the first week of October and then, the whole field was ploughed twice with a disc harrow. Heavy pre-sowing irrigation (about 10 cm) was applied to ensure adequate moisture in the soil profile for seed germination. The field was then prepared by giving two cultivations with a tractor drawn cultivator followed by planking at proper moisture condition to obtain a fine seed bed. Canola (GSC 7) was sown with seed rate of 3.75 kg ha⁻¹ and row spacing of 0.45 m in 4th and 3rd week of October 2017 and 2018, respectively. The whole amount of phosphorus (30 kg P₂O₅ ha⁻¹ as single super phosphate) and potassium (15 kg K₂O ha⁻¹ as muriate of potash) was applied at sowing. In plots with I_0 irrigation regime, the full dose of

nitrogen fertilizer (as urea) was applied at the time of sowing as per treatment while in plots with other irrigation regimes, 50 per cent of N as per treatment was applied at sowing and remaining dose of N was applied prior to first irrigation. In N_0 plots, no nitrogen was applied. First irrigation was applied in I_1 and I_2 irrigation regimes on December 2, 2017 and November 16, 2018 growing seasons, respectively. In I_2 regime, second irrigation was applied on December 27, 2017 and January 11, 2019. Parshall flume was used to apply 70 mm of water as flood irrigation. Extra plants were uprooted manually to maintain plant to plant spacing of 0.1-0.12 m within row. Weeds were controlled by hand weeding, and the crop was protected against hairy and cabbage caterpillar by spraying rogor 30 EC @ 1.0 kg ha⁻¹. Pesticide Actara 25 WG @ 0.1 kg ha⁻¹ in 250 liters of water was sprayed for the control of aphid. The crop was harvested manually in the first week of April.

Measurements

Soil penetration resistance expressed as cone index (CI) was measured with a digital cone penetrometer (CP40II; Rimik Electronics, RFM Australia) down to 0.60 m depth from different sites in CT and DT plots. Determining root growth and root mass density (RMD), soil cores were sampled at flowering stage at 0.15 m depth increments down to 1.80 m soil depth with 0.05 m diameter auger centered at 0.075 m away from plant base (Gajri *et al.* 1994). Roots from each sample were washed in net cloth, cleaned, dried at 65°C and weighed. Dry matter accumulation (DMA) was recorded at 60, 110 and 145 DAS from 0.5 m row length from the second outermost row on either side of each plot. The samples were air dried first and later in an oven at 65±2°C till constant weight. The data of DMA was computed and expressed in t ha⁻¹. Total water use based on irrigation water, rainfall and profile water use was computed during both cropping seasons. Water productivity of canola was calculated as ratio of seed yield to total water use. Seed and stover yield was calculated from area of 6.75 m² per plot. N content was determined by modified micro-Kjeldahl method (Piper 1966) and total N uptake was worked out by multiplying percent N content and yield. Oil yield was calculated by multiplying the oil content (determined by NMR Analyser) in the seed sample of each treatment with its respective seed yield and expressed in t ha⁻¹.

Treatment effects on various parameters were tested for their statistical significance using ANOVA for a factorial split-plot design and comparisons were made at 5% level of significance. Analysis of variance was carried out for various parameters using computer programme CPCS 1

(Cheema and Singh 1991).

Results

Soil penetration resistance

Deep tillage caused a reduction in penetration resistance of soil in the tilled zone and down below (Figure 1). The mean cone index in DT plots and CT plots was 0.4 and 0.6, 1.4 (2.0), 1.6 (2.7), 2.0 (3.0), 2.8 (3.2) and 3.2 (3.2) MPa in 0-0.1, 0.1-0.2, 0.2-0.3, 0.3-0.4, 0.4-0.5 and 0.5-0.6 m of soil depth, respectively.

Rooting depth and density

Roots in canola plants extended up to 1.8 m depth in irrigated plots; while limiting to 1.5 m in I_0 plots. Rooting depth increased with an increase in nitrogen dose as in I_0 plots, N_{100} and N_{75} application forced roots down to 1.5 m deep against 1.2 m obtained in N_{50} and N_0 . Likewise in I_1 plots, roots were obtained up to 1.5 m depth under N_0 however with the application of N fertilizer, roots in canola plants extended to 1.8 m. Tillage, irrigation and nitrogen had substantial effects on root proliferation in both the seasons. Root mass density (RMD) was higher in DT as compared to CT irrespective of the irrigation regimes during both years (Figure 2 and 3). Differences in RMD of DT and CT in the top 0.3 m soil depth was minimal; thereafter difference increased with maximum in 0.6 to 0.9 m soil layer and became almost equal in layers below 1.2 m. This ensured water and nutrient uptake by the root in deep tilled soil. Under different irrigation regimes, the surface layer (0-0.15 m) possessed higher root mass density during both the years (Figure 2 and 3). Root mass density increased with an increase in irrigation frequency up to 0.6 m soil layer. While with increase in soil depth from 0.6 m in I_0 revealed 2.8% and 5.4% higher RMD in comparison to both I_1 and I_2 , respectively. Fertilizer N application also affected RMD with the higher impact of 100 kg N ha⁻¹ as compared to lower N doses in all irrigation regimes (Figure 2 and 3). Increasing the soil depth eventually led to minimal difference in RMD when assessed with varied N doses where highest difference in the root density was noticed in the surface soil layer (0-0.15 m).

Dry matter accumulation

Increase in irrigation frequency resulted in higher dry matter production (Table 3) at different crop growth stages. Two irrigations produced significantly high dry matter by 0.39 and 2.00 t ha⁻¹ over one irrigation and 1.35 and 3.78 t ha⁻¹ over no irrigation, respectively, at 110 and 145 DAS. Tillage exerted a significant effect on dry matter accumulation of canola crop at all

growth stages except 60 DAS. Application of nitrogen also improved DMA significantly at all growth stages. At 145 DAS, N100 treatment resulted in an increase in dry matter by 3.34 t ha⁻¹ over N₀.

Seed and stover yield

The individual effects of irrigation, tillage and nitrogen rates on seed yield of canola were substantial in both cropping seasons but interaction was found non-significant (Table 4). Seed yield recorded during second rabi season (2018-19; 1.59 t ha⁻¹) was higher than that observed during first season (2017-18; 1.36 t ha⁻¹) attributed to favorable weather conditions. Averaged over both seasons, two irrigations recorded 10.8 and 26% higher mean seed yield than one irrigation (1.48 t ha⁻¹) and no post sowing irrigation (1.30 t ha⁻¹), respectively. Deep tilled plots registered a mean increase of 10% in seed yield over CT. Increase in N fertilizer dose resulted in significant improvement in seed yield in canola with mean yield varying from 1.04-1.81 t ha⁻¹. Though the interactive effect of factors was non-significant but numerical data showed that deep tillage with one irrigation (2.04 t ha⁻¹) produced at par yield to conventional tillage with two irrigations (2.02 t ha⁻¹) in N₁₀₀ regime in 2018-19. In conventionally tilled plots, two irrigations with 75 kg N ha⁻¹ (1.83 t ha⁻¹) gave comparable yield to one irrigation with 100 kg N ha⁻¹ (1.88 t ha⁻¹), thus saving either irrigation or N fertilizer. Similar results were obtained in plants with deep tillage.

Stover yield followed the similar trend as seed yield (Table 4). Mean increase by 14% and 9.1% was observed with increase in irrigation frequency from I₀ to I₁ and I₁ to I₂, respectively. On an average, stover yield of 5.76 t ha⁻¹ was obtained with deep tillage which was significantly higher than conventional tillage (5.32 t ha⁻¹). Fertilizer N treatment N₁₀₀ resulted in highest stover yield (6.69 t ha⁻¹) followed by N₇₅ (6.07 t ha⁻¹), N₅₀ (5.43 t ha⁻¹) and N₀ (3.98 t ha⁻¹).

Total water use (TWU) and water productivity (WP)

Total water use was higher in second rabi season (2018-19) owing to higher rainfall during the cropping season (Table 5). Total water use increased with increase in irrigation frequency as well as N dose. Deep tilled plots recorded 2.4% higher TWU against conventionally tilled plots. During 2018-19 cropping season, tillage influenced irrigation and N effects on total water use by the crop (Table 5). The results suggested that two irrigations alone enhanced water use by 103 mm over 298 mm in I₀N₀, while 100 kg N ha⁻¹ increased it by 11 mm, and their

combinations increased water use by 120 mm under CT plots. A corresponding increase in water use with these treatments in DT was 111, 25 and 131 mm over 308 mm in I₀N₀. This implies that deep tillage enhanced the irrigation and N effects on total water use. Water productivity was also significantly affected by irrigation, tillage and N fertilizer (Table 5). WP declined with the number of irrigations during both the cropping seasons. A reduction of 11% was observed in water productivity with two irrigations as compared to no irrigation. Deep tillage resulted in 4.49 kg ha⁻¹ mm⁻¹ of WP which was higher than conventional tillage by 7.2 percent. Nitrogen application influenced WP positively with highest value obtained with N₁₀₀ treatment.

Total N uptake

Superior total N uptake was obtained during second growing season (81.6 kg ha⁻¹) as compared to the first season (79.0 kg ha⁻¹) (Table 6). Two irrigations (I₂) resulted in a significantly higher mean total N uptake of 85.3 kg ha⁻¹ as compared to I₁ (81.0 kg ha⁻¹) and I₀ (74.6 kg ha⁻¹). Deep tilled plots resulted in an increment of 13.3 % in total N uptake of canola plants in contrast to conventionally tilled plots. The maximum value of total N uptake was recorded under N₁₀₀ treatment (105.6 kg ha⁻¹) followed by N₇₅ (89.4 kg ha⁻¹), N₅₀ (75.6 kg ha⁻¹) and N₀ (50.6 kg ha⁻¹). The interactive effect showed that total N uptake recorded under DTN₅₀ treatment was statistically at par to that under CTN₇₅ treatment.

Oil yield

Irrigation frequency increased oil yield with maximum mean value with two irrigations (0.67 t ha⁻¹) followed by one irrigation (0.61 t ha⁻¹) and no irrigation (0.53 t ha⁻¹) (Table 6). Deep tillage resulted in a significantly higher oil yield by 10.5 % ~~than~~ than conventional tillage (0.57 t ha⁻¹). Oil yield significantly increased with successive increments of N doses up to N₁₀₀.

Discussion

Effects of Tillage on soil penetration resistance

When compared to conventional tillage, deep tillage resulted in a significant reduction in soil mechanical resistance in loamy sand and sandy loam soils under soybeans (Arora *et al.* 2011). This reduction in soil mechanical resistance is attributed to soil loosening associated with tillage. Kaur and Arora (2019) also reported that the mean cone index (CI) was 0.6, 1.4, 2.7 MPa in DT plots against 1.4, 3.0 and 3.4 MPa in CT plots for 0-0.2, 0.2-0.4 and 0.4-0.6 m soil depth in maize crop indicating that DT reduced soil penetration resistance.

Comment [AC8]: The same order must be kept in the subheadings of methodology, results and discussion.

Formatted: Highlight

Comment [AC9]: The discussion should delve into the biological and agronomic aspect of the results, clearly showing their impact. Two or three quotes are not enough to support the arguments presented.

Effects of tillage, irrigation and N levels on root growth

Root growth is determined by both plant genetics and soil parameters, and it is highly adaptable to changing environmental conditions (Hodge 2006; Lynch 2011). Higher RMD under DT plots in deeper layers may be attributed to lower soil penetration resistance (Figure 1) by deep ploughing. These results demonstrated that sub-soiling reduces the root distribution at the surface soil and promotes root growth in the deeper layer, thus alleviating root crowding and competition in the topsoil while promoting root growth in the deeper soil layer to improve water and nutrient utilization. These findings are consistent with the studies of Sun *et al.* (2017) and Kaur and Arora (2019) in maize. Caia *et al.* (2014) also reported that loosening of subsoil layer with deep tillage up to 0.5 m, increased root length, surface area, dry weight, diameter and the proportion of roots in the 0.4-0.8 m soil layer.

Formatted: Highlight

Comment [AC10]: Comparative studies must clearly show the context, international or national, that is, include where they were carried out and their direct articulation with this study.

Our results endorse earlier reports on root responses of *Brassica* species to irrigation (Chandra *et al.* 2018). Under restricted irrigation, relatively drier upper soil profile might causes roots to penetrate in soil faster to lower depths in search of moisture; thus increasing root density in deeper layers (Sarkar and Kar 2000).

Formatted: Highlight

The N fertilizer application rate is one of the fundamental components of crop management systems that can regulate the root distribution and certain root traits. An increase in rooting density with increase in N-levels might be due to favorable effect of nitrogen on above-ground plant biomass (Table 3) that also encouraged root growth. Results endorse the findings of Marcinkevičienė *et al.* (2013), significantly higher root biomass of rape in 0-0.1 m soil layer with mineral fertilization than no fertilization. Plant roots develop more intensively in the upper soil layer with fertilizer application while penetrate deeper without fertilization to contact more soil volume in attempt to increase N uptake. Increased root growth with incremented N rates was also reported by Beard *et al.* (2018).

Formatted: Highlight

Effects of tillage, irrigation and N levels on DMA and yield

Mishra *et al.* (2019) also recorded the highest DMA with three irrigations followed by two, one and lowest under no irrigation in mustard. Budzynski *et al.* (2000) also reported higher dry matter accumulation under deep tillage. Sub soil tillage to 0.3-0.45 m soil layer improves soil physical behavior and reduces soil mechanical resistance to root penetration. Importantly, the above-ground stability of plant is enhanced by a well-developed root system (Qamar *et al.* 2013). Sharma *et al.* (2007) observed an increase in dry matter production of mustard with the increase

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

in nitrogen rate and highest DMA was recorded at 80 kg N ha⁻¹. This might be due to higher photosynthesis and translocation of assimilates toward reproductive structure owing to adequate soil moisture.

The results were in conformity with Ray *et al.* (2015) and Shivran *et al.* (2018). Deep tillage provided better root growth and moisture extraction that helped the crop to initially develop adequate source (as reflected by high above-ground biomass) as compared to CT. Deep ploughing significantly increased seed yield of mustard over conventional tillage (Pal and Phogat 2005). Our results corroborate with the findings of Ali *et al.* (2019) in canola. This increase is attributed to enhanced availability of moisture which led to a better nutritional environment at the critical crop growth stages, which in turn resulted in better vegetative growth. These results endorse the findings of Konwar *et al.* (2019). The increase for stover yield in mustard was 30.3% and 49.4% by 80 kg N ha⁻¹ over 40 kg N ha⁻¹ and control, respectively (Kumar *et al.* 2018). Tyagi and Upadhyay (2017) concluded that irrigation frequency increased the consumptive water use considerably; 33.1% and 8.3% increase over no-post sowing irrigation with two and one irrigation, respectively. This was expected because irrigation increased the available water in the soil profile and this facilitated more loss of water through evapotranspiration as compared to no irrigation. Similar results were reported by Konwar *et al.* (2019).

Effects of tillage, irrigation and N levels on TWU and Water Productivity

Tyagi and Upadhyay (2017) concluded that irrigation frequency increased the consumptive water use (CU) considerably; 33.1% and 8.34% increase over no-post sowing irrigation with two and one irrigation, respectively. This was expected because irrigation increased the available water in the soil profile and this facilitated more loss of water through evapotranspiration as compared to no irrigation. Similar results were reported by Konwar *et al.* (2019). This implies that deep tillage enhanced the irrigation and N effects on total water use. The results were in conformity with the findings of Kaur and Arora (2019).

A decrease in WP with the successive increase in the number of irrigations is due to the greater expense of water by evapo-transpiration without a proportionate increase in seed yield. While, under one irrigation, the crop developed a deep root system and utilized some moisture from deeper soil layers than frequent irrigation and achieved higher yield per unit of water which in turn resulted in higher water productivity. Tyagi and Upadhyay (2017) recorded significantly higher WP with the application of one irrigation followed by 2 irrigations and no-post sowing

irrigation. Tao *et al.* (2015) reported that sub soil tillage in spring maize significantly increased mean WP_{ET} by 14 percent on a sandy loam soil in China. Similar results were also reported by Kaur and Arora (2019). At a specific irrigation level, greater evapotranspiration in fertilized plots was primarily associated with stimulating crop growth and increased DMA with greater interception of solar radiation (Mandal and Sinha 2004) as well as increased root biomass. Thus, nutrient application positively influenced WP. The greater increase in seed yield in N_{100} , N_{75} and N_{50} over N_0 and relatively less increase of the corresponding ET have evidently resulted in significantly higher WP, particularly in the case of application of the highest N rate (100 kg N ha^{-1}).

Formatted: Highlight

Effects of tillage, irrigation and N levels on total N uptake and oil yield

Rajput (2017) reported that the highest total nitrogen uptake by Indian mustard was observed with 120 kg N ha^{-1} , which was significantly superior over 0, 40 and 80 kg N ha^{-1} . Pal *et al.* (2009) concluded that the oil yield of mustard can be increased by 41 percent with deep ploughing as compared to conventional tillage. These results endorse early reports of N effects on oil yield in canola (Sukirtee *et al.* 2018).

Conclusion

This study demonstrated that two irrigation and higher nitrogen rate (100 kg N ha^{-1}) along with deep tillage significantly increased the growth, yield and nitrogen uptake of canola oilseed rape. Two irrigations produced 10.8 percent higher mean grain yield over one irrigation and 26.2 percent over no post sowing irrigation. Deep tillage incremented seed yield by 10 percent over conventional tillage. Seed yield significantly increased up to N_{100} with the highest seed yield production of 1.81 t ha^{-1} among nitrogen rates averaged over two years. Water productivity improved with deep tillage and nitrogen dose of 100 kg ha^{-1} . Higher nitrogen rates at low irrigation frequency resulted in yield similar to low nitrogen rates at higher irrigation frequency. Deep tillage with one irrigation produced similar yield as under conventional tillage with two irrigations suggesting saving of irrigation water.

Comment [AC11]: The conclusions should not be a summary of the main results.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author, upon reasonable request.

References

Akmal M, Kiran U, Ali A, Abdin MZ (2014) Enhanced nitrogen assimilation in transgenic

- mustard (*Brassica juncea* L.) over expressing high affinity sulphate transporter gene. *Indian Journal of Biotechnology* **13**, 381-387.
- Ali W, Anwar S, Khan MH, Iqbal J, Kamal A, Ahmad Z (2019) Canola yield response to levels and application methods of nitrogen. *Agrotechnology* **8**, 189.
- Arora VK, Singh CB, Sidhu AS, Thind SS (2011) Irrigation, tillage and mulching effects on soybean yield and water productivity in relation to soil texture. *Agricultural Water Management* **98**, 563-568.
- Beard T, Maaz T, Borrelli K, Harsh J, Pan W (2018) Nitrogen affects wheat and canola silica accumulation, soil silica forms, and crusting. *Journal of Environmental Quality* **47**, 1380-1388.
- Brandt S, Ulrich D, Lafond G, Malhi S, Johnston AM (2002) Management for optimum yield of open pollinated and hybrid canola. In Proc. Soils and Crops Workshop 2002 [CD-ROM]. pp. 21–22 (Feb. 2002. University of Saskatchewan, Saskatoon, SK, Canada)
- Budzynski W, Jankowski K, Szezebiot M (2000) Effects of simplifying soil tillage and weed control on yielding and production cost of winter oilseed rape. *Rośliny Oleiste* **21**, 487-502.
- Caia H, Mab W, Zhanga X, Pingc J, Yana X, Liua J, Yuana J, Wang L, Rena J (2014) Effect of subsoil tillage depth on nutrient accumulation, root distribution, and grain yield in spring maize. *The Crop Journal* **2**, 297-307.
- Chamoro AM, Tamagno LN, Bezus R, Sarandon J (2002) Nitrogen accumulation, partitioning and nitrogen use efficiency in canola under different nitrogen availabilities. *Communications in Soil Science and Plant Analysis* **33**, 493-504.
- Chandra K, Pandey A, Mishra SB, Kavita (2018) Genetic variability of physiological parameters among Indian mustard (*Brassica juncea* L. Czern and Coss) genotypes under non-irrigated and irrigated condition. *The Pharma Innovation Journal* **7**, 517-525.
- Cheema HS, Singh B (1991) Software statistical package CPCS-I. Department of Statistics, Punjab Agricultural University, Ludhiana, India.
- FAO (2019) Oilseeds and Oilseed products. OECD-FAO Agricultural Outlook 2019-2028/FAO 2019, pp. 47.
- Gajri PR, Arora VK, Kumar K (1994) A procedure for determining average root length density in row crops by single-site auguring. *Plant and Soil* **160**, 41-47.

- Hodge A (2006) Plastic plants and patchy soils. *J Exp Bot* 57:401-411.
- Kar G, Singh R, Verma HN (2005) Phenological based irrigation scheduling and determination of crop coefficient of winter maize in rice fallow of eastern India. *Agricultural Water Management* 75, 169-183.
- Kaur R, Arora VK (2019) Deep tillage and residue mulch effects on productivity and water and nitrogen economy of spring maize in north-west India. *Agricultural Water Management* 213, 724-731.
- Konwar K, Sarma A, Bharadwaj K, Boruah P (2019) Productivity and profitability of rapeseed (*Brassica rapa* var. *dichotoma*) under system of rapeseed intensification. *Journal of Oilseed Brassica* 10, 106-111.
- Kumar K, Kumar Y, Katiyar NK (2018) Effect of plant geometry, nitrogen level and antitranspirants on physiological growth, yield attributes, WUE and economics of mustard (*Brassica juncea*) under semi-arid conditions of western Uttar Pradesh. *Journal of Pharmacognosy and Phytochemistry* 7, 226-229.
- Lynch JP (2011) Root phenes for enhanced soil exploration and phosphorus acquisition: tools for future crops. *Plant Physiol* 156:1041-1049.
- Ma BL, Herath AW (2016) Timing and rates of nitrogen fertilizer application on seed yield, quality and nitrogen-use efficiency of canola. *Crop and Pasture Science* 67, 167-180.
- Mandal KG, Sinha AC (2004) Nutrient management effects on light interception, photosynthesis, growth, dry-matter production and yield of Indian mustard (*Brassica juncea*). *Journal of Agronomy and Crop Science* 190, 119-129.
- Mandal KG, Hati KM, Misra AK, Bandyopadhyay KK (2010) Root biomass, crop response and water-yield relationship of mustard (*Brassica juncea* L.) grown under combinations of irrigation and nutrient application. *Irrigation Science* 28, 271-280.
- Marcinkevicienė A, Velicka R, Raudonius S, Kosteckas R (2013) The relationship between root biomass and productivity of spring oilseed rape (*Brassica napus* L.) as influenced by crop density and fertilization. *Zemdirbyste Agriculture* 100, 39-44.
- Mishra J, Singh RK, Yadav D, Das A, Sahoo S (2019) Effect of tillage and irrigation frequency on growth of Indian mustard [*Brassica juncea* (L.) Czernj and Cosson]. *International Journal of Chemical Studies* 7, 2127-2130.
- Moroizumi T, Horino H (2002) The effects of tillage on soil temperature and soil water. *Soil*

Science **167**, 548-559.

Pal D, Phogat VK (2005) Effect of deep tillage and gypsum on yield and N, P, K and S uptake by mustard. *Journal of the Indian Society of Soil Science* **53**, 134-136.

Pal D, Phogat VK, Dahiya R (2009) Effect of deep tillage and gypsum on oil and protein content of mustard. *Research on Crops* **10**, 469-471.

Piper CS (1966) Soil and plant analysis. pp 368. Hans publishers, Bombay.

Prihar SS, Gajri PR, Benbi DK, Arora VK (2000) Intensive cropping: efficient use of water, nutrients and tillage. CRC Press.

Qamar R, Ehsanullah, Rehman A, Ali A, Ghaffar A, Mahmood A, Javeed HMR, Aziz M (2013) Growth and economic assessment of wheat under tillage and nitrogen levels in rice-wheat system. *American Journal of Plant Sciences* **4**, 2083-2091.

Rajput RK (2017) Effect of nitrogen and sulphur levels on nutrients uptake and yield of Indian mustard [*Brassica juncea* L. Czern and Coss.]. *International Archive of Applied Science and Technology* **8**, 29-31.

Rathke GW, Christen O, Diepenbrock W (2005) Effects of nitrogen source and rate on productivity and quality of winter oilseed rape (*Brassica napus* L.) grown in different crop rotations. *Field Crops Research* **94**, 103-113.

Rathore SS, Shekhawat, Kapila, Dass, Anchal, Premi OP, Rathore BS, Singh VK (2019) Deficit irrigation scheduling and superabsorbent polymerhydrogel enhance seed yield, water productivity, and economics of Indian mustard under semi-arid ecologies. *Irrigation and Drainage* **68**, 531-541.

Rathore SS, Shekhawat K, Singh RK (2017) Growth, productivity, input use and economics of Indian mustard as influenced by variable nitrogen doses under micro-irrigation. *Annals of Agricultural Research* **38**(3), 319-324.

Ray K, Sengupta K, Pal AK, Banerjee H (2015) Effects of sulphur fertilization on yield, S uptake and quality of Indian mustard under varied irrigation regimes. *Plant, Soil and Environment* **61**, 6-10.

Said-Al Ahl HAH, Mehanna HM, Ramadan MF (2016) Impact of water regime and phosphorus fertilization and their interaction on the characteristics of rapeseed (*Brassica napus*) and fatty acid profile of extracted oil. *Communications in Biometry and Crop Science* **11**, 64-76.

- Sarkar S, Kar S (2000) Root growth and water use pattern of groundnut at varying soil water regimes. In: Roy G B, Saha D and Datta D K (Eds.), Proceedings of the National Symposium on Principles of Soil Water Conservation, Mohanpur, Nadia, India, pp 1-5.
- Sharma A, Gupta R, Kumar A (2007) Production potential of mustard (*Brassica juncea* L.) under varying sowing method and at different sulphur and nitrogen levels. *Plant Archives* **7**, 893-896.
- Sharma PK, Kharwara PC, Tewatia RK (1990) Residual soil moisture and wheat yield in relation to mulching and tillage during preceding rainfed crop. *Soil and Tillage Research* **15**, 279-284.
- Shivran H, Kumar S, Tomar R, Chauhan GV (2018) Effect of irrigation schedules on productivity and water use efficiency in Indian mustard (*Brassica juncea* L.). *International Journal of Chemical Studies* **6**, 15-17.
- Sukirtee, Singh YV, Ranjan KR, Menka K, Bharteey PK (2018) Interaction effect of nitrogen and sulphur on yield, oil and nutrient content of mustard (*Brassica juncea* L.) in an inceptisol. *International Journal of Chemical Studies* **6**, 1234-1238.
- Sun X, Ding Z, Wang X, Hou H, Zhou B, Yue Y, Ma W, Ge J, Wang Z, Zhao M (2017) Subsoiling practices change root distribution and increase post-anthesis dry matter accumulation and yield in summer maize. *Plos one* **12**, 123-131.
- Tao Z, Li C, Li J, Ding Z, Xu J, Sun X, Zhao M (2015) Tillage and straw mulching impacts on grain yield and water use efficiency of spring maize in Northern Huang-Huai-Hai Valley. *The Crop Journal* **3**, 445-450.
- Tyagi PK, Upadhyay AK (2017) Growth, yield and water use efficiency of Indian mustard (*Brassica juncea* L.) as influenced by irrigation frequency and row spacing. *Journal of Oilseed Brassica* **8**, 27-36.
- White C, Roques S, Berry P (2015) Effects of foliar applied nitrogen fertilizer on oilseed rape (*Brassica napus*). *Journal of Agricultural Science* **153**, 42-55.

Comment [AC12]: Most of the bibliographic citations are very old. The citation pattern is not homogeneous.

Comment [AC13]: Improve table legends.

Table 1 Physico-chemical properties of experimental site

Depth (cm)	Soil separates (%)			Textural class	Bulk Density (Mg m ⁻³)	Water holding capacity (% v/v)		Available N (kg ha ⁻¹)
	Sand	Silt	Clay			FC	-15 bar	
0-15	79.3	11.7	7.6	Loamy Sand	1.38	14.5	4.0	80.3
15-30	80.6	12.3	6.3	Loamy Sand	1.42	15.0	4.5	77.1
30-60	81.9	9.5	8.5	Loamy Sand	1.47	17.0	6.0	60.3
60-90	83.5	7.7	8.8	Loamy Sand	1.53	17.0	6.8	52.8
90-120	84.3	8.4	7.3	Loamy Sand	1.55	18.0	6.5	55.5
120-150	82.8	8.7	8.5	Loamy Sand	1.57	18.2	6.3	42.1
150-180	81.5	9.8	8.7	Loamy Sand	1.59	18.3	6.4	40.1

Table 2 Monthly mean of daily maximum and minimum air temperature (°C) and monthly cumulative pan evaporation (E_p, mm) and rainfall (RF, mm) in different cropping seasons.

Month	2017-18				2018-19				Normal value			
	Temp		E _p	RF	Temp		E _p	RF	Temp		E _p	RF
	Max	Min			Max	Min			Max	Min		
October	33.0	18.5	88.7	0.0	31.3	17.1	96.6	0.0	31.8	16.4	123.4	10.9
November	24.7	11.4	47.4	7.0	27.0	11.8	64.0	2.6	26.6	10.6	81.3	6.7
December	20.8	7.5	50.6	24.0	20.7	5.5	40.0	0.0	20.4	6.5	53.4	17.6
January	18.7	6.2	46.0	18.4	18.5	6.2	43.1	66.0	18.2	5.6	49.5	28.7
February	22.8	9.1	64.4	27.0	20.1	9.2	46.2	95.6	21.0	7.6	48.4	33.3
March	29.3	13.9	125.3	0.0	25.3	11.8	84.8	7.4	26.6	11.7	118.5	21.0
April	35.8	19.9	218.3	10.0	35.1	19.5	170.3	41.6	34.4	17.1	211.4	27.8
Total	-	-	640.7	86.4	-	-	545	213.2	-	-	685.9	146

Table 3 Dry Matter Accumulation (DMA, t ha⁻¹) of canola crop at various stages as influenced by tillage, irrigation and nitrogen levels in different cropping seasons.

Treatment	60 DAS								110 DAS							
	2017-18				2018-19				2017-18				2018-19			
	N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹			
	0	50	75	100	0	50	75	100	0	50	75	100	0	50	75	100
I₀	0.13	0.16	0.26	0.34	0.20	0.29	0.35	0.45	0.30	0.57	0.84	1.24	0.43	0.64	0.92	1.26
CT I₁	0.24	0.69	0.76	0.84	0.25	0.74	0.82	0.87	0.89	1.50	1.76	2.51	0.98	1.61	1.88	2.59
I₂	0.25	0.58	0.75	0.90	0.26	0.65	0.80	0.93	1.36	1.69	2.17	2.98	1.44	1.73	2.25	3.16
DT I₀	0.13	0.18	0.26	0.39	0.23	0.27	0.35	0.39	0.37	0.79	0.94	1.36	0.47	0.90	1.02	1.51
I₁	0.28	0.65	0.79	0.93	0.35	0.69	0.80	0.95	1.02	1.61	1.95	2.85	1.08	1.67	1.98	3.01
I₂	0.30	0.59	0.72	0.86	0.32	0.67	0.82	0.92	1.55	2.08	2.35	2.97	1.63	2.12	2.55	3.21
Means	Year 17-18=0.50, 18-19=0.57								Year 17-18=1.57, 18-19=1.67							
	Tillage CT= 0.52; DT= 0.53								Tillage CT= 1.53; DT= 1.71							
	Irrigation I ₀ =0.27; I ₁ =0.67; I ₂ =0.64								Irrigation I ₀ =0.85; I ₁ =1.81; I ₂ =2.20							
	Nitrogen N ₀ =0.24; N ₅₀ =0.51; N ₇₅ =0.62; N ₁₀₀ =0.73								Nitrogen N ₀ =0.96; N ₅₀ =1.41; N ₇₅ =1.73; N ₁₀₀ =2.39							
LSD	Year=NS								Year=0.03							
(p=0.05)	Tillage= NS								Tillage= 0.09							
	Year ×Tillage=NS								Year ×Tillage=NS							
	Irrigation=0.05								Irrigation=0.10							
	Year × Irrigation= NS								Year × Irrigation= NS							
	Tillage × Irrigation=NS								Tillage × Irrigation=NS							
	Year ×Tillage × Irrigation= NS								Year ×Tillage × Irrigation= NS							
	Nitrogen=0.03								Nitrogen=0.06							
	Year × Nitrogen= NS								Year × Nitrogen= NS							
	Tillage × Nitrogen= NS								Tillage × Nitrogen= NS							
	Year ×Tillage × Nitrogen = NS								Year ×Tillage × Nitrogen = NS							
	Irrigation × Nitrogen= NS								Irrigation × Nitrogen= NS							
	Year × Irrigation × Nitrogen=NS								Year × Irrigation × Nitrogen=NS							
	Tillage × Irrigation × Nitrogen=NS								Tillage × Irrigation × Nitrogen=NS							
	Year × Tillage × Irrigation × Nitrogen=NS								Year × Tillage × Irrigation × Nitrogen=NS							

Comment [AC14]: Improve the legends of the table and its presentation.

Continued....

Treatment	145 DAS							
	2017-18				2018-19			
	N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹			
	0	50	75	100	0	50	75	100
I₀	0.72	1.32	2.14	2.38	0.82	1.43	2.20	2.43
CT I₁	1.92	2.72	4.26	5.34	1.38	2.92	4.58	5.50
I₂	3.10	4.24	6.02	7.09	3.40	4.76	6.28	7.47
I₀	1.65	2.70	2.95	3.78	1.85	2.86	3.18	4.07
DT I₁	2.08	3.81	5.33	6.13	2.23	4.14	5.88	6.77
I₂	4.46	5.69	8.23	8.55	4.60	5.91	8.47	8.74
Means	Year 17-18=4.03, 18-19=4.24							
	Tillage CT= 3.52; DT= 4.75							
	Irrigation I ₀ =2.28; I ₁ =4.06; I ₂ =6.06							
	Nitrogen N ₀ =2.35; N ₅₀ =3.54; N ₇₅ =4.96; N ₁₀₀ =5.69							
LSD	Year=0.04							
(p=0.05)	Tillage= 0.31							
	Year × Tillage=NS							
	Irrigation=0.38							
	Year × Irrigation= NS							
	Tillage × Irrigation=NS							
	Year × Tillage × Irrigation= NS							
	Nitrogen=0.21							
	Year × Nitrogen= NS							
	Tillage × Nitrogen= NS							
	Year × Tillage × Nitrogen = NS							
	Irrigation × Nitrogen= NS							
	Year × Irrigation × Nitrogen=NS							
	Tillage × Irrigation × Nitrogen=NS							
	Year × Tillage × Irrigation × Nitrogen=NS							

Table 4 Canola yield (t ha⁻¹) as influenced by tillage, irrigation and nitrogen levels in different cropping seasons.

Treatment	Seed Yield								Stover yield							
	2017-18				2018-19				2017-18				2018-19			
	N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹			
	0	50	75	100	0	50	75	100	0	50	75	100	0	50	75	100
CT I ₀	0.75	1.09	1.28	1.45	0.87	1.34	1.48	1.69	3.26	4.48	4.98	5.42	3.28	4.78	5.25	6.12
I ₁	0.97	1.21	1.42	1.58	1.03	1.50	1.77	1.88	3.79	4.71	5.71	6.10	3.96	5.48	6.35	7.03
I ₂	1.03	1.37	1.58	1.76	1.16	1.66	1.83	2.02	4.07	5.76	6.22	6.54	4.34	6.08	6.74	7.33
DT I ₀	0.86	1.19	1.36	1.53	0.96	1.43	1.59	1.95	3.66	4.69	5.26	5.51	3.69	5.21	5.89	7.20
I ₁	1.08	1.34	1.63	1.75	1.13	1.62	1.79	2.04	4.22	5.41	6.13	6.57	4.19	5.90	6.58	7.49
I ₂	1.24	1.55	1.74	1.89	1.35	1.82	1.97	2.24	4.44	6.06	6.50	6.73	4.85	6.59	7.24	8.23
Means	Year 17-18=1.36, 18-19=1.59								Year 17-18=5.26, 18-19=5.82							
	Tillage CT= 1.40; DT= 1.54								Tillage CT= 5.32; DT= 5.76							
	Irrigation I ₀ =1.30; I ₁ =1.48; I ₂ =1.64								Irrigation I ₀ =4.91; I ₁ =5.60; I ₂ =6.11							
	Nitrogen N ₀ =1.04; N ₅₀ =1.43; N ₇₅ =1.62; N ₁₀₀ =1.81								Nitrogen N ₀ =3.98; N ₅₀ =5.43; N ₇₅ =6.07; N ₁₀₀ =6.69							
LSD	Year=0.14								Year=NS							
(p=0.05)	Tillage= 0.06								Tillage= 0.19							
	Year × Tillage=NS								Year × Tillage=NS							
	Irrigation=0.07								Irrigation=0.23							
	Year × Irrigation= NS								Year × Irrigation= NS							
	Tillage × Irrigation=NS								Tillage × Irrigation=NS							
	Year × Tillage × Irrigation= NS								Year × Tillage × Irrigation= NS							
	Nitrogen=0.05								Nitrogen=0.19							
	Year × Nitrogen= 0.08								Year × Nitrogen= 0.27							
	Tillage × Nitrogen= NS								Tillage × Nitrogen= NS							
	Year × Tillage × Nitrogen = NS								Year × Tillage × Nitrogen = NS							
	Irrigation × Nitrogen= NS								Irrigation × Nitrogen= NS							
	Year × Irrigation × Nitrogen=NS								Year × Irrigation × Nitrogen=NS							
	Tillage × Irrigation × Nitrogen=NS								Tillage × Irrigation × Nitrogen=NS							
	Year × Tillage × Irrigation × Nitrogen=NS								Year × Tillage × Irrigation × Nitrogen=NS							

Table 5 Total Water Use (mm) and Water Productivity (kg ha⁻¹ mm⁻¹) as influenced by tillage, irrigation and nitrogen levels in different cropping seasons.

Treatment	TWU								Water productivity							
	2017-18				2018-19				2017-18				2018-19			
	N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹			
	0	50	75	100	0	50	75	100	0	50	75	100	0	50	75	100
I₀	250	253	256	257	298	301	305	308	3.02	4.30	4.98	5.65	2.91	4.46	4.85	5.48
CT I₁	319	323	327	329	357	359	363	365	3.04	3.74	4.35	4.79	2.89	4.19	4.88	5.16
I₂	360	367	374	376	401	404	414	417	2.85	3.74	4.24	4.68	2.91	4.12	4.41	4.83
DT I₀	249	254	258	259	308	316	325	333	3.43	4.69	5.28	5.92	3.11	4.53	4.89	5.84
I₁	314	317	321	322	368	371	373	376	3.44	4.24	5.09	5.42	3.07	4.38	4.79	5.41
I₂	363	371	382	385	420	426	434	440	3.41	4.17	4.55	4.92	3.23	4.27	4.55	5.09
Means	Year 17-18=316, 18-19=366								Year 17-18=4.33, 18-19=4.34							
	Tillage CT= 337; DT= 345								Tillage CT= 4.19; DT= 4.49							
	Irrigation I ₀ =283; I ₁ =344; I ₂ =396								Irrigation I ₀ =4.58; I ₁ =4.31; I ₂ =4.12							
	Nitrogen N ₀ =334; N ₅₀ =339; N ₇₅ =344; N ₁₀₀ =347								Nitrogen N ₀ =3.11; N ₅₀ =4.24; N ₇₅ =4.74; N ₁₀₀ =5.27							
LSD	Year=NS								Year=NS							
(p=0.05)	Tillage= 0.15								Tillage= 0.15							
	Year × Tillage=NS								Year × Tillage=NS							
	Irrigation=0.19								Irrigation=0.19							
	Year × Irrigation= NS								Year × Irrigation= NS							
	Tillage × Irrigation=NS								Tillage × Irrigation=NS							
	Year × Tillage × Irrigation= NS								Year × Tillage × Irrigation= NS							
	Nitrogen=0.15								Nitrogen=0.15							
	Year × Nitrogen= NS								Year × Nitrogen= NS							
	Tillage × Nitrogen= NS								Tillage × Nitrogen= NS							
	Year × Tillage × Nitrogen = NS								Year × Tillage × Nitrogen = NS							
	Irrigation × Nitrogen= NS								Irrigation × Nitrogen= NS							
	Year × Irrigation × Nitrogen=0.27								Year × Irrigation × Nitrogen=0.27							
	Tillage × Irrigation × Nitrogen=NS								Tillage × Irrigation × Nitrogen=NS							
	Year × Tillage × Irrigation × Nitrogen=NS								Year × Tillage × Irrigation × Nitrogen=NS							

Table 6 Effect of irrigation, tillage and nitrogen rates on oil yield (t ha⁻¹) and total N uptake (kg ha⁻¹) of canola

Treatment	Oil Yield (t ha ⁻¹)								Total N uptake (kg ha ⁻¹)							
	2017-18				2018-19				2017-18				2018-19			
	N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹				N- rate, kg ha ⁻¹			
	0	50	75	100	0	50	75	100	0	50	75	100	0	50	75	100
CT I ₀	0.32	0.46	0.54	0.61	0.35	0.53	0.59	0.66	41.66	62.86	74.58	90.52	43.88	67.64	76.77	91.88
I ₁	0.41	0.51	0.60	0.66	0.41	0.60	0.70	0.74	48.29	65.07	89.33	100.47	48.93	72.99	88.91	101.78
I ₂	0.44	0.58	0.67	0.75	0.47	0.66	0.73	0.80	47.93	77.37	90.57	100.55	51.33	78.91	91.20	104.36
DT I ₀	0.36	0.51	0.58	0.64	0.38	0.57	0.63	0.76	46.30	73.53	87.95	109.43	49.41	74.68	88.88	113.97
I ₁	0.46	0.57	0.69	0.74	0.45	0.65	0.71	0.80	52.83	81.06	91.57	110.97	55.55	81.82	94.84	112.33
I ₂	0.53	0.66	0.74	0.81	0.54	0.73	0.76	0.89	59.20	83.87	98.28	112.27	62.30	87.72	100.01	118.65
Means	Year 17-18=0.58, 18-19=0.63								Year 17-18=79.0, 18-19=81.6							
	Tillage CT= 0.57; DT= 0.63								Tillage CT= 75.3; DT= 85.3							
	Irrigation I ₀ =0.53; I ₁ =0.61; I ₂ =0.67								Irrigation I ₀ =74.6; I ₁ =81.0; I ₂ =85.3							
	Nitrogen N ₀ =0.43; N ₅₀ =0.59; N ₇₅ =0.66; N ₁₀₀ =0.74								Nitrogen N ₀ =50.6; N ₅₀ =75.6; N ₇₅ =89.4; N ₁₀₀ =105.6							
LSD	Year=NS								Year=2.2							
(p=0.05)	Tillage= 0.023								Tillage= 3.0							
	Year × Tillage=NS								Year × Tillage=NS							
	Irrigation=0.028								Irrigation=3.7							
	Year × Irrigation= NS								Year × Irrigation= NS							
	Tillage × Irrigation=NS								Tillage × Irrigation=NS							
	Year × Tillage × Irrigation= NS								Year × Tillage × Irrigation= NS							
	Nitrogen=0.023								Nitrogen=2.2							
	Year × Nitrogen= NS								Year × Nitrogen= NS							
	Tillage × Nitrogen= NS								Tillage × Nitrogen= NS							
	Year × Tillage × Nitrogen = NS								Year × Tillage × Nitrogen = NS							
	Irrigation × Nitrogen= NS								Irrigation × Nitrogen= NS							
	Year × Irrigation × Nitrogen=NS								Year × Irrigation × Nitrogen=NS							
	Tillage × Irrigation × Nitrogen=NS								Tillage × Irrigation × Nitrogen=NS							
	Year × Tillage × Irrigation × Nitrogen=NS								Year × Tillage × Irrigation × Nitrogen=NS							

Fig. 1: Effect of tillage on soil penetration resistance (MPa) distribution in 0-60 cm soil layer

Fig. 2: Effect of tillage and nitrogen rates on depth-wise root mass density (RMD, μgcm^{-3}) of canola under different irrigation regimes during cropping season 2017-18.

Fig. 3: Effect of tillage and nitrogen rates on depth-wise root mass density (RMD, μgcm^{-3}) of canola under different irrigation regimes during cropping season 2018-19.





Fig. 1: Effect of tillage on soil penetration resistance (MPa) distribution in 0-60 cm soil layer

Fig. 2: Effect of tillage and nitrogen rates on depth-wise root mass density (RMD, μgcm^{-3}) of canola under different irrigation regimes during cropping season 2017-18.











Fig. 3: Effect of tillage and nitrogen rates on depth-wise root mass density (RMD, μgcm^{-3}) of canola under different irrigation regimes during cropping season 2018-19.

Comment [AC15]: Take into account the recommendations made previously for tables and figures.

UNDER PEER REVIEW