

Estimation of Actual Evapotranspiration (ET_a) of major crops of adistributary of Mahanadi canal command using CROPWAT 8.0 Model.

ABSTRACT:

This paper aims to estimate Actual Evapotranspiration of major crops by using cropwat 8.0 software in 2A canal command of Mahanadi of Dhamtari district. Actual evapotranspiration is a key process of hydrological cycle and a sole term that links land surface water balance and land surface energy balance. Irrigation is an essential part of different crops because rainfall is not enough for irrigated farmland. Long term daily meteorological data including rainfall, maximum temperature, minimum temperature, relative humidity, wind speed and sunshine hours of IMD station data was used as input data in modified Penman method and CROPWAT 8.0 model. The crop stage data, including the value of K_c in, K_c mid, K_c end for the selected crop was obtained from department of irrigation the rooting depth, critical depletion and crop height of different crops are taken from FAO Irrigation and Drainage Paper 56. The Penman Monteith method was used to estimate ET_o and ET_c respectively. The major crop in 2A canal command area are Paddy, Wheat, Chickpea, Summer Paddy. Based on the intensive study of this paper, daily basis meteorological weather data recorded from 2007 to 2021 were used to obtain the result. The study detects that Penman–Montieth method is the best method to estimate ET_a because of its inclusion of parameters in calculation.

Key words: Reference Evapotranspiration, Crop coefficient, Modified Penman method, CROPWAT 8.0, Crop water requirement, Actual Evapotranspiration.

1. INTRODUCTION

Evapotranspiration is a process by which water is lost by evaporation and transpiration. In real world conditions, the evaporation and transpiration can occur at the same time and it is very difficult to differentiate between the two. When the crop is small, soil evaporation is the primary source of evapotranspiration however, once the crop has matured and fully covers the soil, transpiration becomes the primary source of water loss. Temperature, Relative humidity, movement of wind and movement of air, availability of Soilmoisture, different type of crops available are parameters affecting evapotranspiration. Evapotranspiration calculation is critical not only for the analysis of climate emergency and the assessment of water supplies, and include for crop water demand, drought forecast and tracking effective water resource output and use. ET , from an agricultural perspective, decides the volume of water to be supplied artificially. ET estimation is essential since it defines the size of irrigation channels and other irrigation. Evapotranspiration can be measured directly or indirectly and depend on weather data and soil

Formatted: Font: (Default) Times New Roman, 12 pt, Bold

water balance. These methods are broadly known as empirical (e.g., Thornthwaite method, Blaney and Criddle) or physical methods (e.g., Penman method ; Montheith, and FAO Penman Montheith method (Allen et al., 1998). To better estimate crop water requirements, the Food and Agriculture Organization recommends the use of CROPWAT software, CROPWAT is frequently used for crop evapotranspiration, reference crop evapotranspiration, irrigation scheduling, and cropping patterns prediction. The study emphasis on using CROPWAT model for estimating actual evapotranspiration.

2. STUDY AREA.

The present study will be carried out in Mahanadi Command area Dhamtari and Kurud Block of Northern part of Dhamtari district of Chhattisgarh State. The area lies between 20.7404 and 21.0829 N latitudes and 81.4750 and 81.8530 E longitudes (Fig. 1). The geographical extension of the study area is 975 sq km representing around 29% of the district's geographical area. One of the major rivers of North India is Mahanadi, which originates from the Sihawa hill, located on the east of Nagri tehsil. Mahanadi is the principal river of the Dhamtari district along with its tributaries and Kharun on western boundary of block respectively. Dhamtari district is agriculturally intensive due to the establishment of good network of irrigation canals from New Rudri Barrage, which was constructed on the ~~Mahanadi river~~ Mahanadi River. The major crop of this district and kurud block is paddy in kharif season and in Rabi season mostly summer paddy whereas paddy is grown in command areas of tank, and wheat, chickpea, pulses (mainly gram and millets) and oil seeds are taken. 2A Canal whose length is 22200 M and Cultural command area is 12461.58 and Discharge (Q) = 11.40 CUMEC.

Comment [Pa1]: It cannot be that an introduction contains only a bibliographical reference

Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Formatted: List Paragraph, Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"




Fig.1. Location map of the study area

3. MATERIAL AND METHODS

Following are data used in CROPWAT8.0 software to estimate Actual Evapotranspiration for each crop (Table 1):

Table 1.List 1 Model Input data for CROPWAT8.0

Climate Data	Minimum Temperature
	Maximum Temperature
	Relative Humidity
	Wind Speed
	Sunshine Hours
Rainfall Data in mm	Daily
Crop Data	K _c value for paddy ,wheat, chickpea, summer paddy

All the input data are required for model to calculate actual evapotranspiration (ET_a). All the daily Climate data are taken from IMD (India Meteorological Department) of 15 years and rainfall data is also taken daily basis of 15 years as well as crop data is of 15 years daily basis.

3.1.METHODOLOGY:

3.1.1. DATA:

The input for the computation of ET_a requires meteorological variables such a maximum temperature, minimum temperature, relative humidity, wind speed,sunshine hours and rainfall. The meteorological database has been generated for 15 years from (2007- 2021) on daily basis. Using the CROPWAT software reference evapotranspiration was calculated for the entire study area which uses FAO56 Penman Monteith method. Soil data have been collected from Department of Soil Science and Agricultural Chemistry, IGKV, Raipur. Crop data have been collected from Agriculture Department of dhamtariblock.

3.1.1. CROPWAT:

CROPWAT 8.0 developed by FAO, based on FAO Irrigation and Drainage Paper 56 named FAO56. FAO56 adopted the P - M (Penman - Monteith) method as a global standard to estimate ETo from meteorological data. The Penman–Monteith equation is used for computation of daily

Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: List Paragraph, Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"

Formatted Table

Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Formatted: List Paragraph, Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"

Formatted: Font: Bold

Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Formatted: List Paragraph, Outline numbered + Level: 3 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.75"

reference evapotranspiration. Penman-Monteith equation is mathematically expressed as shown in equation (1)

$$ET_0 = 0.408\Delta (R_n - G) + \gamma \frac{900}{T+273} U(e_s - e_a)$$

$$\Delta + \gamma(1+0.34U)$$

Where,

ET_0 : reference crop evapotranspiration

[mm/day]

R_n : net radiation at the crop surface

[MJ/ m² /day]

G: soil heat flux [MJ/(m² .d)]

T : average air temperature [° C]

U : wind speed measured at 2 m height[m/s]

e_s :saturation vapour pressure [kPa]

e_a : actual vapour pressure [kPa]

($e_s - e_a$) : saturation vapour pressure deficit [kPa]

Δ : slope of the vapor pressure curve,

[kPa/ ° C]

γ : psychrometric constant, [kPa/ °C]

900: conversion factor.


The FAO CROPWAT program incorporates procedures for reference crop evapotranspiration and crop water requirements and allow the simulation of crop water use under various climates, crop and soil conditions.

3.1.2. CROPDATA:

The major cultivated crops in the study area are paddy, wheat, chickpea, summer paddy. Crop coefficient values (K_c) are taken from available published data. K_c values for initial, development, mid-season and late-season growth stages of different crops are used.

Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Formatted: List Paragraph, Outline numbered + Level: 3 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.75"



[Fig.2.Methodology for Estimation of Actual Evapotranspiration](#)

3.1.4. Actual Evapotranspiration:

Actual evapotranspiration is mathematically a product of reference evapotranspiration and crop coefficient, which is given in the equation below:-

$$ET_a = ET_0 \times K_c$$

where,

$ET_a = \text{Actualevapotranspiration}$

$ET_o = \text{Reference evapotranspiration}$

$K_c = \text{Crop Coefficient}$

The crop coefficient changes with the growing stages of the crop. The value of K_c for any crop is most likely to be less in planting stage and reaches a maximum at mid season. In the present study wheat, paddy, chickpea, summer paddy crop is chosen for determination of evapotranspiration. The K_c values for different growth stages as per FAO 56 is shown in Table 1.

TABLE Table+2. Crop coefficient for various growing stages of different crop:-

Sl. No.	Crops	Growing stages of crops			
		Initial	Development	Mid season	Late season
1.	Paddy	1.10	1.20	1.20	1.05
2.	Wheat	0.30	1.15	1.15	0.30
3.	Chickpea	0.40	1.0	1.0	0.35
4.	Summer paddy	1.05	1.20	1.20	0.70

4. RESULTS AND DISCUSSION:-

The reference evapotranspiration was calculated using the CROPWAT 8.0 Model and actual evapotranspiration was calculated by multiplying reference evapotranspiration with the crop coefficient. The input data provided for CROPWAT model includes minimum temperature, maximum temperature, latitude, longitude, altitude, sunshine hours and wind velocity. The input data was collected and analysed for a decade starting from 2007 to 2021 in the Dhamtari region. The daily potential evapotranspiration was obtained and tabulated using the CROPWAT 8.0 model as shown in Tables.

Comment [Pa2]: The results and discussion part is very poorly presented, only tables have been shown and there is no discussion part

Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Formatted: List Paragraph, Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"

Table 23: Reference Evapotranspiration of the study area by CROPWAT model for Rice Crop

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2007	2.18	2.88	3.46	3.89	4.11	3.77	2.92	2.98	3.06	1.84	2.55	2.18
2008	2.12	2.55	3.26	4.07	3.95	3.04	3	2.92	3.47	3.23	2.4	2.04
2009	2.23	2.76	3.06	3.8	4.37	4.18	2.91	3.26	3.54	3.02	2.33	2.08
2010	2.07	2.81	3.47	4.26	4.55	4.04	3.22	3.38	3.38	3.24	2.58	2.03
2011	2.29	2.79	3.64	4.33	4.78	3.8	3.44	2.99	2.97	3.38	2.6	2.08

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

2012	1.85	2.78	3.15	4.2	4.29	3.8	2.81	2.83	3.18	3.24	2.41	2.15	Formatted: Font: (Default) Times New Roman
2013	2.12	2.81	3.34	4.17	4.54	3.67	2.97	2.8	3.25	2.72	2.57	2.12	Formatted: Font: (Default) Times New Roman
2014	2.12	2.81	3.34	4.13	4.41	3.84	3.04	3.18	2.93	3.07	2.5	1.96	Formatted: Font: (Default) Times New Roman
2015	2.56	2.94	3.27	3.69	3.64	3.21	3.09	3.24	3.61	3.5	2.82	1.75	Formatted: Font: (Default) Times New Roman
2016	1.94	2.19	3.15	4.22	4.59	4.28	3.07	3.01	2.89	3.05	2.52	2.06	Formatted: Font: (Default) Times New Roman
2017	2.29	2.86	3.34	3.85	4.4	4.07	3.04	3.37	3.58	3.32	2.75	2.4	Formatted: Font: (Default) Times New Roman
2018	2.21	2.66	3.2	4	4.5	4.13	2.66	2.61	3.33	3.39	2.65	1.72	Formatted: Font: (Default) Times New Roman
2019	1.69	2.57	2.95	3.35	3.62	3.43	3.18	2.96	3.05	3.04	2.39	1.65	Formatted: Font: (Default) Times New Roman
2020	2.04	2.54	3.02	3.49	3.93	3.38	3.18	2.78	2.55	2.66	2.5	1.6	Formatted: Font: (Default) Times New Roman
2021	1.68	2.01	2.72	2.69	3.47	3.37	3.05	2.85	2.57	3.23	2.38	2.17	Formatted: Font: (Default) Times New Roman

Table 34: Reference Evapotranspiration of the study area by CROPWAT model for Chickpea Crop

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2007	2.18	2.88	3.46	3.89	4.11	3.77	2.92	2.98	3.06	1.84	2.55	2.18
2008	2.12	2.55	3.26	4.07	3.95	3.04	3	2.92	3.47	3.23	2.4	2.04
2009	2.23	2.76	3.06	3.8	4.37	4.18	2.91	3.26	3.54	3.02	2.33	2.08
2010	2.07	2.81	3.47	4.26	4.55	4.04	3.22	3.38	3.38	3.24	2.58	2.03
2011	2.29	2.79	3.64	4.33	4.78	3.8	3.44	2.99	2.97	3.38	2.6	2.08
2012	1.85	2.78	3.15	4.2	4.29	3.8	2.81	2.83	3.18	3.24	2.41	2.15
2013	2.12	2.81	3.34	4.17	4.54	3.67	2.97	2.8	3.25	2.72	2.57	2.12
2014	2.12	2.81	3.34	4.13	4.41	3.84	3.04	3.18	2.93	3.07	2.5	1.96
2015	2.56	2.94	3.27	3.69	3.64	3.21	3.09	3.24	3.61	3.5	2.82	1.75
2016	1.94	2.19	3.15	4.22	4.59	4.28	3.07	3.01	2.89	3.05	2.52	2.06
2017	2.29	2.86	3.34	3.85	4.4	4.07	3.04	3.37	3.58	3.32	2.75	2.4
2018	2.21	2.66	3.2	4	4.5	4.13	2.66	2.61	3.33	3.39	2.65	1.72
2019	1.69	2.57	2.95	3.35	3.62	3.43	3.18	2.96	3.05	3.04	2.39	1.65
2020	2.04	2.54	3.02	3.49	3.93	3.38	3.18	2.78	2.55	2.66	2.5	1.6
2021	1.68	2.01	2.72	2.69	3.47	3.37	3.05	2.85	2.57	3.23	2.38	2.17

Table 45: Reference Evapotranspiration of the study area by CROPWAT model for Wheat Crop

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2007	2.18	2.88	3.46	3.89	4.11	3.77	2.92	2.98	3.06	18.4	2.55	2.18
2008	2.12	2.55	3.26	4.07	3.95	3.04	3	2.92	3.47	3.23	2.4	2.04
2009	2.23	2.76	3.06	3.8	4.37	4.18	2.91	3.26	3.54	3.02	2.33	2.08

2010	2.07	2.81	3.47	4.26	4.55	4.04	3.22	3.38	3.38	3.24	2.58	2.03
2011	2.29	2.79	3.64	4.33	4.78	3.8	3.44	2.99	2.97	3.38	2.6	2.08
2012	1.85	2.78	3.15	4.2	4.29	3.8	2.81	2.83	3.18	3.24	2.41	2.15
2013	2.12	2.81	3.34	4.17	4.54	3.67	2.97	2.8	3.25	2.72	2.57	2.12
2014	2.12	2.81	3.34	4.13	4.41	3.84	3.04	3.18	2.93	3.07	2.5	1.96
2015	2.56	2.94	3.27	3.69	3.64	3.21	3.09	3.24	3.61	3.5	2.82	1.75
2016	1.94	2.19	3.15	4.22	4.59	4.28	3.07	3.01	2.89	3.05	2.52	2.06
2017	2.29	2.86	3.34	3.85	4.4	4.07	3.04	3.37	3.58	3.32	2.75	2.4
2018	2.21	2.66	3.2	4	4.5	4.13	2.66	2.61	3.33	3.39	2.65	1.72
2019	1.69	2.57	2.95	3.35	3.62	3.43	3.18	2.96	3.05	3.04	2.39	1.65
2020	2.04	2.54	3.02	3.49	3.93	3.38	3.18	2.78	2.55	2.66	2.5	1.6
2021	1.68	2.01	2.72	2.69	3.47	3.37	3.05	2.85	2.57	3.23	2.38	2.17

Table 56: Reference Evapotranspiration of the study area by CROPWAT model for Summer Rice Crop

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2007	2.18	2.88	3.46	3.89	4.11	3.77	2.92	2.98	3.06	1.84	2.55	2.18
2008	2.12	2.55	3.26	4.07	3.95	3.04	3	2.92	3.47	3.23	2.4	2.04
2009	2.23	2.76	3.06	3.8	4.37	4.18	2.91	3.26	3.54	3.02	2.33	2.08
2010	2.07	2.81	3.47	4.26	4.55	4.04	3.22	3.38	3.38	3.24	2.58	2.03
2011	2.29	2.79	3.64	4.33	4.78	3.8	3.44	2.99	2.97	3.38	2.6	2.08
2012	1.85	2.78	3.15	4.2	4.29	3.8	2.81	2.83	3.18	3.24	2.41	2.15
2013	2.12	2.81	3.34	4.17	4.54	3.67	2.97	2.8	3.25	2.72	2.57	2.12
2014	2.12	2.81	3.34	4.13	4.41	3.84	3.04	3.18	2.93	3.07	2.5	1.96
2015	2.56	2.94	3.27	3.69	3.64	3.21	3.09	3.24	3.61	3.5	2.82	1.75
2016	1.94	2.19	3.15	4.22	4.59	4.28	3.07	3.01	2.89	3.05	2.52	2.06
2017	2.29	2.86	3.34	3.85	4.4	4.07	3.04	3.37	3.58	3.32	2.75	2.4
2018	2.21	2.66	3.2	4	4.5	4.13	2.66	2.61	3.33	3.39	2.65	1.72
2019	1.69	2.57	2.95	3.35	3.62	3.43	3.18	2.96	3.05	3.04	2.39	1.65
2020	2.04	2.54	3.02	3.49	3.93	3.38	3.18	2.78	2.55	2.66	2.5	1.6
2021	1.68	2.01	2.72	2.69	3.47	3.37	3.05	2.85	2.57	3.23	2.38	2.17

Table 67: Weightage Crop coefficient for major crops of the study area.

Sl. NO.	Crops	Weightage K _c
1	Rice	1.030
2	Chickpea	0.782
3	Wheat	0.831
4	Summer Rice	1.030

Table 78: Actual Evapotranspiration (ET_a mm/day) values obtained from CROPWAT model of all major Crops

Year	RICE	WHEAT	CHICKPEA	SUMMER
2007	3.4834	2.4453	2.0691	3.4031
2008	3.2259	2.3572	1.9815	3.2857
2009	3.4834	2.3090	1.9549	3.3413
2010	3.5555	2.4420	1.9971	3.5349
2011	3.4154	2.5068	2.0864	3.6729
2012	3.2609	2.3373	1.9502	3.3516
2013	3.1744	2.4254	2.0081	3.4978
2014	3.3083	2.4138	2.0284	3.4628
2015	3.4299	2.3971	2.0707	3.3166
2016	3.3578	2.2026	1.8547	3.3145
2017	3.5802	2.3938	2.0457	3.4484
2018	3.3207	2.4054	2.0691	3.4134
2019	3.2259	2.0414	1.8125	2.9210
2020	2.9973	2.1178	1.8218	3.0941
2021	3.1044	1.7787	1.6450	2.5441

5. CONCLUSIONS :

Water requirements of crops varies with climate, soil type and crop variety. CROPWAT 8.0 model is the tool for irrigation planning and management because of considering many input parameters. Evapotranspiration provides the necessary information regarding water requirements for growing different crops in different seasons. The results show that the crop water requirement of different crops will consequently help improving the management of water resources and productivity. The CROPWAT 8.0 model gives sufficiently accurate result and reduce the calculation and also consumes less time. The major crop in 2A canal command area are Paddy,

Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Formatted: List Paragraph, Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"

Wheat, Chickpea, Summer Paddy. Based on the intensive study of this paper, daily basis meteorological weather data recorded from 2007 to 2021 were used to obtain the result. The study detects that Penman–Montieth method is the best method to estimate ETa because of its inclusion of parameters in calculation.

REFERENCES

A. Hashem, “Performance Evaluation and Development of Daily Reference Evapotranspiration Model”, Irrigation and Drainage Sys Eng 2016.

C. Y. Xu and V. P. Singh, “Evaluation and generalization of radiationbased methods for calculating evaporation, Hydrological processes”, Vol 14, Issue 2, pp 339–349, 2000.

Comment [Pa3]: They are not mentioned in text

G. Allen, L. S. Pereira, D. Raes and M. Smith, “Crop evapotranspiration guidelines for computing crop water requirements” FAO Irrigation and drainage paper 56. Food and Agriculture Organization, Rome 1998.

G. Lindstrom, B. Johansson, M. Persson, M. Gardelin and S. Bergström, “Development and test of the distributed HBV-96 hydrological model”, Journal of Hydrology, Volume 201, Issues 1–4, pp 272-288, 20 December 1997.

G. Stancalie, A. Marica and L. Toullos, “Using earth observation data and CROPWAT model to estimate the actual crop evapotranspiration”, Physics and Chemistry of the Earth, Parts A/B/C, vol 35(1-2), pp 25–30, 2013

K. T. Zeleke, L. J. Wade, “Evapotranspiration Estimation using Soil Water Balance, Weather and Crop Data”, Evapotranspiration: Remote Sensing and Modeling, pp 41-58, 2012.

L. Zhao, J. Xia, C. Xu, Z. Wang, L. Sobkowiak and C. Long , “Evapotranspiration estimation methods in hydrological models”, journal of Geographical Sciences, vol 23,pp 359-369,2013.

T. S. Bajirao and H. W. Awari, “Estimation of reference evapotranspiration for Parbhani district”, International Journal of Agricultural Engineering, vol 10, pp 51-54, 2017.

V. Singh, Vijay Kumar and Avinash Agarwal, “Reference Evapotranspiration by Various Methods for Kashmir Valley”, Journal of Indian Water Resources Society, Vol. 26, pp 3- 4,2006.

Comment [Pa4]: They are not mentioned in text