

Review Article

Cabbage (*Brassica oleracea*) Production in Kenya: A review of its economic importance, ecological requirement and production constraints

Abstract

Cabbage is an important vegetable containing a considerable amounts of water, protein, lipids, carbohydrates, fibre, vitamins, assorted minerals, phenols and glucosinolates. In Kenya, cabbage production is done mainly by smallholder farmers for food and sale. The crop is mainly grown within 800m and 2900m above sea level altitude. It plays an important role towards achieving improved human nutrition and health conditions, reducing poverty through food security and enhancing ecosystem resilience as a source of income, human food and livestock feed. Statistics show that though average area under production has been growing since 2017, average production has been fluctuating and consumption increasing steadily. Profitable production of the crop within the country is mainly constrained by myriad of factors such as climate change effects, pest and disease prevalence, poor agronomic practices, land degradation, soil health deterioration, lack of structured marketing systems and poor research- extension- farmer linkages. Possession of limiting climate smart agriculture knowledge and skills, challenging access to credit facilities and agro processing technologies as well as narrow post-harvest loss management knowledge and skills was identified as other key cabbage production constraints within the country. It was however noted that the country has potential to achieve profitable cabbage farming if adoption of climate smart technologies, farmer- extension linkage, government policies, credit facilities as well as structured market system were streamlined. The current work reviews cabbage crop, with emphasis on its biology, economic importance, ecological requirements, current production status in Kenya and production constraints.

Keywords: Cabbage, production constraints, climate change.

1.0 Introduction

Cabbage is an important vegetable widely consumed in the world. Its production varies around the world with China being the leading producer (32,800,000 tonnes) followed by India (8,500,000 tonnes), and Russia (3,309,315 tonnes) [1]. The origin of cabbage is not clear, however, it is suggested to have originated from the South and Western coast of Europe or the Eastern Mediterranean, Asia minor region, or ancient Greek [2,3]. According to Dixon [4] and Leike [5], the crop was first cultivated in the Netherlands in the 14th century, and later spread to other European and Mediterranean countries. It was introduced in Africa in the 16th century by European colonialists [4,6].

In Kenya, cabbage is cultivated by both small-scale and medium-scale farmers. It is grown in all parts that fall between 800m and 2900m above sea level [7,8]. The major producing areas include; Kiambu, Nakuru, Nyandarua, Nyeri, Muranga, Narok, Kerinyaga, Laikipia, Kisii, Nyamira, Meru, Bomet, Elgeyo Marakwet, Muranga, Trans-Nzoia, Makeni, Laikipia, Bungoma, and Kericho [7, 9,10,11]. In the year 2020, the area under production was 30,258Ha amounting to 977,054 tonnes, while in the year 2019 it was 33, 473Ha amounting to 991,549 tonnes. In the years 2018, 2017, and 2016 the production area was 22,892Ha, 26,807Ha, and 25,926Ha respectively, amounting to 620, 523 tones, 713 360 tones, and 733,817 tones in that order [11,12,13].

Numerous varieties are grown in Kenya, however, the most popular include; Gloria F1, Copenhagen, Sunny F1, Karen F1, Riana F1, Queen F1, Polo F1, Pruktor F1, Serena F1, Sugar Loaf, Green Challenger F1, Golden Acre, and Rosy F1. Other varieties available in the market include; Prize Drumhead, Fortuna, K-Y Cross, Early Market F1, Super Master F1, Oxylus, Star 3308 F1, Santa F1,

Rotan F1, Field Winner F1, Globe Master Hybrid F1, Fanaka F1, Amigo, Amuko Si F1, Barak A F1, Blue Dynasty F1, Early Drum Head, Green Coronet F1, Fiona F1, Zawadi F1 and Tristar among others [14, 15]. Most of these cabbage varieties are either consumed locally or transported to major towns within the country. The left over and over supply are utilized as dairy cattle and heifers feed.

2.0 Botany of Cabbages

Taxonomically, cabbage is classified under kingdom Plantae, division Spermatophyta, class Dicotyledonae, family Cruciferae/Brassicaceae or (mustard), and genus Brassica [3,16]. Within this genus, there are many species, which include; *Brassica oleracea* var. *capitata* (white cabbage) and *Brassica rapa* subsp. *Pekinensis* (Napa cabbage) [3, 16,17, 18]. Within a species, there are different categories of cabbage depending on their colour. These categories are green, while others are red/purple. Green cabbages range from very dark to very light and these light green-coloured cabbages are referred to as "white" cabbages because the inner leaves are white. Red/purple cabbages range widely in colour, with some appearing deep purple [5]. Within each cabbage category, various varieties have been bred to suit different ecological zones. Phenotypically they are leafy biennial crops, but they are grown as annual vegetables that are 40 – 60cm tall Brassica [3,16]. Some of these varieties form an oval shape when mature, while others are round shaped [19].

When cabbage matures, the basal leaves form a rosette of 7 to 15 sessile outer leaves each 250 to 350 × 200 to 300mm in size. The upper leaves form a compact flattened globosely ellipsoidal head that is 100 to 300mm in diameter that is formed as a result of fleshy leaves overlapping on one another around a single growing point. The stems are unbranched, 200 to 300mm long and are gradually thickened upwards. The single growing point has a ramified system of thin roots, where several fibrous roots are emerging from the taproot. About 90% of these roots are found in the upper 200 to 300 mm of the soil to the ground surface and some laterals penetrate down to 1.5 to 2m deep [16]. Closely related members of cabbages (*Brassica oleracea* Capitata and *Brassica rapa* subsp. *Pekinensis*) include; kale and collards (*Brassica oleracea* Acephala), Chinese broccoli (*Brassica oleracea* Alboglabra), cauliflower (*Brassica oleracea* Botrytis), Brussels sprouts (*Brassica oleracea* Gemmifera), kohlrabi (*Brassica oleracea* Gongylodes), broccoli (*Brassica oleracea* Italica), Field mustard (*Brassica rapa* subsp. Oleifera), Choy sum (*Brassica rapa* subsp. parachinensis) Bomdong (*Brassica rapa* var. glabra), Bok choy (*Brassica rapa* subsp. Chinensis) [16,17,18].

3.0 Cabbage Ecological Requirements

Cabbage require relatively cool and humid climate [20]. The optimum temperatures for growth and development range from 15°C to 25°C. It is fairly resistant to frost and can survive temperatures as low as -3°C without damage [14,16, 21]. Water requirement is variety depended but vary from 380mm to 500mm per crop throughout the growing period. Crop water use, however, increases with growth period with a peak towards the end of the season during head maturity. Soil moisture less than 380mm causes limping and wilting while moisture higher than 500mm causes root rot leading to crop death [22]. Irrigation and water management or conservation practices are therefore important to ensure adequate soil moisture in areas prone to high evaporation [23]. This crop is equally sensitive to flooding, therefore, during the rainy season, appropriate drainage canals must be in place to drain off the excess water after heavy rain [14].

Light plays a key role in cabbage growth. It is crucial in determining photo-morphogenesis and photosynthetic biosynthesis [24]. Adequate light intensity is therefore required for maximum photosynthesis. Where light intensity is insufficient, photosynthetically active radiation (PAR), leaf chlorophyll content, and photosynthetic rate will decrease resulting in reduced formation of carbohydrates, leading to stem elongation and etiolation [25]. Sufficient light at photosynthetic photon flux density (PPFD) of 10 $\mu\text{mol m}^{-2} \text{s}^{-1}$ is therefore needed to suppress the succulent elongation to increase dry matter deposition [26].

According to Adeniji [20], cabbage crop is termed to be a heavy feeder that needs plenty of nutrients. It therefore, thrives well under deep, well-drained fertile loamy soils with approximately 600mm rooting depth and adequate amounts of nutrients such as nitrogen, phosphorus and potassium. Although Nitrogen plays critical role in vigorous growth and all metabolic activities, excess levels causes loose head formation and internal decay. Phosphorus on the other hand plays a critical role in metabolism,

germination and root development [27, 28]. Low Potassium (K) on the other hand result in marginal necrosis and lower head quality, but when it is in excess it can cause the heads to open. The plant has high Sulfur requirement and it is sensitive to Calcium, Magnesium, and Boron deficiency. The general fertilizer recommendation is therefore 100 - 150kg/ha N: 50 – 65kg/ha P: 100 - 130kg/ha K. The fertilizer is applied in splits that are site specific. For example, three splits can be applied as follows: a basal, 3 weeks after transplanting, and during heading. In a situation where soils are sandy addition of manure and a mulching layer of 15 to 20cm deep is recommended [14,16].

Ideal soil pH for the crop ranges from 5.5 to 7.5 [14, 15,16]. Where the soil pH is above 6.5, the leaves become dark, and the leaf margins die back while pH below 5.5, causes chlorosis and retarded growth. In saline soils, the crop is highly susceptible to blackleg disease [15,16]. Drainage is recommended in waterlogged soils and leaching of salts in saline conditions [23].

4.0 Economic Importance of Cabbages

According to Wambani *et al.* [29], and Kibata [30], cabbage is among the popular leafy vegetable that is widely consumed by both rural and urban dwellers in Kenya. It is highly nutritious comprising of 92% water, 1% proteins, 0.1% lipids, and 5% carbohydrates [31]. It also contains 2% fiber, Vitamin A, C, and K, and mineral ions, like calcium, iron, sodium, zinc, magnesium, phosphorus, and potassium, which are crucial in human nutrition and health [32,33]. Cabbage also contains phenolics and glucosinolates [methylsufinylbutyl isothiocyanate (sulforaphane, SFN) and phenethyl isothiocyanate (PEITC); and indole-3-carbinol (I3C) and 3,3'-diindolylmethane (DIM)] that are anti-carcinogenic for effective reduction of cancer-related diseases and prevent type 2 diabetes, cardiovascular diseases, inflammation of digestive systems, cataracts, and Alzheimer's disease [34,35]. The main cancer types cabbages is named to be active against include: lung cancer, gastric cancer, breast cancer, prostate cancer, colon cancer, rectum cancer, liver cancer, and pancreatic cancer [1]. A high intake of cabbages also ensures sufficient levels of iodine in the body of humans which can aid the proper functioning of the brain, thyroid gland, and nervous system [36].

The cabbage industry serves as a source of employment for many Kenyans ranging from small scale farmers to middle men and agribusiness players [8]. This empowers rural population especially women and youths who are largely unemployed, have little capital and limited access to land, working under labour constraints. In return money obtained from this enterprise contributes to food and nutritional security at the household level as well as enabling women and the youths to attain some degree of financial independence within the family budget [37,38,39].

5.0 Production Constraints of Cabbage Farming in Kenya

Despite the growing population and increased demand for food and nutrition security within the country, cabbage industry growth has been constrained over the years. Cabbage farming has been challenged by myriad of factors such as: Pest and disease prevalence, impact of climate change, land degradation, soil health decline, erratic rains leading to water scarcity, weed infestation, land tenure systems and land fragmentation, unstructured marketing systems, post-harvest losses, limited primary processing technologies and services, lack of harvesting technologies as well as poor agronomic technical knowledge [14, 15,40, 41].

5.1 Pest and Disease Prevalence

Cabbage is very susceptible to pest infestations in the field, which causes huge losses [40,42]. The major insect pests infesting cabbages in Kenya and other African states include: the diamondback moth (*Plutella xylostella*), the cabbage webworm (*Helula undalis*), white butterfly (*Pieris brassicae*), cabbage aphid (*Brevycoryne brassicae*), cabbage looper (*Trichoplusia ni*) and green peach aphids (*Myzus persicae*) [15; 40,42]. However, the diamondback moth, (*Plutella xylostella*), and cabbage webworm (*Helula undalis*) are the most destructive insect pests of cabbage crops [15,43,44]. These insect pests infest the *Brassica oleracea* crop at different stages of growth, causing huge destruction during the growth stages and eventually resulting in huge yield losses [42,44]. Other cabbage pests that attack at seedling stage include; cutworm (*Agrotis spp*) flea beetles (*Phyllotreta striolata*), cabbage root maggot (*Delia radicum*), ants (*Formicidaees ssp.*), cabbage head caterpillar (*Crocidolomia pavonana*), and spider mites (*Tetranychus spp*).

Diseases negatively affect cabbage yields if not properly managed. The most disastrous diseases are classified as fungal, and bacterial. Fungal diseases include; Leaf Spot (*Alternaria brassicae*), Blight (*Alternaria brassiciola*), Downy Mildew (*Peronospora parasitica*), Damping off (*Pythium debaryanum*), Sclerotinia rot/ White Mould (*Sclerotinia sclerotiorum*) Yellows or Fusarium Wilt (*Fusarium oxysporum* f. sp *conglutinans*), and Black Leg (*Phoma lingam*). Conversely, common bacterial diseases include; Bacterial Soft Rot (*Pectobacterium carotovorum*), Black Rot (*Xanthomonas campestris* pv. *campestris*), Wire Stem (*Rhizoctonia solani*), and Clubroot of Cabbage (*Plasmodiophora brassicae*), [14,15,45]. Most fungal diseases are commonly field based while bacterial diseases are both field and post-harvest.

5.2 Climate Variability and Limited Climate Smart Agriculture Knowledge and Skills

Considering that optimal crop growth and yield necessitate provision of all nutrients in correct forms and amounts, water plays a major role as a solvent for all soil chemical reactions that eventually determine germination, nutrient uptake, metabolisms, assimilation, photosynthesis and all other plant biochemical processes [46,47]. Most of the cabbage production regions within the country have been experiencing either unpredictable, increased or reduced rainfall over the past years. The constant off season increased rainfall or on season erratic coupled with limited climate smart agricultural practices knowledge and adoption among other factors constraint cabbage production within production areas leading to increased wastage due to high levels of moisture during maturity, or increased disease infections or reduced yields. [14,41]. Cabbage being a cool and humid climate crop, the optimum temperatures for growth and development range from 15°C to 20°C, therefore temperatures above 24°C delay maturity and vegetative growth and number of leaves increase leading to the formation of loose heads or failure of head formation [19,48,49]. This makes Kenyan farmers in cabbage producing areas to depend heavily on cooler seasons for cabbage production.

5.3 Competing farm enterprises

The demand for highly valued food has become the driving factor for resource allocation which consequently influences resources allocated to cabbage production [50]. Competing farm enterprises as a result of pressure from food demand by the growing human population has created challenges in cabbage production with farmers, shifting to production of vegetables with faster maturity rate and higher revenue [51,52].

5.4 Land degradation and declining soil health

Land degradation interacting with climate change represents one of the biggest and most urgent challenges for humanity today. According to FAO reports, land degradation within Kenya which is commonly in form of poor soil health is attributed to poor agricultural activities. The activities include: i) Poor fertility management and nutrient cycling (ii) intensification (iii) land subdivision leading to limited fallow period (iv) leaching of soil nutrients into deeper soil horizons in areas that receive relatively high amounts of (v) soil erosion (vii) nutrient mining; (viii) physical, chemical and biological degradation of the soil through acidification and soil pollution [53,54,55,56,57,58]. According to Weil and Brady [46], adequate availability of nutrients in rightful proportions is paramount in crop growth and productivity. Availability of these nutrients for crop uptake is, however, determined by prevailing soil physical, chemical and biological properties which are often compromised in un healthy soils. Owing to the fact that the mineralogy of most cabbage growing regions is acidic in nature, poor soil health management predispose the soils to acidity problem leading to poor availability of certain essential macro nutrients and biota as well as toxicity of some micronutrients [46,57,59]. These situation calls for continuous soil health management through application of amendments such as organic fertilizers, inorganic fertilizers and lime as well as adoption of integrated soil fertility management climate smart technologies. This is challenging in majority of cabbage growing areas because farmers grow cabbage as mono-crop, season after season and the inputs are also shared with other high value crops. This leads to nutrient mining leading to poor yields and return to capital. Farmers who use commercial fertilizers also often use straight fertilizers that don't provide all limiting nutrients within their soils.

5.5 Poor Structured supply chain and market Systems

According to Dyeyi and Zenda [60], access to profitable marketing of agricultural produce rely on infrastructure, supply chain and institutions. Availability of better transport network enables access to

markets hence minimizing information dissymmetry about input and output quality and prices. There, however, exists poor coordination between supply chain actors of most perishable farm products, along with inefficient retail cold chain management practices within the country; identified as the vital causes of cabbage and other perishable crop produce wastage. Both commercial and small-scale farmers utilize considerable time in coordinating the supply chain of their cabbages to reach markets but large percentage of their produce is often spoilt before it reaches target market. This cabbage spoilage, deterioration and wastage, results in revenue losses for players within the cabbage value chain whose burden is often carried by producers. According to Chamberlin and Jayne [61], market access within most agricultural production areas have improved slightly over the years with large numbers of independent traders buying directly from rural farms. This mechanism has however been a drawback to farmers with some traders either under weighing commodities or offering low prices. At times farmers incur losses due to impassable roads during rain seasons leading to destruction of cabbages and other fresh produce before reaching the market.

5.6 Access to Agro Processing Technologies

According to Kirsten et al. [62] and KALRO [14], farmers access to agro processing technologies is crucial to guarantee profitability of a product after harvesting; something that limits crop production in most African countries. Most cabbage farmers within the country either don't have access to agro processing facilities or have access to old and outdated processing methods. This expose them to high post-harvest losses leading to low enterprise profitability.

5.6 In Adequate Access to Agricultural Information and Extension Services

Access to agricultural information is critical and very crucial to increased agricultural productivity [63]. According to Yaseen [64], limited information sources in rural areas constrains farmer's agricultural production and growth. Public agricultural extension officers offer significant role of linking farmers with current technologies and market information. Currently agricultural extension services are dimly offered to farmers, yet the government spends on agricultural research as a proportion of GDP. By 2016 it was 0.48%, approximately one-third of its value in 2006 [65]. The research- extension – farmer link in Kenya has however been deteriorating over the years with the current extension: farmer ratio standing at 1800:1 compared to the FAO recommendation of 400:1 [66,67,68,69]. As a result, many farmers have limited sources of useful and reliable, soil and water management, agronomic and marketing information leading to low uptake of technologies, crop production and yields in the region.

According to Ndambiri *et al.* [70] and Dhakshana and Rajandran [71], there exists a positive relationship between new technologies adoption level and the availability of credit. Availability of credit eases the cash constraints and enabling farmers to purchase inputs such as improved cultivars, fertilizer, pesticides with ease. Lack of security among horticultural farmers in Kenya, however limit their access to credit facilities [72]. Additionally, though government policies play a major role in cushioning farmers, unlike other crops, there exists limited specific legal or regulatory framework for most horticultural crops production [73].

5.7 Land Fragmentation and land tenure challenges

High population densities within Kenya highlands imply that farm sizes per farmer have continuously declined small with average total farm size near the slopes of Mt. Kenya being between 1.0 and 2.0 hectares. Several enterprises are allocated within this small sized pieces limiting mechanization and economic viability. This also encourages intensification while limiting integrated soil fertility and integrated pest and disease management.

6.0 Conclusion and Recommendations

The review indicates that cabbage plays a significant role achieving improved human nutrition and health conditions, lowering poverty through food security and income generation. They act as source of human food and nutrition by playing a role in the management of diseases such as cancer-related diseases, type 2 diabetes, cardiovascular diseases, and inflammation of the digestive systems. Statistics show that although area under production has been increasing over the years. Production on the other hand has not increased significantly and it is constrained by myriad of factors. Some of the factors that constrain cabbage production within Kenya include: Pests and disease prevalence, climate change effects, soil health decline due to nutrient mining without adequate replenishment. Other factors such as poor access

to agricultural information, credit facilities, poor structured marketing system as well as access to agro processing technologies play a key role in constraining cabbage production within the country. Land fragmentation, poor post-harvest losses management and delivery systems were also identified to be key limiting factors. The review revealed that the country has a potential to support maximum cabbage production. This however requires adoption of climate smart technologies, improved post-harvest handling technologies, farmer- extension linkage, government policies, credit facilities as well as structured market system.

8.0 References

- 1) Moreb N, Murphy A, Jaiswal S, & Jaiswal AK. Cabbage. *Nutritional Composition and Antioxidant Properties of Fruits and Vegetables*, 2020; 33-54.
- 2) Maggioni L, Von Bothmer R, Poulsen G, Branca F. Origin and domestication of cole crops (*Brassica oleracea*): linguistic and literary considerations. *Economic botany*, 2010; 64(2):109-123.
- 3) Maggioni L. Domestication of *Brassica oleracea* L. Faculty of Landscape Architecture, Horticulture and Crop Production Science, 2015; 74Pgs.
- 4) Dixon G. *Vegetable brassicas and related crucifers: Crop Production Science in Horticulture Series*, 14. Willingford: UK. 2007.
- 5) Leike H. Cabbage (*Brassica oleracea* var. *capitata* L.). In *Crops II* 1988; pp. 226-251.
- 6) Maggioni L, Von Bothmer R, Poulsen G, Lipman E. Domestication, diversity and use of *Brassica oleracea*., based on ancient Greek and Latin texts. *Genetic resources and crop evolution*, 2018; 65(1): 137-159.
- 7) Macharia I, Löhr B, De Groote H. Assessing the potential impact of biological control of *Plutella xylostella* (diamondback moth) in cabbage production in Kenya. *Crop Protection*, 2005; 24(11): 981-989.
- 8) Gitange J O, Nyangeri J, Maobe S. An economic analysis on the use of hot water seed treatment, mulching and pruning in the control of black rot disease in cabbage with in Kisii County. *Int. J. Curr. Microbiol. App. Sci*, 2018; 7(12): 413-426.
- 9) Muriuki AW, Schnier HF, Recke H, Kanyanjua SM. The effect of farmyard manure and fertilisers on cabbage yields in Kenya. *East African Agricultural and Forestry Journal*, 2002; 68(1): 41-49.
- 10) Mwatabu E, Were JO, Anjichi VE. Response of Selected Cabbage Varieties to *Alternaria brassicicola* under Greenhouse Conditions. Faostat 2018
- 11) Horticultural Crops Directorate. Manual. 2019- 2020 Validated Horticulture Report
- 12) Horticultural Crops Directorate. Manual. 2018- 2019 Validated Horticulture Report
- 13) Horticultural Crops Directorate. Manual. VALIDATED REPORT 2017-2018
- 14) KALRO. Inventory of climate smart agriculture technologies, innovations and management practices for cabbage value chain. KALRO Headquarters, Nairobi kenya. 2022; 386 pgs.
- 15) Japan International Cooperation Agency. Cabbage production. Smallholder Horticulture Empowerment & Promotion Project for Local and Up-Scaling (SHEP PLUS). 2019; https://www.jica.go.jp/project/english/kenya/015/materials/c8h0vm0000f7o8cj-att/materials_21.pdf
- 16) Bewick TA. Cabbage: uses and production. *Florida Cooperative Extension Service Fact Sheet*, 1994; 712.
- 17) Burt J, Phillips D, Gatter D. *Growing chinese cabbage in Western Australia*. Department of Agriculture and Food, Western Australia, Perth. Bulletin, 2006; 4673, 1-23.
- 18) Daly P, Tomkins B. Production and postharvest handling of Chinese cabbage. *Rural Industries Research and Development Corporation (RIRDC), Barton ACT, Australia, RIRDC Research Paper*, 1997; 97(1); 1-40.
- 19) Mudzingwa S, Muzemu S, Chitamba J. Pesticidal efficacy of crude aqueous extracts of *Tephrosia vogelii* L., *Allium sativum* L. and *Solanum incanum* L. in controlling aphids (*Brevicoryne brassicae* L.) in rape (*Brassica napus* L.) n controlling aphids (*Brevicoryne brassicae* L.) in rape (*Brassica napus* L.). *Journal of Research in Agriculture*, 2013; 2(1); 157-163.
- 20) Adeniji OT, Swai I, Oluoch MO, Tanyongana R, Aloyce A. Evaluation of head yield and participatory selection of horticultural characters in cabbage (*Brassica oleracea* var. *capitata*). *Journal of Plant Breeding and Crop Science*, 2010; 2(8):243-250.
- 21) Daugovish O, Cahn M, Koike S, Natwick E, Cantwell M, Takele E. Cabbage Production in California. *California Agriculture and Natural Resources*, 2008; 1-5.

- 22) Food and Agriculture Organization of the United Nations (FAO). Retrieved from <https://www.fao.org/faostat/en/#compare>. Accessed 31th May, 2022.
- 23) Kelley WT, Granberry DM, Boyhan GE, Langston DB, Adams DB, MacDonald G, Westberry GO (Eds). Commercial production and management of cabbage and leafy greens. *University of Georgia College of Agricultural and Environmental Sciences*, 2009; 1-45.
- 24) Avercheva OV, Berkovich YA, Erokhin AN, Zhigalova TV, Pogosyan SI, Smolyanina SO. Growth and photosynthesis of Chinese cabbage plants grown under light-emitting diode-based light source. *Russian Journal of Plant Physiology*, 2009; 56(1):14-21.
- 25) Fukuoka N, Yoshioka H, Shimizu E, Fujiwara T. Effect of shading cabbage (*Brassica oleracea* var. capitata) seedlings on their physiological processes and rooting ability after transplanting to the nursery. *Journal of the Japanese Society for Horticultural Science*, 1996; 65(3):545-551.
- 26) Sato F, Yoshioka H, Fujiwara T. Effect of dim light on growth and photosynthetic ability of cabbage plug-transplants during low-temperature storage. *Environment Control in Biology*, 2002; 40(3):317-320.
- 27) Hawkesford M, Horst W, Kichey T, Lambers H, Schjoerring J, Skrumsager IM, White P. Function of macronutrients. p. 135-178. In Marschner, P. (ed.) *Marschner's mineral nutrition*. 3rd ed. Elsevier Ltd., Oxford, UK. 2012.
- 28) Muindi EM. Understanding soil phosphorus: A review. *International Journal of Soil Science*. 2019;31(2):1-18.
- 29) Wambani H, Nyambati EM, Kamidi M, Mulati J. Participatory evaluation of cabbage varieties as a source of food and income for smallholder farmers in north western Kenya. In *Proceedings of the 8th African Crop Science Society Conference, El-Minia, Egypt*, 27-31. 2007.
- 30) Kibata GN. The diamondback moth: a problem pest of brassica crops in Kenya. *The Management of Diamondback Moth and Other Crucifer Pests. Proceedings of the 3rd International Workshop, Kuala Lumpur*, 1997; 29: 47-53.
- 31) Butnariu M, Butu A. Chemical composition of Vegetables and their products. *Handbook of Food Chemistry*, 2015;1-49.
- 32) Singh B K, Sharma SR, Singh B. Variation in mineral concentrations among cultivars and germplasms of cabbage. *Journal of plant nutrition*, 2009; 33(1):95-104.
- 33) Cervantes-Godoy D, Dewbre J. Economic importance of agriculture for poverty reduction. 2010.
- 34) Sarikamış G. Glucosinolates in crucifers and their potential effects against cancer. *Canadian journal of plant science*, 2009; 89(5): 953-959.
- 35) Abdull Razis AF, Noor N M. Cruciferous vegetables: dietary phytochemicals for cancer prevention. *Asian Pacific Journal of cancer prevention*, 2013; 14(3):1565-1570.
- 36) Pandey V, Chura A, Pandey HK. Cabbage: A Storehouse of Nutraceuticals. In *Medicinal Plants*, 2018;1-16. CRC Press.
- 37) Lenné J. Promotional Opportunities for Outputs from the Peri-urban African Vegetable Cluster Projects. *A report to the CPP*. 2002.
- 38) Ota M, Lenné J. Effectiveness of commercial horticulture networks as pathways for poverty alleviation, exploitation and promotion of pest management technologies in Kenya. *Natural Resources International, Aylesford, Kent, UK*.2003.
- 39) Oruko L, Ndun'gu B. Final Socio-Economic Report for Peri-urban Vegetable IPM Thematic Cluster. A report to the DFID Crop Protection Programme. *Natural Resources International, Aylesford, Kent, UK*.2001.
- 40) Baidoo PK, Adam J I. The effects of extracts of *Lantana camara* (L.) and *Azadirachta indica* (A. Juss) on the population dynamics of *Plutella xylostella*, *Brevicoryne brassicae* and *Hellula undalis* on cabbage. *Sustainable Agriculture Research*, 2012; 01(2): 229-234.
- 41) Kenya Meteorological Department (KMD). Extreme weather events in Kenya between 2011& 2020. https://meteo.go.ke/sites/default/files/downloads/Extreme%20Climate%20Events_Kenya%202011_to_2020.pdf
- 42) Baidoo PK, Mochiah MB. Comparing the effectiveness of garlic (*Allium sativum* L.) and hot pepper (*Capsicum frutescens* L.) in the management of the major pests of cabbage *Brassica oleracea* (L.). *Sustainable Agriculture Research*, 2016; 5(2):83- 90.
- 43) Furlong MJ, Wright DJ, Dossdall LM. Diamondback moth ecology and management: problems, progress, and prospects. *Annual review of entomology*, 2013; 58: 517-541.

- 44) Labou B, Bordat D, Brevault T, Diarra K. Spatiotemporal distribution and impact of diamondback moth parasitoids in the Dakar Niayes in Senegal. *International Journal of Biological and Chemical Sciences*, 2017; 11(3): 1288-1298.
- 45) Sharma A, Rathore JP, Ali A, Qadri I, Hussain SM, Angmo T. Major diseases and pathogen ecology of cabbage. *The Pharma Innovation Journal*, 2018; 7(7): 574-578.
- 46) Weil R, Brady NC. The nature and properties of soil. 15th edition. Pearson. 2017: 1104pp
- 47) Muindi EM, Wamukota AW, Mulinge JM, Okello N, Wekesa G, Ahmed H. Onion (*Allium cepa* L.) Performance as influenced by manure and fertilizer in titanium mined soils. *Sustainable Agricultural Research*, 2022; 11(3):27-37.
- 48) Ngoye BO, Saado H, Gachari CW. Who could have seen this coming? The Kenya Red Cross society and the drought appeal of 2019. *Emerald Emerging Markets Case Studies*. 2021.
- 49) Daniel KA, Muindi EM, Gogo EO, Muti S. Performance of Brassica rapa and Brassica oleracea under Shade net within Coastal Environment. *Journal of Agriculture and Ecology Research International*, 2022; 23(3): 45-59.
- 50) Hasanov S, Nomman A. Agricultural efficiency under resources scarcity in Uzbekistan: A Data Envelopment Analysis. *Business and Economic Horizons (BEH)*, 2011; 4(1232-2016): 81-87.
- 51) Shiundu KM, Oniang'o RK. Marketing African leafy vegetables: Challenges and opportunities in the Kenyan context. *African Journal of food, agriculture, nutrition and development*, 2007; 7(4): 1-17.
- 52) Masayi N, Netondo GW. Effects of sugarcane farming on diversity of vegetable crops in Mumias Division, Western Kenya. 2012.
- 53) Lal R. Restoring soil quality to mitigate soil degradation. *Sustainability*, 2015;7(5): 5875-5895.
- 54) Tamene L, Amede T, Kihara J, Tibebe D, Schulz S. A review of soil fertility management and crop response to fertilizer application in Ethiopia: towards development of site-and context-specific fertilizer recommendation. *CIAT Publication*. 2017.
- 55) Muindi EM, Mrema JP, Semu E, Mtakwa PW, Gachene CK. Phosphate sorption characteristics and external phosphorus requirements of Nitisols in Central Kenya Highlands. *International Journal of Soil Science*, 2017; 12(3):113-119.
- 56) Esilaba AO, Miriti J, Gitari J, Mangale N, Kathuku-Gitonga AN, Njiru E, Muindi EM, Nassiuma ...et al., KCEP-CRAL Integrated soil fertility and Water Management Extension Manual. Kenya Agricultural and Livestock Research Organization, Nairobi, Kenya. 2021; 61Pgs.
- 57) Muindi EM, Mrema J, Semu, E, Gachene, C. Soil Chemical Properties as Affected by Tillage Method, Phosphorus and Lime Application. *International Journal of plant and soil sciences*, 2023; 35(16): 213-221.
- 58) Muindi EM, Semu E, Mrema JP, Mtakwa PW, Gachene CK, Njogu MK. Soil Acidity Management by Farmers in the Kenya Highlands. *Journal of Agriculture and Ecology Research International*, 2016; 5(3): 1-11.
- 59) Muindi EM, Mrema JP, Semu E, Mtakwa PW, Gachene CK, Njogu MK. Phosphorus adsorption and its relation with soil properties in acid soils of Western Kenya. *International Journal of Plant Science*, 2015;4(3):203-211.
- 60) Chamberlin J, Jayne TS. Unpacking the meaning of 'market access': evidence from rural Kenya. *World development*, 2013; 41:245-264.
- 61) Dyeyi MI, Zenda M. Challenges in the supply chain of cabbage in Nelson Mandela Bay Metropolitan municipality, Eastern Cape, South Africa. *South African journal of agricultural extension*,2022; 50(2): 2022. <http://dx.doi.org/10.17159/2413-3221/2022/v50n2a11282>
- 62) Kirsten J, Mapila M, Okello J. De S. Managing agricultural commercialization for inclusive growth in Sub-Saharan Africa. *Global Development Network Working Paper No. 60*. 2013.
- 63) Kelil A, Mamo D, Seyoum D. *Factors influencing smallholder dairy farmers'access to agricultural information in ada'a woreda, east shoa zone, central ethiopia* (Doctoral dissertation, Haramaya university).2017.
- 64) Yaseen M, Xu S, Yu W, Hassan S. Farmers' access to agricultural information sources: Evidences from rural Pakistan. *Journal of Agricultural Chemistry and Environment*, 2016; 5(1): 12-19.
- 65) Birch I. Agricultural productivity in Kenya: barriers and opportunities.2018.
- 66) Kondylis F, Mueller V, Zhu J. Seeing is believing? Evidence from an extension network experiment. *Journal of Development Economics*, 2017; 125: 1-20.

- 67) MOALF. Situational analysis of the agriculture sector in Kenya. Final Report. Nairobi, Kenya. 2020: 154Pgs
- 68) Kazungu FK, Muindi EM, Mulinge JM. Overview of sorghum (*sorghum bicolor.L.*), its economic importance, ecological requirements, and production constraints in Kenya. *Journal of Plant and Soil Sciences*,2023; 35(1):62-71
- 69) Muchomba MK, Muindi EM, Mulinge JM. Overview of greengram (*Vigna radiata L.*) crop, its economic importance, ecological requirements and production constraints in Kenya. *Journal of Agriculture and Ecology Research International*, 2023; 24(2):1-11.
- 70) Ndambiri HK, Ritho C, Ng'ang'a, SI, Kubowon PC, Mairura FC, Nyangweso PM, Cherotwo FH. Determinants of economic growth in Sub-Saharan Africa: A panel data approach. *International Journal of Economics and Management Sciences*, 2012; 2(2): 18-24.
- 71) Dhakshana A, Rajandran KVR. Challenges and problems on farmers' access to agricultural credit facilities in Cauvery Delta, Thanjavur District. *St. Theresa Journal of Humanities and Social Sciences*, 2018; 4(1): 50-62.
- 72) Nyachae SM, Warue B. Horticultural Financing Lending Policies on Performance of Financing Institutions in Kenya. *The University Journal*, 2020: 2(1): 57-70.
- 73) Government of Kenya. National Horticultural Policy. Agricultural Sector Coordination Unit, Nairobi, Kenya. 2012; 76 Pg.