

Response of NPK and FYM levels on Physicochemical Properties of Soil under carrot crop (*Daucus carota*) in an Inceptisol of Prayagraj district, Uttar Pradesh, India

ABSTRACT

An experiment was conducted to investigate the “Response of NPK and FYM levels on the Physicochemical Properties of Soil under carrot crop (*Daucus carota*) in an Inceptisol of Prayagraj district, Uttar Pradesh, India” during the Rabi season of 2022 at the Research Farm of the Department of Soil Science and Agricultural Chemistry, NAI, SHUATS, Prayagraj. The experimental having two factors with three levels of N P K @ 0%, 50% and 100% ha⁻¹, and three levels of F Y M @ 0%, 50% and 100% ha⁻¹ which was layout in randomized block design (RBD). The results showed that the physicochemical properties of soil were significantly affected with the application of FYM in conjunction with NPK fertilizers. This was evident through a significant increase in organic carbon percentage, per cent pore space, and higher levels of available nitrogen, phosphorus, and potassium in the soil. Additionally, the combination of NPK and FYM led to a decrease in bulk density, particle density, and pH of soil. In conclusion, the experiment demonstrated that the combination of NPK and FYM positively influenced the soil health under carrot cultivation.

Key words: Nantes, FYM, NPK, Carrot, Soil Properties, Yield

1. INTRODUCTION

Soil is a dynamic non-renewable natural resource that is vital to life. There are connections between soil and land use, water flow, water quality, and vegetation productivity have an important relation with soil. The ability of the soil to supply all of the necessary plant nutrients in readily available forms and in an appropriate balance is referred to as soil fertility, whereas soil productivity is the outcome of a number of factors. The nutrients and physical characteristics can have a direct impact on yields in cultivated fields. **Ahmad et. al (2014)**

Soil plays an important role in carrot production by providing nutrients, water and physical support for growth. To grow healthy and straight carrots, it is important to provide the right soil conditions. Loose soil helps the roots to grow straight down without obstruction, while clumps or hard clayey chunks can cause deformities. Carrots require soil that is rich in Nitrogen potassium and phosphorus to encourage healthy root growth. Soil fertility management *via.* organic and inorganic fertilizers can improve fertility and increase Carrot

yield. The ideal pH level for carrots is between 6 to 6.8. However, carrots can grow well in pH level anywhere between 5.5 to 7.5. **Yadav et. al (2010)**

Carrot (*Daucus carota*), is a member of the family Apiaceae and important root vegetable crops of the world. It is grown in spring, summer and autumn in temperate countries and during winter in tropical and subtropical countries. It is mainly a cold requiring crop but it can also be grown at a relatively higher temperature without much difficulty. In India it is grown during winter when the temperature ranges from 11.17 to 28.9°C. Risk of growing carrot is also much less due to fewer problems of diseases and insects' pests. **Yadav et. al (2010)**

Fertilizer plays a crucial role in carrot cultivation as it affects growth and yield attributes. Nitrogen increases vegetative growth and promotes carbohydrate synthesis, while phosphorus stimulates root diameter and increases growth rate. Potassium is important for root development and the manufacture of sugar and starch. Improper doses of N, P, and K can negatively impact growth and yield, and excessive nitrogen can cause root splitting and reduce marketable yield. Proper application of manure and fertilizers, especially NPK, can increase crop yield. **Singh et. al (2007)**

The use of organic manures such as FYM, vermicompost, and poultry manure has gained prominence in recent years as they can improve crop productivity and maintain soil fertility. FYM is a valuable organic amendment for soil improvement in agriculture. When applied to soil, FYM enriches it with organic matter, enhancing soil structure, water holding capacity, and nutrient retention. The slow release of nutrients from FYM supplies plants with essential elements over time, while also improving nutrient availability in the soil. FYM acts as a pH buffer, regulating soil pH and creating optimal conditions for plant growth. Additionally, it enhances water retention, drainage, and infiltration, making it beneficial in water-scarce and waterlogged conditions. FYM promotes the growth of beneficial microorganisms, contributing to nutrient cycling, organic matter decomposition, maintaining the ideal C:N ratio and disease suppression. Its positive impacts on soil quality and plant productivity have been widely recognized, leading to increased soil fertility, higher crop yields, and reduced reliance on synthetic inputs. **Verma et. al (2018)**

Considering the importance of soil fertility and the role of NPK and FYM in carrot cultivation, this experiment was conducted to investigate the effects of different levels of NPK and FYM on the physicochemical properties of soil under carrot crop cultivation. The study aimed to evaluate the changes in organic carbon percentage, pore space, pH, bulk density, particle density, and the availability of nitrogen, phosphorus, and potassium in the

soil. The results would provide valuable insights into the combined influence of NPK and FYM on soil health and optimize the management practices for carrot cultivation.

2. MATERIALS AND METHODS

The field experiment was conducted at Research Farm of Soil Science and Agricultural Chemistry at Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj which is situated at 25°24'23" N latitude, 81°50'38" Longitude and at the altitude of 98 meter above the sea level.

The fieldwork was done in the Prayagraj district, which is part of the subtropical belt and has semi-arid climatic conditions with both winter and summer temperature extremes. "The maximum temperature of the location reaches up to 46°- 48°C and seldom falls as 4°-5°C. The relative humidity ranges between 20% to 94%. The average annual rainfall in Prayagraj is around 900-1100 mm annually. The minimum temperature during the crop season was to be 6.9°C and the maximum is to be 28.57°C. The maximum humidity was to be 89% and minimum was to be 93.28%". The present research investigation was setup in randomized block design (RBD) with nine treatment combinations, which are replicated three times and randomly allocated in each replication, dividing the research site into twenty- seven plots. In this study, NPK and FYM was applied in three different doses. Sowing of the Carrot crop was carried out on the 19th Nov, 2022, by hand. The seed variety Nantes was sown at a rate of 4 kg ha⁻¹ and at a row-to-row spacing of 25 cm and plant-to-plant spacing of 5 cm. The recommended doses of NPK were applied @60:80:70 kg ha⁻¹. The graded level of NPK were applied through Urea, Single super phosphate and Murate of potash. Half dose of nitrogen and full dose of phosphorus and potassium were applied basally at the time of sowing. In addition to these applications, FYM was used as a basal dose at 0, 10, and 20 t ha⁻¹ for the treatment.

The soils from each plot were separately collected, air-dried, ground, and passed through a 2-mm-size sieve for laboratory analysis. Soil samples were analyzed for bulk density, particle density, Percentage pore space, and waterholding capacity (**Muthuvel et al., 1992**), pH (**Jackson, 1958**), EC (**Wilcox, 1950**), Percentage Organic Carbon (**Walkley and Black, 1947**), Available Nitrogen (**Subbiah and Asijja, 1956**), Available Phosphorus (**Olsen et al., 1954**) and Available Potassium (**Toth and Prince, 1949**) before sowing and after harvest of the crop. Findings of the Soil analysis before sowing of the crops at depth

of 0-15 and 15-30 cm are bulk density (1.32 and 1.35 Mg m⁻³), particle density (2.52 and 2.53 Mg m⁻³), pore space (43.79 and 41.54%), water holding capacity (40.19 and 39.27 %), pH (7.24 and 7.26) EC (0.29 and 0.27 dS m⁻¹), organic carbon (0.35 and 0.34 %), available nitrogen (224.84 and 221.25 kg ha⁻¹), available Phosphorus (21.51 and 18.41kg ha⁻¹) and available potassium (188.39 and 185.82kg ha⁻¹).

Table: 1. Treatment Combinations

| TREATMENT | TREATMENT COMBINATION | Symbol |
|----------------|---------------------------|---------------------------------------|
| T ₁ | (Absolute control) | NPK ₀ FYM ₀ |
| T ₂ | NPK @ 0 % + FYM @ 50 % | NPK ₀ FYM ₅₀ |
| T ₃ | NPK @ 0 % + FYM @ 100 % | NPK ₀ FYM ₁₀₀ |
| T ₄ | NPK @ 50 % + FYM @ 0 % | NPK ₅₀ FYM ₀ |
| T ₅ | NPK @ 50 % + FYM @ 50 % | NPK ₅₀ FYM ₅₀ |
| T ₆ | NPK @ 50 % + FYM @ 100 % | NPK ₅₀ FYM ₁₀₀ |
| T ₇ | NPK @ 100 % + FYM @ 0 % | NPK ₁₀₀ FYM ₀ |
| T ₈ | NPK @ 100 % + FYM @ 50 % | NPK ₁₀₀ FYM ₅₀ |
| T ₉ | NPK @ 100 % + FYM @ 100 % | NPK ₁₀₀ FYM ₁₀₀ |

3. Result and Discussion

3.1. Physical and Chemical analysis of soil

The results showed that the treatment T₉ (NPK₁₀₀FYM₁₀₀) was recorded maximum physio chemical characters such as pore space (%), water holding capacity (%), EC (dS m⁻¹), organic carbon (%), available nitrogen (kg ha⁻¹), available Phosphorus (kg ha⁻¹) and available potassium (kg ha⁻¹). While bulk density (Mg m⁻³), particle density (Mg m⁻³) and pH were recorded maximum in T₁ (NPK₀ FYM₀).

Table 2: Response of NPK and FYM levels on bulk density, particle density, percent pore space, and water holding capacity of soil.

| Treatments | BD (Mg m ⁻³) | | PD (Mg m ⁻³) | | Pore space (%) | | WHC (%) | |
|-----------------------------------|--------------------------|----------|--------------------------|----------|----------------|----------|---------|----------|
| | 0-15 cm | 15-30 cm | 0-15 cm | 15-30 cm | 0-15 cm | 15-30 cm | 0-15 cm | 15-30 cm |
| NPK ₀ FYM ₀ | 1.32 | 1.35 | 2.52 | 2.53 | 43.79 | 41.54 | 40.19 | 39.27 |

| | | | | | | | | |
|---------------------------------------|------|------|------|------|-------|-------|-------|-------|
| NPK ₀ FYM ₅₀ | 1.29 | 1.31 | 2.47 | 2.52 | 45.29 | 43.92 | 42.10 | 40.7 |
| NPK ₀ FYM ₁₀₀ | 1.26 | 1.28 | 2.44 | 2.48 | 46.45 | 44.93 | 43.09 | 41.87 |
| NPK ₅₀ FYM ₀ | 1.31 | 1.33 | 2.51 | 2.54 | 44.43 | 42.84 | 41.28 | 40.02 |
| NPK ₅₀ FYM ₅₀ | 1.28 | 1.30 | 2.46 | 2.50 | 45.56 | 44.55 | 42.27 | 41.03 |
| NPK ₅₀ FYM ₁₀₀ | 1.26 | 1.28 | 2.43 | 2.46 | 47.16 | 45.76 | 44.01 | 42.74 |
| NPK ₁₀₀ FYM ₀ | 1.30 | 1.32 | 2.50 | 2.52 | 44.80 | 43.41 | 41.77 | 40.54 |
| NPK ₁₀₀ FYM ₅₀ | 1.27 | 1.29 | 2.45 | 2.50 | 46.12 | 44.52 | 42.76 | 41.49 |
| NPK ₁₀₀ FYM ₁₀₀ | 1.25 | 1.27 | 2.42 | 2.44 | 47.70 | 46.52 | 44.49 | 43.08 |

Table 3: Response of NPK and FYM levels on pH, EC (dSm⁻¹) and organic carbon (%) of soil

| Treatments | pH | | EC (dSm ⁻¹) | | OC (%) | |
|---------------------------------------|---------|----------|-------------------------|----------|---------|----------|
| | 0-15 cm | 15-30 cm | 0-15 cm | 15-30 cm | 0-15 cm | 15-30 cm |
| NPK ₀ FYM ₀ | 7.24 | 7.26 | 0.29 | 0.27 | 0.35 | 0.34 |
| NPK ₀ FYM ₅₀ | 7.19 | 7.25 | 0.32 | 0.30 | 0.44 | 0.42 |
| NPK ₀ FYM ₁₀₀ | 7.03 | 7.08 | 0.34 | 0.32 | 0.52 | 0.50 |
| NPK ₅₀ FYM ₀ | 7.22 | 7.25 | 0.31 | 0.28 | 0.38 | 0.37 |
| NPK ₅₀ FYM ₅₀ | 7.17 | 7.18 | 0.32 | 0.31 | 0.46 | 0.44 |
| NPK ₅₀ FYM ₁₀₀ | 6.98 | 7.01 | 0.35 | 0.32 | 0.52 | 0.51 |
| NPK ₁₀₀ FYM ₀ | 7.2 | 7.23 | 0.30 | 0.30 | 0.41 | 0.40 |
| NPK ₁₀₀ FYM ₅₀ | 7.13 | 7.15 | 0.33 | 0.30 | 0.48 | 0.46 |
| NPK ₁₀₀ FYM ₁₀₀ | 6.95 | 6.98 | 0.36 | 0.33 | 0.54 | 0.52 |

Table 4: Response of NPK and FYM levels on Available Nitrogen (kg ha⁻¹), Phosphorus (kg ha⁻¹) and Potassium (kg ha⁻¹) of soil

| Treatments | N (kg ha ⁻¹) | | P (kg ha ⁻¹) | | K (kg ha ⁻¹) | |
|-----------------------------------|--------------------------|----------|--------------------------|----------|--------------------------|----------|
| | 0-15 cm | 15-30 cm | 0-15 cm | 15-30 cm | 0-15 cm | 15-30 cm |
| NPK ₀ FYM ₀ | 224.84 | 221.25 | 21.51 | 18.41 | 188.39 | 185.82 |

| | | | | | | |
|---------------------------------------|--------|--------|-------|-------|--------|--------|
| NPK ₀ FYM ₅₀ | 228.54 | 225.35 | 22.68 | 19.62 | 192.28 | 188.46 |
| NPK ₀ FYM ₁₀₀ | 234.42 | 229.20 | 23.56 | 20.23 | 194.52 | 192.09 |
| NPK ₅₀ FYM ₀ | 239.35 | 235.78 | 24.47 | 21.85 | 197.25 | 195.32 |
| NPK ₅₀ FYM ₅₀ | 243.62 | 240.85 | 24.93 | 23.03 | 200.66 | 198.24 |
| NPK ₅₀ FYM ₁₀₀ | 246.81 | 243.21 | 25.48 | 23.67 | 203.59 | 201.95 |
| NPK ₁₀₀ FYM ₀ | 252.12 | 249.35 | 26.01 | 24.50 | 206.54 | 204.12 |
| NPK ₁₀₀ FYM ₅₀ | 256.93 | 252.97 | 26.64 | 25.18 | 209.32 | 206.35 |
| NPK ₁₀₀ FYM ₁₀₀ | 258.46 | 257.58 | 27.18 | 25.40 | 211.39 | 208.94 |




Fig: 1. Response of NPK and FYM levels on Available Nitrogen (kg ha^{-1}), Phosphorus (kg ha^{-1}), Potassium (kg ha^{-1}) of soil

CONCLUSION

The results of experiment concluded as the application of NPK and FYM was found to improve soil health references to carrot. Treatment T₉ (NPK₁₀₀FYM₁₀₀) was found optimal for improving soil properties like Pore space, Water holding capacity, organic carbon, pH, Electrical conductivity and Available Nitrogen, Phosphorus, Potassium of soil.

REFERENCE

- Ahmad, T., Amjad, M., Iqbal, Q., Nawaz, A. and Iqbal, Z. (2014).** Integrated nutrient management practices improve growth and yield of carrot (*Daucus carota* L.). *Bulgarian J. Agric. Sci.*, 20 (6):1457-1465.
- Yadav, S. K., Khokar, U.V. and Yadav, R.P. (2010).** Integrated nutrient management for carrot (*Daucus carota* L.) cultivation. *Indian J. Horti.*, 67 (4):445-449.
- Singh, B.N., Singh, A.P., Singh, T. and Singh, N.K. (2007).** Integrated nutrient management in carrot (*Daucus carota* L.). *Prog. Agric.*, 7 (1/2):84-86.
- Verma, B. K., Verma, R. K., Yadav, R. S., & Yadav, A. K. (2018).** Role of organic amendments for soil health improvement. *Journal of Pharmacognosy and Phytochemistry*, 7(4), 1481-1485.
- Walkley, A. J. and Black, J. A. (1934)** An examination of method for determination of organic carbon and nitrogen in soil. *J. Agric. Sci. (England)*, **25**: 589-609
- Olsen, S. R., Cole, C.V., Watanable, F.S. and Dean, D. C. (1954)** Estimation of available phosphorus in soil by extraction with sodium bicarbonate. *Circ. U.S Dept. Agric.*
- Balloom, A.F., Balloom, M.A. and Qayyum, S.M. (1993).** Influence of phosphorus and potassium fertilizer levels with standered dose of nitrogen on the productivity of carrot (*Daucus carota* L.) *Sharhad J. Agric.*, 1:(21-25)
- Bhattarai, B. P. and Maharjan, A. (2013).** Effect of organic nutrient management on the growth and yield of carrot (*Daucus carota* L.) and soil fertility status, *Nepalese J. Agric. Sci.*, 11:16-25.
- Kumar, P., Meghwal, P. R. and Painuli, D. K. (2014).** Effect of organic and inorganic nutrient sources on soil health and quality of carrot (*Daucus carota* L.). *Indian J. Hort.*, 71 (2):222-226.

- Sharma, A., Sharma, R.P., Sood, S. and Sharma, J.J. (2003).** Influence of integrated use of nitrogen, phosphorus, potassium and FYM on the yield attributing traits and marketable yield of carrot (*Daucus carota* L.) under the high hills dry temperate conditions of north western Himalayas. *Indian J. of Agril. Sci.*, 73 (9):500-503.
- Sunandarani, N and Mallareddy, K. (2007).** Effect of different organic manures and inorganic fertilizers on growth, yield and quality of carrot (*Daucus carota* L.). *Karnataka Journal of Agricultural Sciences*, 20 (3):686-688.
- Uddin, A.S.M.M., Hoque, A.K.M.S., Shahiduzzaman, M., Sarker, P.C., Patwary, M.M.A. and Shiblee, S.M.A. (2004).** Effect of nutrients on the yield of carrot (*Daucus carota* L.). *Pakistan Journal of Biological Sciences*, 7 (8):1407-1409.
- Vithwel and Kanaujia, S.P. (2013).** Integrated nutrient management on productivity of carrot (*Daucus carota* L.) and fertility of soil. *SAARC J. Agri.*, 11 (2):173-181.
- Zakir, H.M., Sultana, M.N. and Saha K.C. (2012).** Influence of commercially available organic vs inorganic fertilizers on growth yield and quality of carrot (*Daucus carota* L.), *jour. Environ. Sci. & natural resources*, 5 (1):39-45.