

Phenotypic characterization of plant growth promoting microbial isolates from rice rhizosphere and phyllosphere

Abstract

The present investigation was carried out during 2021-22 at Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya (Uttar Pradesh). Results revealed that phenotypic characterization of plant growth promoting microbial isolates from rice rhizosphere and phyllosphere have significant effects on physicochemical properties of soil. On the other hand, microbial isolates have a significant effect on microbial biomass carbon (MBC) and total protein content of the soil. Applying microbial isolates has a positive impact on microbial population in terms of total rhizospheric bacterial population ($0.77 - 8.4 \times 10^6$ cfu g⁻¹ soil). Whereas, fungal population ($1.0 - 51.0 \times 10^3$ sfu g⁻¹ and actinomycetes ($1.7 - 97 \times 10^3$ cfu g⁻¹ of soil). Whereas, total number of phyllospheric bacterial population of rice plant leaves and clumps were $0.53 - 6.4 \times 10^6$ cfu g⁻¹, leaf fungal population $1.0 - 73 \times 10^2$ sfu g⁻¹, leaf Actinomycetes population $0.67 - 1.2 \times 10^3$ cfu g⁻¹. A total of 6 bacterial stains, 2 Actinomycetes and 3 fungal strains were isolated from the rhizospheric soil. Similarly, 4 bacterial stains, 2 Actinomycetes and 2 fungal strains were also isolated from the phyllosphere of the rice field by serial dilution effect on plating techniques. Plant growth promoting (PGP) traits were evaluated. The highest IAA (Indole Acetic Acid) production ($23.75 \mu\text{g ml}^{-1}$) was recorded under the RRS5 (Rice Rhizospheric Sample 5) and followed by RRS3 which was produced at $18.76 \mu\text{gml}^{-1}$. The isolated microorganisms can be used as an effective bio inoculant either individually or in different combinations for the formulation of different multi potent biofertilizers for plant growth promotion substances as well as control of plant diseases in rice crop.

Keywords: Actinomycete, Bacteria, Fungi, Phyllosphere, Plant growth promoting traits, Rice, Rhizosphere

Introduction

Rice (*Oryza sativa* L.) is an important staple cereal crop in India. It is cultivated on an area of 45 million hectare with a total production of 122 million tonnes and average productivity of 4.08 million tonnes ha⁻¹. In Uttar Pradesh, rice is growing on 5.74 million ha area with a total production of 15.52 million tonnes/year and average productivity (27.04 q ha⁻¹). But comparatively it is lower than other states like Punjab (40.35 q ha⁻¹), Haryana (33.34 q ha⁻¹) and West Bengal (28.51 q ha⁻¹). Therefore, there is an urgent need to enhance productivity of this crop in the state by using beneficial microorganisms in Rhizosphere and Phyllosphere (USDA 2022). Rice plays a vital role in national food grain supply and contributed about 43% of total food grain production and 46% of total cereal production of the country. It is mainly grown in states like Assam, West Bengal, Uttar Pradesh, Haryana, Punjab, Andhra Pradesh, Tamil Nadu and in other states of the country. It is cultivated globally across the divergent agro-ecosystems and requires divergent climatic and edaphic conditions for its better growth. Therefore, there is an

urgent need to increase the yield of this crop through development of plant growth promoting isolates which can promote enzymatic activities in soil horizons. Diverse groups of microorganisms colonize the rhizospheric soils as well as the phyllosphere region of rice plant, including those that colonize the zone around the root (rhizosphere), some others that dwell on the root surfaces (rhizoplane) as well as those microbes found on the aerial parts (phyllosphere) (Knief *et al.*, 2011 and Dong *et al.* 2019) have significant role on rice plant growth and development. Plant growth promoting rhizobacteria (PGPR) play a crucial role in stimulating plant growth substances in several ways including solubilization, mineralization and fixation of nutrients, phytohormone production, and pathogen suppression (Gupta *et al.* 2015). The groups of bacteria that are associated with plant roots have a vital role on the rice plant development and yield (Mir, *et al.* 2021).

Materials and method

A Field experiment was conducted at Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya (Uttar Pradesh) during kharif season of 2021-22, which is located at 81.837461 °N latitude and 26.545184 °E longitude and mean sea level height 322 feet. Samples from rice rhizosphere and phyllosphere were randomly collected from each site. Twelve samples were collected from the entire paddy field at 70 days after transplanting (DAT). Samples were collected just before the fruiting stage in the crop. Whole paddy plant was carefully uprooted along with adhering soil, without breaking the secondary and tertiary roots and followed by chopping off the whole shoots. The samples of shoots and roots of the rice plant along with the adhering rhizospheric soil was placed into separate polythene bags and then labelled and tied in order to minimize the evaporation losses. The roots were shaken to dislodge and separate loosely adhering soil aggregates around primary, secondary and tertiary roots and the adhering soil was collected and stored in a refrigerator at 4°C temperature. The shoot samples were directly taken in laminar flow hoods and the leaves and stem samples cut into small pieces with the help of surface sterilized scissors and imprinted aseptically into nutrient agar and PDA (Potato Dextrose Agar) plates were kept separately for screening of rice phyllosphere bacteria, fungi and actinomycetes, respectively. Soil samples were analysed for determination of pH, EC, organic carbon content, microbial biomass carbon (MBC) and total protein content, a portion from each stored composite soil sample was taken for air drying, grinding followed by passing through 2 mm sieves before physico-chemical analysis of the soil. Collected soil samples of Rhizospheric were estimated for the counting of microbial population through using methods described by Aneja (2018). By preparation of serial dilutions up to 10⁻⁸ and further plating into culture media plates specific for bacteria, fungi and actinomycetes. Viable microbes were developed into colonies or spores on respective media after incubation. The number of colonies was the same as the number of organisms contained in the sample multiplied by the dilution factor.

$$\text{Organisms per g of soil samples} = \frac{\text{Number of colonies (Average of 3 replicated)}}{\text{Amount plated X Dilution factor}}$$

The following common culture media was used for plating of serially diluted rhizospheric and phyllospheric samples for counting of bacterial colonies, fungal spores and actinomycetes spores. The culture media was used for plating the dilutions and for isolating, purification and sub-culturing of microbial colonies were listed as their respective nutrient composition. The common culture media was used for plating of serially diluted rhizospheric and phyllospheric samples for counting of bacterial colonies, fungal spores and actinomycetes spores.

Results and Discussion

The highest total bacterial count (8.4×10^6 cfu g⁻¹ soil) was recorded in RRS6 treatment. Whereas, RRS10 treatment had the lowest number of bacterial populations (0.77×10^6 cfu g⁻¹ soil). However, the highest fungal population (51×10^3 sfu g⁻¹ soil) was found in the RRS2 treatment while the least number of fungi (1×10^3 sfu g⁻¹ soil) were observed in RRS1 treatment. The actinomycetes population in the tested soil samples ranges between 97×10^3 cfu g⁻¹ with RRS3 to 1.7×10^3 cfu g⁻¹ soil with RRS6. Similarly, in phyllosphere samples, the maximum bacterial population was recorded (6.4×10^6 cfu g⁻¹ sample) in RPS4 and the lowest bacterial count was in sample of RPS7 (0.53×10^6 cfu g⁻¹). The maximum fungal population was recorded (32×10^2 sfu g⁻¹) in RPS2 and the lowest fungal count was observed in RPS1 (1×10^2 sfu g⁻¹) sample. But the actinomycetes population in casein starch agar plates was varied among the samples with no colonies detected in plates which were prepared with the samples like RPS1, RPS3, RPS6 and RPS7, respectively. Among the sample plates of actinomycetes could not be detected. However, the highest number of actinomycetes were noticed in RPS4 (1.2×10^3 cfu g⁻¹ sample) and lowest was observed in RPS2 (0.67×10^3 cfu g⁻¹ sample). The variation in microbial load of rhizospheric soil samples has been found due to addition of inputs with varying levels of organic ingredients and other soil inputs (Mukherjee *et al.* 2021). The low microbial load and their diversity in the phyllosphere may be attributed due to biocontrol activities of the resident phyllosphere bacteria, fungi and actinomycetes. Etesami (*et al.* 2017) also isolated gram-positive bacteria from the rhizosphere and inside the roots of rice; they characterized them for their plant growth promoting (PGP) traits and antifungal activity against some rice plant pathogenic fungi. The results showed the endophytic and rhizosphere isolates had different PGP traits and antifungal activity. Similar findings have been also observed by Rani *et al.* 2021. Only one rhizosphere isolates and one endophytic isolate showed highly inhibitory effects against the mycelial growth of all fungal rice pathogens tested. The best bacterial isolates, based on multiple PGP traits and inhibitory effects against the mycelial growth of all fungal rice pathogens, were identified. Krishna (*et al.* 2012) have shown that the population of bacteria, fungi and actinomycetes decreased significantly with increased soil depth. The depth wise profile of total microbial abundance was well matched with the vertical profiles of soil nutrients and organic carbon. Microbial numbers were highest in surface soil where organic matter and nutrients were highest as reported by Wickramasinghe (*et al.* 2021). Further, the presence of more plant roots in the top soil had a strong effect on microbial load.

Table 1 Microbial population in soil and leaf samples

Treatment	Soil			Treatment	leaf		
	Bacteria (cfu g ⁻¹ soil)	Fungi (sfu g ⁻¹ soil)	Actinomycetes (cfu g ⁻¹ soil)		Bacteria (cfu g ⁻¹ soil)	Fungi (sfu g ⁻¹ soil)	Actinomycetes (cfu g ⁻¹ soil)
RRS1	1.3×10 ⁶	1×10 ³	6×10 ³	RPS-1	1.0×10 ⁶	1×10 ²	ND
RRS2	6.0×10 ⁶	51×10 ³	50×10 ³	RPS-2	4.2×10 ⁶	32×10 ²	0.67×10 ³
RRS3	2.5×10 ⁶	26×10 ³	97×10 ³	RPS-3	2.0×10 ⁶	21×10 ²	ND
RRS4	ND	26×10 ³	6.5×10 ³	RPS-4	6.4×10 ⁶	6.0×10 ²	1.2×10 ³
RRS5	7.2×10 ⁶	3×10 ³	8.1×10 ³	RPS-5	0.58×10 ⁶	21×10 ²	ND
RRS6	8.4×10 ⁶	11×10 ³	1.7×10 ³	RPS-6	0.53×10 ⁶	24×10 ²	ND
RRS7	0.78×10 ⁶	32×10 ³	2.2×10 ³	RPS-7	0.62×10 ⁶	73×10 ²	ND
RRS8	0.82×10 ⁶	30×10 ³	2.4×10 ³	RPS-8	2.9×10 ⁶	30×10 ²	1.18×10 ³
RRS9	0.81×10 ⁶	44×10 ³	2.1×10 ³	RPS-9	3.3×10 ⁶	26×10 ²	1.05×10 ³
RRS10	0.77×10 ⁶	50×10 ³	1.8×10 ³	RPS-10	2.3×10 ⁶	24×10 ²	ND
RRS11	0.82×10 ⁶	41×10 ³	2.2×10 ³	RPS-11	1.8×10 ⁶	18×10 ²	1.07×10 ³
RRS12	0.98×10 ⁶	36×10 ³	2.0×10 ³	RPS-12	2.6 x 10 ⁶	12 x 10 ⁶	0.74×10 ³

Where, ND: No colonies detected on the plates




Fig 1 General view of Bacteria, Fungal and Actinomycetes isolates from the rice

Conclusions

It is showed that rhizospheric samples have the highest bacterial count (8.4×10^6 cfu g⁻¹ soil) and actinomycetes (97×10^3 cfu g⁻¹ soil) in RRS6. While the Phyllosphere samples have

highest bacterial count (4.2×10^6 cfu g⁻¹) in RPS2. Maximum fungal population in plant sample (73×10^2 sfu g⁻¹) was perceived in RPS7 sample. A total of 6 bacterial, 2 actinomycetes and 3 fungal strains were isolated from rhizospheric soil samples. Besides, 4 bacterial, 2 actinomycetes and 2 fungal strains were also isolated from Phyllosphere samples of rice crop through serial dilution plating technique.

Disclaimer

This paper is an extended version of a Thesis document of the same author.

The Thesis document is available in this link:
<https://shodhganga.inflibnet.ac.in:8443/jspui/handle/10603/442738>

REFERENCES

- Aneja K R. 2018. Laboratory manual of microbiology and biotechnology. 2nd edition. New age International Publishers, New Delhi. Pp:242-70.
- Dong C J, Wang L L, Li Q, Shang, Q M. 2019. Bacterial communities in the rhizosphere, phyllosphere and endosphere of tomato plants. *PloS one*: 14(11): e0223847.
- Etesami H, Alikhani H A. 2017. Evaluation of gram-positive rhizosphere and endophytic bacteria for biological control of fungal rice (*Oryza sativa* L.) pathogens. *European Journal of Plant Pathology* 147(1): 7-14.
- Gupta G, Parihar S, Ahirwar N, Snehi S, Singh V .2015. Plant growth promoting Rhizobacteria (PGPR): current and future prospects for development of sustainable agriculture. *Journal of Microbial & Biochemical Technology* 32(6):1-29.
- Knief C, Delmotte N, Vorholt J A 2011. Bacterial adaptation to life in association with plants-A proteomic perspective from culture to in situ conditions. *Proteomics*. 11(15): 3086-05.
- Krishna M P, Rinoy Varghese, Hatha M A A. 2012. Depth-wise variation of microbial load in the soils of the midland region of Kerala: a function of important soil physicochemical characteristics and nutrients. *Indian Journal of Education and Information Management* 3: 126-129.
- Mir M I, Hameeda B, Quadriya H, Kumar B K , Ilyas N, Zuan A T K , Sayyed R Z. 2021. Multifarious Indigenous Diazotrophic Rhizobacteria of Rice (*Oryza sativa* L.) Rhizosphere and Their Effect on Plant Growth Promotion. *Frontiers in Nutrition*. 8: 781764.

Mukherjee B, Roy S, Dhara A, Dutta, S. 2021. Screening and optimization of indole-3-acetic acid production by bacterial strain isolated from the rice rhizosphere and its effects on plant growth. *Environmental and Experimental Biology*. 19: 141–149.

Rani V, Bhatia A, Nain L, Tomar G S, Kaushik R. 2021. Methane utilizing plant growth-promoting microbial diversity analysis of flooded paddy ecosystems of India. *World Journal of Microbiology and Biotechnology*. 37(4):1-22.

Wickramasinghe W R K D W K V, Girija D, Gopal K S. 2021. Native Multiphasic Plant Growth Promoting Rhizobacteria Consortia for Growth and Yield of Rice. *Research Journal of Agricultural Sciences* 12(2): 601–606.

USDA. 2022. World Agriculture Production United States Department of Agriculture.

UNDER PEER REVIEW