

## Original Research Article

### Evaluation of Rice (*Oryza sativa*) under Agro-climatic Zones of Prayagraj

Formatted: Font: Italic

Comment [AM1]: I suggest change the title: Comparative evaluation of rice (*Oryza sativa*) hybrids under Prayagraj agro-climatic conditions

#### ABSTRACT

A field experiment was conducted at Crop Research Farm, Naini Agriculture Institute, Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during *Kharif*, 2022 on sandy loamy soil. The experiment was laid out in Randomized Block Design, replicated thrice, consisting of ten treatments using the rice hybrid cultivars: *i.e.*, R-24, R-40, R-48, R-52, R-77, R-107, R-111, R-120, R-127, R-145. The field experiment results revealed that R-48 has significantly increased the growth parameters *viz.*, Plant height (111.25 cm), Number of tillers (14.50), Plant dry weight (38.43 g/plant), and yield attributes as Number Tillers/m<sup>2</sup> (338.65), Panicle length (29.60 cm), Grain yield (5.16 t/ha) and Straw yield (13.31 t/ha). Also maximum gross return (₹116264 /ha), net return (₹67218/ha), and B:C benefit cost ratio (1.37). Therefore, the rice hybrid ~~was also recorded in~~ R-48 ~~would be able recommended to farmers after further trails.~~

**Keywords:** *Rice, Hybrid, Rice, Growth, Yield, Kharif, Summer, Uttar-Pradesh-*

#### 1. INTRODUCTIONntroduction:

~~“Rice (*Oryza sativa* L.)”~~ The Rice is one of the major staple cereals with more than 3.5 billion people depending on rice for more than 20 % of their daily calorie intake (IRRI, Africa Rice and CIAT, 2010). It is the most crucial cereal food crop of India, which occupies about 24% of gross cropped area of the country. It contributes 42% of total food grain production and 45% of total cereal production of the country. China and India are major rice growing countries in the world. Krishnakumar *et al.*, (2005) have reported that in India, rice is cultivated on an area of 42.2 million hectares with a production of 110.9 million tons of rice, while China produces 187.45 million tons of rice on the area of 33.1 million hectares. India has a long history of rice cultivation and stands first in rice area and second in rice production, after China (Yadav *et al.*, 2010).

Comment [AM2]: Find the manner to cite in References.

The genetic classification of rice plant belongs to the genus *Oryza* L., of family Poaceae Graminae (Graminae Poaceae). The *Oryza* genus ~~contains~~ ~~is divided into~~ four species complexes: *O. sativa* L., *O. officinalis* Wall., *O. ridley* Hook. f., and ~~the species~~ *O. granulata* Nees & Arn. ex G. Watter. There are three *Oryza sativa* types: Indica, Japonica and Javanica. The Indica rice type (*O. sativa* subsp. ~~Indica~~ *Shig. Kato*) ~~is are~~ grown throughout the tropical and sub-tropical region, ~~The~~ Indica rice ~~is are~~ grown in around 90% of world rice area. ~~and~~ The Japonica rice type (*O. sativa* subsp. *japonica* Shig. Kato) ~~is varieties are~~ grown throughout the temperate zone, ~~and~~ The Javanica rice type ~~is the name given to~~ *O. sativa* subsp. *japonica* ~~are~~ growing mainly in the part of Indonesia ~~lines~~.

Formatted: Font: Not Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Not Italic

Formatted: Font: Italic

Formatted: Font: Not Italic

Comment [AM3]: I suggest to put this paragraph beginning the INTRODUCTION, Then, continuous with The rice is one of the major staple cereals with more than.....

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Not Bold

Comment [AM4]: Find the manner to cite in References.

Comment [AM5]: It is not cited in References.

According Directorate of Economics and Statistics (2017- 2018) India is ~~the~~ major rice growing country in world with an area of 43.79 million hectares, having production 112.91 million tonnes and productivity of 2.572 t/ha (Directorate of Economics and Statistics 2017- 2018). In Uttar Pradesh, 5.9 million ha and production 13.27 million tonnes with an average productivity of 2447 kg/ha and production of 14.63 million tonnes (Agriculture Statistics, 2016). It is estimated that the rice production must increase by 114 million tons by 2035, but farmers must achieve it under significant threats from climate change (Suzanne et al., 2012) coupled with decreasing amount of available agricultural land, labour, and water for agriculture and increased costs of all inputs. Increasing global food production with minimal adverse impact on resources and the environment is the greatest challenge for food security (Ladha et al., 2015).

## 2. MATERIALS AND METHODS:

This experiment was carried out during kharif season 2022 at Crop Research Farm, Department of Agronomy, Naini Agriculture Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, (U.P.), India, which is located at 25° 28' 42" N latitude, 81° 50' 56" E longitude and 98 m altitude above the mean sea level. This area is situated on the right side of the river Yamuna by the side of Prayagraj Rewa Road about 5 km away from Prayagraj city. The climate of the region is semi-arid subtropical. The experimental field had a sandy loam texture??, pH.....?? a level of organic carbon (0.87%), available Nitrogen (225 kg/ha), Phosphorus (41.8 kg/ha) and Potassium (261.2 kg/ha). ~~The climate of the region is semi arid subtropical.~~

Comment [AM6]: If you have a meteorological service, near the place, you may be able to add data - e. g., rainfall, temperature, wind- in the year you did the field experiment. I suggest this because is important if results are compared with some repetition of this field experiment.

The experiment was conducted in Randomized block design consisting of 10 hybrids (R-24, R-40, R-48, R-52, R-77, R- 107, R-111, R-120, R-127, R-145) with three replications and was allocated randomly in each replication.

Comment [AM7]: What surface m<sup>2</sup> was used as experimental field ?; May you put a photograph ?

To reduce crop-weed competition one hand weeding was carried out at 35 days after sowing (DAS). Two irrigations were provided at 40 days interval.

The

The observations pertaining to growth attributes were recorded using standard procedure at 20 days intervals, and presented at 100 DAS. Yield parameters were observed on the day of harvest, 23<sup>rd</sup> November, 2022.

All the attributes were recorded and analyzed statistically by using appropriate analysis of variance adopting Gomez and Gomez (1984).

### 3. Experimental design

The experiment was conducted in Randomized block design consisting of 10 hybrids i.e., from R-24 to R-145 with 3 replications and was allocated randomly in each replication.

## 4. 3. RESULT AND DISSCUSSION:

### 4.1 3.1 Growth parameters

#### 4.1.1 3.1.1 Plant height (cm)

At 100 DAS the significantly and higher plant height was observed in R-107.48 (111.25 cm) (Table 1). However, R-40 (110.73 cm), R-24 (108.34 cm), R-77 (108.24 cm) and R-127 (110.28 cm) were statistically at par with R-107.305. Genetic makeup of the variety is a huge contributing factor which have also been reported by Haqueet al. (2015). Increase in plant height may be due to synchronized availability of essential plants nutrients to the crop especially nitrogen for a longer period during its growth stages. The result conformed with Deshpande and Devasenpathy, (2011). Also, reason for maximum plant height might be due to more favourable weather condition associated. Land was criticized by the higher growing degree days and hydrothermal units gained in these hybrids was found by Bahure et al. (2019).

#### 4.1.2 3.1.2 Numbers of tillers/hill

At 100 DAT the highest number of tillers was observed in R-48 (14.50) (Table 1). However, R-127 (14.30), R-240 (13.3047), and R-2440 (13.1730) were statistically at par with R-48.305. The significant differences could due to the variation in genetic make up of the high yielding varieties that might be influenced by heredity. A comparative study of the performance of rice hybrids exhibited that R-48.305 recorded significantly higher tillers/hill. The significant differences could due to the variation in genetic make-up of the high yielding varieties that might be influenced by heredity. This was consistent with Chowdhery et al. (1993).

#### 4.1.3 3.1.3 Plant dry weight (g/plant)

At 100 DAT the highest dry weight was observed in R-48 (38.43 g/plant) (Table 1). However, R-107 (37.44 g/plant), were statistically at par with R-48.305. The probable reason for maximum dry matter accumulation depends upon the photosynthesis and respiration rate, which finally increases the plant growth with respect to increased plant height, leaf area and tillers/hill, etc. Thus, the

Formatted: Indent: Left: 0", Space Before: 0 pt

Comment [AM8]: It is not cited in References.

Formatted: Indent: Left: 0.6", No bullets or numbering

Formatted: Indent: Left: 0.35", No bullets or numbering

Formatted: Normal, Indent: Left: 0.1", No bullets or numbering

Formatted: Font: 12 pt, Bold

Formatted: Font: Bold

Comment [AM9]: It is not cited in References

Comment [AM10]: It is not cited in References

Comment [AM11]: Bahure et al. (2019) found the higher growing degree days and hydrothermal units are gained by these hybrids.

Formatted: Highlight

Formatted: Indent: Left: 0.1", No bullets or numbering

Formatted: Not Highlight

Formatted: Indent: Left: 0.47", No bullets or numbering

treatment which attained maximum growth, also accumulated higher dry matter similar result have also been

reported by Kumar (2016). The other reason of high dry matter accumulation in might be due to the significant increase in morphological parameters which responsible for the photosynthetic capacity of the plant thereby increasing the straw yield. The result conformed with Bozorgi et al. (2011).

#### 4.2.4.1 3.2 Yield parameters:

##### 4.2.14.1.1 3.2.1 Number of Tillers/meter<sup>2</sup>

The highest ~~value~~tillers/m<sup>2</sup> was observed in R-48 (338.65 tillers/m<sup>2</sup>-), (Table 2). However, R-40 (320.63 tillers/m<sup>2</sup>-) ~~and~~ R- 120 (323.74 tillers/m<sup>2</sup>-) ~~were~~as statistically at par with R-48. The ~~high tillering capacity is the~~ probable reason for high yielding varieties ~~have high tillering capacity~~. Similar findings are also reported by Yadav et al. (2004). Wang et al. (2016) reported that the unequal distribution of photo-synthetically active radiation (PAR) was the source of heterogeneity in individual tiller yields, in that early emerging superior tillers pre-empted the uppermost light source, and shaded the late emerging tillers under limited light conditions.

##### 4.2.2 3.2.2 Panicle Length

~~In Table 2 is shown the R-305~~ recorded significantly higher ~~in~~ panicle length ~~4hill~~ (29.60 cm) ~~in R-48~~. However, ~~R-40 (28.23cm),~~ R-107 (29.12 cm) and ~~R-40 (28.23cm) R-111(27.59 cm)~~ were statistically at par with R-48. The nitrogen level exerted a significant ~~favourable~~ effect ~~of~~ on panicle length ~~on rice hybrids~~ rice. ~~Thus, in the study had significantly produced the longest panicle among the hybrids under experiment~~. The significant differences in panicle length among the ~~hybrid-rice hybrids varieties~~ could also be attributed to their genetic make-up. ~~R~~The results confirm the findings of Rahman et al. (2013).

##### 4.2.3 3.2.3 Grain yield (t/ha)

The data (Table 2) showed the significantly highest grain yield ~~was observed~~ in R-48 (5.16 t/ha). However, R-40 (4.73 t/ha) and R-77 (4.85 t/ha) were statistically at par with R-48. The higher grain yield/hill under variety might be due to the optimum utilization of nutrients. The hybrids of short duration ~~high yielding~~ have the potential to give the maximum grain yield ~~respect to the~~ rest of the varieties. The other reason of the high yield of ~~R-48 variety~~ is due to the better growth ~~parameters (tillers/hill, plant dry weight) attribute~~ resulting to produce higher grain yield. Similar findings were reported by ~~Ranjitha et al. (2013)~~.

##### 4.2.4 3.2.4 Straw yield (t/ha)

The data ~~in Table 2,~~ showed the significantly highest straw yield ~~was observed~~ in R- 48 (13.31 t/ha). However, R- 145 (12.25 t/ha) ~~was~~ere statistically at par with R-48. According to the findings by Padmavathi ~~;- (1997)~~, it shows that the capability of ~~hybrid-rice hybrids~~ to utilize more nitrogen ~~is~~

Formatted: Indent: Left: 0.1", No bullets or numbering

Formatted: Indent: Left: 0.1", No bullets or numbering

Comment [AM12]: It is not cited in References.

Formatted: Not Highlight

Formatted: Indent: Left: 0.1", No bullets or numbering

~~expressed through the expression of~~ better growth ~~brought by the~~ ~~with~~ beneficial effect on nutrient uptake and physiological ~~metabolism~~ ~~growth~~ increasing the straw yield.

### **3.3 Economics:**

The result showed (Table 3) that ~~{Table 3}~~ the maximum gross return (116264 INR/ha), net return (67218 INR/ha) and B:C ratio (1.37) was recorded in R-48 as compared to other ~~rice h~~ hybrids.

### **5. 4. CONCLUSION:**

The concluded experiment showed that ~~the rice~~ hybrid R-48 was found to be ~~the~~ best for obtaining maximum grain yield. It also fetched the maximum gross return, net return and ~~Benefit cost: C~~ ratio as compared to other hybrids. Since the findings are based on the research done in one season. Further trials are needed to conform more precise results. ~~The Heneerice~~ hybrid R-48 can be recommended to farmers after further trails.

Formatted: Indent: Left: 0.35", No bullets or numbering

UNDER PEER REVIEW

**Table 1. Field evaluation of ~~different varieties on~~ growth attributes of Rice Hybrids [at 100 days after sowing](#).**

S. No	Hybrids	Growth parameters <a href="#">at 100 DAS</a>		
		Plant Height (cm)	Tillers/hill (No.)	Dry weight (g/plant)
1.	R-24	108.34	13.17	33.36
2.	R-40	110.73	13.30	32.69
3.	R-48	104.97	14.50	38.43
4.	R-52	103.68	10.97	3398
5.	R-77	108.24	12.34	33.76
6.	R-107	111.25	10.43	37.44
7.	R-111	107.30	11.70	35.45
8.	R-120	104.20	10.50	34.57
9.	R-127	110.28	14.30	35.21
10.	R-145	103.20	11.97	32.93
	F-test	S	S	S
	SEm±	1.35	0.47	0.78
	CD (p=0.05)	4.01	1.39	2.34

Table 2. Field evaluation of ~~different varieties on~~ yield attributes of Rice Hybrids at 100 days after sowing.

S. No.	Hybrids	Growth Parameter <u>at 100 DAS</u>			
		<u>No. Tillers/m<sup>2</sup></u>	<u>Panicle length (cm)</u>	<u>Grain yield (t/ha)</u>	<u>Straw yield (t/ha)</u>
1.	R-24	273.81	25.28	3.14	9.46
2.	R-40	320.63	28.23	4.73	10.59
3.	R-48	338.65	29.60	5.16	13.31
4.	R-52	280.22	24.99	3.66	8.83
5.	R-77	254.22	26.29	4.85	10.17
6.	R-107	307.43	29.12	3.47	8.34
7.	R-111	308.88	27.59	3.64	9.84
8.	R-120	323.74	25.43	2.43	7.10
9.	R-127	246.36	23.20	3.37	9.98
10.	R-145	255.43	20.20	2.90	12.25
	F-test	S	S	S	S
	SEm±	9.89	0.76	0.19	0.53
	CD (p=0.05)	29.40	2.27	0.59	1.58

Formatted: Font: 12 pt, Bold

**Table 3. Field evaluation of different varieties on Economics of Rice Hybrids**

S. No.	Hybrids	Economics			
		Cost of cultivation (INR/ha)	Gross return (INR/ha)	Net return (INR/ha)	B:C ratio
1	R-24	49046	71444	22398	0.45
2	R-40	49046	106081	57035	1.16
3	R-48	49046	116264	67218	1.37
4	R-52	49046	82,222	33176	0.67
5	R-77	49046	108343	59297	1.20
6	R-107	49046	77941	28895	0.58
7	R-111	49046	82196	33150	0.67
8	R-120	49046	55085	6039	0.12
9	R-127	49046	68463	19418	0.39
10	R-145	49046	107250	58204	1.18

**Comment [AM13]:** This column without differences is not appropriated to stay in the Table. I suggest to eliminate.

**REFERENCES** [See the Journal Guide lines for authors to cite and complete the cites in References](#)

1. Bozorgi H.R., Faraji, A., Danesh R.K., Keshovarz A., Azarpour E. and Tarighi, F. 2011. Effect of plant density on yield and yield components of rice. *World applied Science journal* **12**(11):2053-2057.
2. Deepa G, Singh V, Naidu KA. 2008. Nutrient composition and physicochemical properties of Indian medicinal rice – Njavara. *Food Chem.* **2008**;106(1):165-71
3. Deshpande, H. H., and Devasenapathy, P. 2011. Effect of green manuring and organic manures on yield, quality and economics of rice (*Oryza sativa* L.) under lowland condition. *Karnataka Journal of Agricultural Sciences* **23**(2): 235-238.
4. Krishnakumar, S., R. Nagarajan, S.K. Natarajan, D. Jawahar and B.J. Pandian. 2005. NPK Fertilizers for Hybrid Rice (*Oryza sativa* L.) Productivity in Alfisols of Southern Districts of Tamil Nadu. *Asian J. Plant Sci.* 4(6): 574-576. <https://doi.org/10.3923/ajps.2005.574.576>
5. Kumar, S. N. 2016. Evaluate the establishment techniques on growth and yield of rice, *Agricultural Research Communication Centre, Agriculture Science Digest* **36**(2): 110- 113
6. Ladha JK, Rao AN, Raman AK, Padre AT, Dobermann A, Gathala M, Kumar V, Saharawat Y, Sharma S, Piepho HP, Alam MM, Liak R, Rajendran R, Reddy CK, Parsad R, Sharma PC, Singh SS, Saha A, Noor S. (2015.) Agronomic improvements can make future cereal systems in South Asia far more productive and result in a lower environmental footprint. *Glob Chang Biol* **22**: 1054–1074.
7. Marri P. R., Sarla N., Reddy I. V. and Siddiq F. A. 2005. Identification and mapping of yield and yield related QTLs from an Indian accession of *Oryza rufipogon*, *BMC : genetics* **13**: 33-39.
8. [Directorate of economics and statistics. 2017-2018](#) Ministry of Agriculture, Farmers Welfare Department of Agriculture, Cooperation and Farmers Welfare. ~~Directorate of economics and statistics. 2021-22.~~ First advance estimates of production of foodgrains for; [2021-22](#)

**Comment [AM14]:** It is not cited in text of manuscript

**Comment [AM15]:** It is not cited in text of manuscript

**Comment [AM16]:** It is cited in the text. In References you must find the indication for cite it correctly.

9. Padmavathi, P. 1997. Studies on relative performance of conventional hybrid rice varieties under various levels of nitrogen, plant population and planting patterns. phd thesis, Indian agricultural research institute, New Delhi.

**Comment [AM17]:** It is not cited in text of manuscript

10. Rahman M.M., Islam M.T., Faruq A.N., Akhtar N., Ora N. and Uddin M.M. 2013. Evaluation of Some Cultivated Hybrid Boro Rice Varieties Against BLB, ShB and ALS Diseases Under Natural Epiphytic Conditions Middle-East *Journal of Scientific Research* 15(1): 146-151

11. The probable reason for high yielding varieties have high tillering capacity. Similar findings are also reported by **Yadav et al. (2004)**. **Wang et al. (2016)** reported that the unequal distribution of photo- synthetically active radiation (PAR) was the source of heterogeneity in individual tiller yields, in that early emerging superior tillers pre-empted the uppermost light source, and shaded the late emerging tillers under limited light conditions.

**Comment [AM18]:** What is this???? It is not a bibliographic references

12. The significant differences could due to the variation in genetic make-up of the high yielding varieties that might be influenced by heredity. This was consistent with **Chowdhery et al. (1993)**.

**Comment [AM19]:** What is this?? It is not the manner of cite a bibliographic references

13. Virmani SS, Mao CX, Hardy B. Hybrid rice for food security, poverty all deviation and environmental protection, Proc. In 4 th International Symposium on Hybrid Rice, Hanoi, Vietnam; c2002-2003. pp. 407.

14. Wang, Y., Ren, T., Lu, J. W., Ming, R., Li, P. F., Saddam, H., Cong, R. H. and Li, X. K. 2016. Heterogeneity in rice tillers yield associated with tillers formation and nitrogen fertilizer. *Agronomy Journal* 108: 1717–1725.

**Comment [AM20]:** I did not find cited in the text.

15. Yadav, S. K., Suresh, B. G., Praveen, P. and Kumar. 2010. Assessment of genetic variability, correlation and path association in rice (*Oryza sativa* L.). *Journal of bioscience* 18: 1-8.

**Formatted:** Font: Not Bold