

### Original Research Article

## Effect of organic and inorganic fertilizers on the growth and yield of Tomato (*Solanum Lycopersicum*) under protected cultivation.

Comment [i1]: Delete

Comment [i2]: Delete

### **ABSTRACT**

Comment [i3]: The abstract needs to be rewritten

The present experiment was carried out during 2022 in Central Horticulture Research Farm of Department of Horticulture, SHUATS, Prayagraj. The experiment was conducted in Randomized Block Design with 10 treatments replicated thrice. The treatments were T<sub>0</sub> Recommended dose of fertilizers (RDF), T<sub>1</sub> 75% (RDF) +25% (Vermicompost), T<sub>2</sub> 50% (RDF) + 50% (Vermicompost), T<sub>3</sub> 25% (RDF) +75% (Vermicompost), T<sub>4</sub> 75% (RDF) + 25% (FYM), T<sub>5</sub> 50 % (RDF) +50% (FYM), T<sub>6</sub> 25 % (RDF) +75% (FYM), T<sub>7</sub> 75% (RDF) + 25% (Poultry Manure), T<sub>8</sub> 50% (RDF) +50% (Poultry Manure), T<sub>9</sub> 25% (RDF) +75% (Poultry Manure). On the basis of our finding the best treatment was found to be T<sub>6</sub> (25% RDF + 75% FYM) which was found best in terms of growth parameters, yield parameters and quality parameters among different treatment combinations of tomato. Also the best B:C ratio 3.84 was found in T<sub>6</sub>(25% RDF + 75% FYM).

Comment [i4]: There is no need to mention them in the abstract, it is preferable to focus on the results only

**Keywords:** *Vermicompost, FYM, Poultry manure, yield, growth, Randomized block design.*

### **INTRODUCTION**

Tomato, botanically known as *Solanum lycopersicum* L. or *Lycopersicon esculentum* Mill. belongs to the family *Solanaceae* bearing chromosome number  $2n=2X=24$  and originated from South America. It is one of the most popular and widely grown vegetable crops throughout the world and treated as "protective food" universally. It is rich source of vitamins, vegetable protein and minerals and holds a glorious position among vegetable after the potato and sweet potato. Tomato known as "love of apple in England". Tomato is used as soup, salad, pickles, ketchup, puree, sauces, tomato paste, tomato juice and other products. The pulp and juice of tomato fruit are digestible and mild aperients, a promoter of gastric secretion and a blood purifier.

Comment [i5]: Why are there no sources in the text? The researcher must rewrite the introduction according to the research conditions in terms of entering the names of the sources for each paragraph.

The tomato plants typically grow 1-3 meters in height and have a weak stem that often sprawls over the ground and vines over other plants. Flower are generally borne in clusters of 4 to 8 but small fruited type may have 30 to 50 flowers per cluster. Tomato plants are dicots,

Comment [i6]: One of the most well-liked and widely-cultivated vegetable crops, it is regarded as "protective food" everywhere. After the potato and sweet potato, it ranks highly among vegetables and is a great source of vitamins, vegetable protein, and minerals. "Love of apple in England" is the nickname for the tomato. Tomatoes are used in a variety of items, including soup, salads, pickles, ketchup, puree, sauces, tomato paste, and tomato juice. The tomato fruit's pulp and juice are easily digestible, mild aperients that encourage gastric secretion and purify the blood.

and grow as a series of branching stems, with a terminal bud at the tip that does the actual growing.

Tomato plays a major role in human nutrition, fruit contain 93.1% water, 1.9% protein, 0.3 g fat, 0.7% fibre, 3.6% carbohydrates, 23 calorie, 320 I.U vitamin A., 0.07 mg vitamin B1, 0.01 mg vitamin B2, 31 mg vitamin C, 20 mg calcium, 36 mg phosphorus and 0.8 mg iron. Tomato has valuable vitamins and cholesterol. Approximately 20–50 mg of lycopene per 100g of fruit weight can be obtained from tomato. The best fruit color and quality is obtained at a temperature range of 21-24°C. Tomato is one of the versatile crop in the world because of its fast and wide climate adaption and it is universally treated as protective food. Tomato contributes to a healthy, well balanced diet. They are rich in minerals, vitamins, essential amino acids, sugars, dietary fiber and it has many other uses tomato seed contain 24% of oil is used as salad oil and in the manufacture of margarine.

Generally, food security in world is very important because the large population and the better living standard need more food. In conventional farming, farmers using inorganic fertilizers for nutrient availability to increase crop production. However, for long term of period the usage of inorganic fertilizer could reduce soil fertility and crop productivity. Although organic manure is eco-friendly, it renders lower yield in comparison to inorganic fertilizers. In this regard, by applying organic manure in combination within organic fertilizer we can increase production as well as improve soil health. In addition, this combined application maximizes the use of available organic resources and minimizes the use of expensive inorganic fertilizers (Manral and Saxena, 2003; Ghosh *et al.*,2004). We hypothesized that incorporation of organic and inorganic fertilizers may provide better yield and quality of crop with improvement in soil health. Therefore, the present study was conducted to evaluate the performance of tomato varieties with different combinations of organic and inorganic fertilizers.

#### **MATERIALS AND METHODS: -**

The present investigation was done to understand the effect of organic and inorganic fertilizers at different doses combination on growth, yield and quality of tomato under protected cultivation variety Romon. The experiment was carried out at Horticultural Research Farm (HRF), Department of Horticulture, Naini Agricultural Institute SHUATS, Prayagraj, U.P., during the Rabiseason of 2022. The different combination doses of organic

**Comment [i7]:** The tomato is one of the most adaptable crops in the world due to its rapid and widespread climatic adaptation, and it is regarded as a food that provides protection everywhere. A diet that is healthy and well-balanced benefits from tomatoes. They are abundant in dietary fiber, carbohydrates, vital amino acids, minerals, and vitamins. Tomato seeds, which have a 24% oil content, are used to make margarine and as salad dressing.

**Comment [i8]:** In general,

**Comment [i9]:** standards require

**Comment [i10]:** In conventional agriculture, farmers use inorganic fertilizers to increase nutrient availability to increase crop production.

**Comment [i11]:** in the long term, the use of inorganic fertilizers can reduce soil fertility and crop productivity. Although organic fertilizer is environmentally friendly, it produces less than inorganic fertilizers. In this regard, by spreading organic manure with organic manure,

**Comment [i12]:** Find out

**Comment [i13]:** Tomato plants

manures and inorganic fertilizers mentioned in table 1 and replicated thrice. Observations were recorded at different stages of growth periods.

Comment [i14]: the growing season

**Physical properties of soil at Horticulture Research field (SHUATS).**

**Chemical composition of the soil**

S. No.	Particulars	Mechanical Properties
1	Sand	48.15
2	Silt	21.34
3	Clay	30.51
4	Textural class	Sandy loam

S. No.	Ingredients	Quantity
1	Soil pH	6.9
2	Organic carbon (%)	0.358
3	Available Nitrogen	212.56 kg/ha
4	Available Phosphorus	14.59 kg/ha
5	Available Potash (K)	225.1 kg/ha

?

Comment [i15]: titles are given to the tables

Notation	Treatment combination
T <sub>0</sub>	100% Recommended dose of fertilizers (RDF) ( 180:100:80)
T <sub>1</sub>	75% (RDF) +25% (Vermicompost)
T <sub>2</sub>	50% (RDF) + 50% (Vermicompost)
T <sub>3</sub>	25% (RDF) +75% (Vermicompost)
T <sub>4</sub>	75% (RDF) + 25% (FYM)
T <sub>5</sub>	50 % (RDF) +50% (FYM)
T <sub>6</sub>	25 % (RDF) +75% (FYM)
T <sub>7</sub>	75% (RDF) + 25% (Poultry Manure)
T <sub>8</sub>	50% (RDF) +50% (Poultry Manure)
T <sub>9</sub>	25% (RDF) +75% (Poultry Manure)

**Table No 1:- Treatment details:-**

Comment [i16]: It is preferable to give titles to the tables separately

**Table No 2 :- Effect of organic and inorganic fertilizer on Plant height (cm), flowering parameters and yield related traits in tomato**

Comment [i21]: Move the table to the results section

Treatment Notation	Treatment details	Plant height(120 DAT)	Days to first flowering	Days to 50% flowering	No. of flowers/cluster	Fruit set per cluster	No. of flower clusters/plant	Number of Fruit per plant	Fruit yield per plant (kg)	Fruit yield (250 sq. meter) (quantal)
T <sub>0</sub>	100% Recommended dose of fertilizer	248.18	31.05	49.11	7.45	5.75	10.02	57.61	4.65	55.79
T <sub>1</sub>	75% RDF + 25% Vermicompost	250.22	29.32	48.89	8.12	6.14	10.93	67.11	5.52	66.23
T <sub>2</sub>	50% RDF + 50% Vermicompost	250.62	29.90	46.33	8.32	6.23	10.83	67.47	5.57	66.82
T <sub>3</sub>	25% RDF + 75% Vermicompost	253.42	29.27	45.33	9.02	6.26	11.06	69.23	5.73	68.74
T <sub>4</sub>	75% RDF + 25% FYM	257.08	29.78	47.11	9.95	7.17	11.09	79.51	6.84	82.05
T <sub>5</sub>	50% RDF + 50% FYM	255.55	29.08	46.77	10.12	6.42	11.36	72.93	6.07	72.82
T <sub>6</sub>	25% RDF + 75% FYM	259.22	27.25	45.22	10.58	7.48	13.23	98.96	8.63	103.53
T <sub>7</sub>	75% RDF + 25% Poultry manure	258.98	28.08	47.21	10.25	7.19	12.96	93.18	8.03	96.33
T <sub>8</sub>	50% RDF + 50% Poultry manure	251.39	28.99	48.56	7.79	6.93	10.6	73.45	6.16	73.90
T <sub>9</sub>	25% RDF + 75% Poultry manure	254.72	30.18	48.67	10.22	6.92	12.06	83.45	7.11	85.29
<b>'F' test</b>		S	S	S	S	S	S	S	S	S
<b>S.E. (m) ±</b>		1.319	0.757	0.765	0.692	0.299	0.848	0.776	0.209	0.542
<b>C.D. at 5%</b>		2.722	1.563	1.579	1.428	0.618	1.751	1.602	0.431	1.118
<b>C.V.</b>		22.840	12.392	12.306	11.984	5.186	14.696	12.825	3.618	9.380

Comment [i18]: fruits

Comment [i20]: fruits

Comment [i19]: fruits

Comment [i17]: Delete , needless to mention it

under

protected

condition.

## Results and Discussion :-

### Plant height

The height of plant significantly varied among different treatment combinations. The maximum plant height (259.22 cm) at 30 DAT was observed with treatment T<sub>6</sub> (T<sub>6</sub>-25% RDF + 75% FYM) followed by T<sub>7</sub> (75% RDF + 25% Poultry manure) with 258.98 cm. Minimum plant height (248.18 cm) was observed in T<sub>0</sub> (100% Recommended dose of fertilizer), while the remaining treatments are moderate in their growth habit.?

Comment [i22]: The highest plant height was recorded

Comment [i23]: The lowest plant height was recorded

Comment [i24]: You need a source to prove the results

### Earliness parameter :-

#### Days to First Flowering and Days to 50% Flowering.

Maximum days to 1<sup>st</sup> flowering and 50% flowering was recorded (31.05, 49.11) in T<sub>0</sub> (100% RDF) followed by (30.18, 48.67) in T<sub>9</sub> (25% RDF + 75% Poultry manure) and minimum (27.25, 45.22) in T<sub>6</sub> (25% RDF + 75% FYM).?

Comment [i25]: You need a source to prove the results

#### No. of Flower/Cluster, Fruit set/ Cluster, No. of flower cluster/ Plant.

The effect of suitable doses of organic and inorganic fertilizer on number of flower/cluster of tomato is very obvious and consistent. There was significant difference among the doses of the different treatments, among the treatment applied the maximum number of flower/cluster was recorded in T<sub>6</sub> (25% RDF + 75% FYM) with (10.58) which was followed by T<sub>7</sub> (75% RDF + 25% Poultry manure) with (10.25) which was significantly superior over T<sub>0</sub> (100% Recommended dose of fertilizer) with (7.45) number of flower/cluster. . There was significant difference among the doses of the different treatments, among the treatment applied the maximum number of flower/cluster was recorded in T<sub>6</sub> (25% RDF + 75% FYM) with (7.48) which was followed by T<sub>7</sub> (75% RDF + 25% Poultry manure) with (7.19) which was significantly superior over T<sub>0</sub> (100% Recommended dose of fertilizer) with (5.75) fruit set/cluster. among the treatment applied the maximum number of flower/cluster was recorded in T<sub>6</sub> (25% RDF + 75% FYM) with (13.23) which was followed by T<sub>7</sub> (75% RDF + 25% Poultry manure) with (12.96) which was significantly superior over T<sub>0</sub> (100% Recommended dose of fertilizer) with (10.02) flower cluster/plant.?

Comment [i26]: Fruits

Comment [i27]: Delete

Comment [i28]: Delete

Comment [i29]: You need a source to prove the results

## Yield parameters:-

### Number of Fruit per plant

The effect of suitable doses of organic and inorganic fertilizer on number of fruit per plant of tomato is very obvious and consistent. There was significant difference among the doses of the different treatments, among the treatment applied the maximum number of fruit per plant was recorded in T<sub>6</sub> (25% RDF + 75% FYM) with (98.96) which was followed by T<sub>7</sub> (75% RDF + 25% Poultry manure) with (93.18) which was significantly superior over T<sub>0</sub> (100% Recommended dose of fertilizer) with (57.61) number of fruit per plant. Similar findings were reported by Yadav *et al.*, (2017); Majumdar *et al.*, (2018); Singh *et al.* (2018) in tomato.

Comment [i30]: Fruits

Comment [i31]: Delete

Comment [i32]: Fruits

Comment [i33]: Delete

Comment [i34]: Fruits

Comment [i35]: Fruits

### Fruit yield/Plant (kg)

The effect of suitable doses of organic and inorganic fertilizer on fruit yield per plant of tomato is very obvious and consistent. There was significant difference among the doses of the different treatments, among the treatment applied the maximum fruit yield per plant of tomato was recorded in T<sub>6</sub> (25% RDF + 75% FYM) with (8.63 kg) which was followed by T<sub>7</sub> (75% RDF + 25% Poultry manure) with (8.03 kg) which was significantly superior over T<sub>0</sub> (100% Recommended dose of fertilizer) with (4.65) kg.

Comment [i36]: Fruits

Comment [i37]: Delete

Comment [i38]: fruits

Comment [i39]: delete

Comment [i40]: delete

Comment [i41]: You need a source to prove the results

### Yield/250 sq. m (q)

The effect of suitable doses of organic and inorganic fertilizer on fruit yield per 250 m<sup>2</sup> of tomato is very obvious and consistent. There was significant difference among the doses of the different treatments, among the treatment applied the maximum fruit yield per 250 m<sup>2</sup> of tomato was recorded in T<sub>6</sub> (25% RDF + 75% FYM) with (103.53 Qt) which was followed by T<sub>7</sub> (75% RDF + 25% Poultry manure) with (96.33 Qt) which was significantly superior over T<sub>0</sub> (100% Recommended dose of fertilizer) with (55.79) Qt.

Comment [i42]: Delete

Comment [i43]: fruits

Comment [i44]: delete

Comment [i45]: delete

Comment [i46]: You need a source to prove the results

## Conclusion:-

Hence, it is concluded that T<sub>6</sub> (25% RDF + 75% FYM) was found best in terms of growth parameters and yield of tomato under protected cultivation.

**REFERENCE: -**

**Ali, M. R., Mehraj, H. and Jamal Uddin, A. F. M. 2015.** Effects of foliar application of zinc and boron on growth and yield of summer tomato. *Journal of Biological and Agricultural Research*.**06**(01): 512-517.

**Anonymous, (2015).** *Horticultural Statistics at a Glance*, Horticulture statistics division department of agriculture, cooperation & farmers welfare ministry of agriculture & farmers welfare government of India.

**Basavarajeshwari, C. P., Hosamani, R. M., Ajjappalavara, P. S., Naik, B. H., Smitha, R. P. and Ukkund, K. C. (2008).** Effect of Foliar Application of Micronutrients on Growth and Yield components of Tomato (*Lycopersicon esculentum* Mill.). *Karnataka Journal of Agricultural Sciences*.**21**(3): 428-430.

**De Candolle A.** Origin of cultivated plants. *Hafner Publication Co.*, New York, (Reprint of 2<sup>nd</sup> edition. 1959).

**Directorate of Economics and Statistics, (2020-21)** Ministry of Agriculture & Farmers Welfare (DAC & FW), Government of India, 2020-21).

**Dixit, A. Dhananjay, S., Tinku, K. S. and Pappu, L. B. (2018).** Effect of Foliar Application of Some Macro and Micronutrients on Growth and Yield of Tomato (*Solanum lycopersicum* L.) cv. ArkaRakshak. *International Journal of Current Microbiology and Applied Science*.**6**: 197-203.

**FAOSTAT (2020).** [www.fao.org/faostat/en/data/CC](http://www.fao.org/faostat/en/data/CC). Food Supply- Crops Primary Equivalent. Visited on 15/12/2021.

**sFrancis, F.J. (2000).** Encyclopedia of Food Science and Technology. John Wiley and Sons Inc.

**Gopalan B.V., Sastri Rama S.C. and Balasubramanian K.P. (2004).** Nutritive value of Indian foods. *National Institute of Nutrition, Indian Council of Medical Research*, Hyderabad, India.

**Habibullah, S. N., Saravaiya, Y. N., Tandel, S. K., Patel, B. N. and Golakiya, P. D. (2017).** Effect of Foliar Application of Micronutrients on Growth and Yield of Tomato under Protected Culture. *Trends in Biosciences*.**10**(14): 2491-2495.

**Haleema, B., Abdur, R. and Hussain, A. S. (2017).** Effect of Calcium, Boron and Zinc Foliar Application on Growth and Fruit Production of Tomato. *Sarhad Journal of Agriculture*.**30**(1): 19-30.

**Comment [i47]:** The researcher must enter the sources into the body of the research because the sources do not exist and it is not clear which source belongs to any paragraph

**Kumar, N. M., Ajay, K. P. and Amin, M. B. (2017).** Growth and Yield of Solanaceous Vegetables in Response to Application of Micronutrients – A Review. *International Journal of Innovative Science, Engineering & Technology*.**3**(2): 611-626.

**Kumar, N. S., Ahirwar, K., Khare, A. K., Mishra, U. S., Singh, D. and Jatav, R. (2016).** Effect of micronutrients and bio-fertilizers on yield and yield attributes of tomato. *International Journal of Agriculture Sciences*.**8**(38): 1780-1782.

**Meena, E. C., Maji, S. J., Meena, K., Govind, R., Kumawat, K. R., Meena, S. K. and Sodh, K. (2015).** Improvement of Growth, Yield and Quality of Tomato (*Solanum lycopersicum* L.) cv. Azad T-6 with Foliar Application of Zinc and Boron. *International Journal of Bio-resource and Stress Management*.**6**(5):598-601.

**NHB(2013).** nhb.gov.in/statistics/2013-14. Area and Production of Horticulture Crops- All India. Visited on 08/12/2021.

**Pandiyan, R., Sathyamurthy, V. A. and Pugalenti, L. (2018).** Effect of foliar application of Micronutrients on Growth, and Quality of Tomato. *International Journal of Trend in Scientific Research and Development*.**2**(4): 1988-1992.

**Reddy, G. P. D., Reddy P. S. R., Chandramohan, G. R. and Thanuja, S. G. (2018).** Effect of Foliar Application of Micronutrients on Growth and Yield Parameters in Tomato (*Solanum lycopersicon* L.). *International Journal Pure Applied Bio-sciences*.**6**(2): 929-934.

**Saravaiya, S. N., Wakchaure, S. S., Jadhav, P. B., Tekale, G. S., Patil, N. B., Dekhane, S. S. and Patel, D. J. (2014).** Influence of foliar application of micronutrients on tomato (*Lycopersicon esculentum* Mill.) Cv. "Gujarat tomato 2". *International Journal of Development Research*.**4**(8): 1539-1542.

**Sathyamurthy, V. A., Shanmugasundaram, T., Rajasree, V. and Arumugam, T. (2017).** Effect of Foliar Application of Micronutrients on Growth, Yield and Economics of Tomato (*Lycopersicon esculentum* Mill.). *Madras Agricultural Journal*.**104**(4-6): 188-193.

**Sivaiah, N. K., Swain, S. K., Sandeep, V. V. and Raju, B. (2013).** Effect of Foliar Application of Micronutrients on Growth Parameters in Tomato (*Lycopersicon esculentum* Mill.). *Discourse Journal of Agriculture and Food Sciences*.**1**(10): 146-151.

**Singh, S. Heralal, B., Suresh, C. G., Hareram, K., Manisha, J. and Upendar, K. B. (2018).** Effect of Foliar Spray of Micronutrients on Uptake of Micronutrients in (*Solanum esculentum* Mill.) cv. Navoday. *International Journal of Current Microbiology and Applied Sciences*. **7**: 930-933.

**Swetha, K., Saravanan, S. and Lalu, N. B. (2018).** Effect of micronutrients on fruit quality, shelf life and economics of tomato (*Solanum lycopersicum* L.) cv. pkm-1. *Journal of Pharmacognosy and Phytochemistry*.**7**(5): 3018-3020