

Original Research Article

Risk Factors of Mastitis and Antibiotic Susceptibility of *Staphylococcus* spp. and *Escherichia coli* isolated from cows kept by farmers in Githunguri Sub-County, Kenya

ABSTRACT

Mastitis poses a major challenge in dairy cattle farming as it leads to production and economic losses. Some mastitis causing bacteria have shown resistance to some antibiotic and the risk factors associated with the antimicrobial resistance (AMR) are not well understood. The current study assessed the prevalence and risk factors associated with occurrence of mastitis in dairy cows kept by small-holder farmers in Githunguri Sub-County, Kenya. Further, susceptibility of the isolated *Staphylococcus* spp. and *Escherichia coli* to commonly used antibiotics was evaluated. The study design was cross sectional where 91 milk samples from 40 farms were sampled from two wards in the study area. Structured questionnaires were used to assess risk factors of mastitis. The milk obtained from lactating cows was first subjected to California mastitis test (CMT) and then taken to microbiology laboratory for isolation and identification of *Staphylococcus* spp. and *Escherichia coli* using standard bacteriological tests. Antibiotics sensitivity of the isolated bacteria was examined using disc diffusion method. Based on CMT, the overall prevalence of mastitis was 50.5%. The prevalence of mastitis was associated ($p < 0.05$) with hygiene levels, being highest (69.2%) in the dairy cattle whose udder and leg hygiene were dirty. Highest prevalence of mastitis ($p < 0.05$) was found in cows who were not milked regularly (52.4%) compared to those who were milked regularly (50%) ($p < 0.05$). Cows which were cleaned using tap water had higher prevalence (53.3%) of mastitis compared to those cleaned using water from bore holes (45.2%) ($p < 0.05$). Cows that produced between 21-30 liters of milk per day had the highest prevalence (66.7%) of mastitis, while those producing between over 30 liters milk per day had lower prevalence (25%) ($p < 0.05$). The prevalence of *Staphylococcus* spp. and *E. coli* were 30.8% and 12.1%, respectively. *Staphylococcus* spp. isolates showed varied resistance to all the tested antibiotics with the highest resistance being against Oxytetracycline (57.7%) and Tetracycline (38.5%) but were sensitive to Vancomycin (100%), Ampicillin (100%), Chloramphenicol (96.2%), Penicillin (92.3%) and Gentamycin (92.3%). All the *Escherichia coli* isolates were sensitive to Chloramphenicol, Kanamycin, Gentamycin and

Oxytetracycline but were resistant to Ampicillin (88.9%) and Vancomycin (88.9%) amongst other antibiotics. In conclusion, half of the sampled cows had mastitis whose risk factors was associated with poor hygiene and irregular milking of the cows. The isolated bacteria showed varied resistance to commonly used antibiotics. To reduce the prevalence of the mastitis and AMR, there is need for extension service workers to train farmers on improved management of the mastitis.

Key words: Mastitis, prevalence, risk factor, antibiotic sensitivity, *Staphylococcus aureus*, *Escherichia coli*.

1. Introduction

Livestock production sector in Kenya contributes about 10% to Kenya's National Gross Domestic Product (GDP) and 42% to agricultural sector GDP [1]. In the country, Kiambu County is one of the leading milk producing areas and has several dairy processors [2]. In the county, the dairy sector is a source of income and employment to many people [3]. However, local extension officers in their reports have highlighted that mastitis pose a major problem in the dairy cattle industry where it causes production loss, increased cost of management, treatment and mortality in severe cases. However, the magnitude of the disease as well as risk factors have not been well studied.

In other counties in Kenya, where dairy cows are kept, the prevalence of mastitis ranges from 44% to 89% [4,5,6,7,8]. In these studies, the highest prevalence of mastitis has been observed in cows which are aged, having higher parities, poor body condition, having previous udder infections and pendulous teats of the udder. Other risk factors for mastitis include use of poor milking techniques, inappropriate milking personnel or machine and poor housing [9,10]. Thus, the strategies which have been advocated for control of mastitis include improvement of udder hygiene, treatment of subclinical cases, observation of sanitary conditions like proper cleaning of milking utensils and culling of chronic affected cows [11].

In recent years, an increase in the incidence of mastitis in the dairy herds have led to over-use of antibiotics which have resulted in emergence and spread of antimicrobial resistance [12]. Other risk factors associated with emergence of AMR include misusing and underdosing of antibiotics [13,14]. According to a previous study [15], all (100%) isolates from milk of dairy goats from Thika Sub- County in Kenya were resistant to Penicillin. Further, recent studies in some counties

in Kenya showed that bacteria isolated from milk of cows had high burden of resistance to common antibiotics [4]. Globally, this situation cuts across most of countries and thus bodies like FAO and WHO have called for increased AMR surveillance and studies to determine the risk factors associated with the phenomenon. The current study assessed prevalence of mastitis in dairy cows kept by small holder farmers in Githunguri Sub- County in Kiambu bacteria and thereafter determined the susceptibility profile of *Staphylococcus* spp. and *E. coli* to antibiotics commonly used in treatment of mastitis. The two bacteria were chosen as they have been reported to be highly prevalent in other studies in Kenya and cause severe mastitis [4].

2.0 MATERIALS AND METHODS

2.1 Study area

This study was carried out in Githunguri Sub- County, which is situated in Kiambu County between Latitude $0^{\circ} 55' 0''$ S and Longitude $37^{\circ} 4' 60''$ E. It has an elevation of 1530 meters above sea level. The area receives a mean annual rainfall of 1380mm. The average temperature of the study area is 24.6°C [16]. The sub-county covers an area of 173.50 square Kilometer and consists of five wards: Githunguri, Githiga, Ikinu, Komothai and Ngewa. It has a population of about 165,232 people. The livestock population numbers in the sub-county are 9,111 sheep, 57,008 cattle, 6,513 goats [17]. Small holder or small-scale dairy production system dominates in Githunguri area, whereby the exotic dairy cattle are mainly fed on home grown forage and commercial concentrates [18]. Most of the farmers in the area sell their milk through the Githunguri Farmers' Cooperative Society, which is the Kenya's third biggest dairy operation, producing 230,000 kilograms of milk per day [19,20].

2.2 Study design and sampling strategy

A cross-sectional survey was conducted to determine the prevalence and risk factors of mastitis, and occurrence of AMR in dairy cattle. The sample size was determined based on the formula described in [21] for low animal populations. The population of lactating cows in the study area was less than 10,000. Using the formula, the sample size was 91 lactating cows. These animals were obtained from 40 households in Githunguri and Ngewa wards. The farms were randomly

sampled from a list of farms supplied by the extension agent from Githunguri Farmers' Cooperative Society.

2.3 Milk sample collection and handling

Milk was collected aseptically from lactating cows as previously described previously [15]. California Mastitis Test (CMT) was done at the field level where a four-well plastic paddle was used. Foremilk was discarded, and then about 3ml milk was drawn into each well. An equal volume of CMT reagent was added, and gently agitated. After mixing of milk and the reagent, the result was read as 1 (negative), 2 (positive), 3 (positive) depending on the gel formation [22]. Thereafter, 10ml milk was collected into sterile universal bottles, labelled and then transported in cool box to JKUAT Medical Microbiology Laboratory for analysis.

2.4 Determination of presence of *Staphylococcus* spp. and *E. coli* in sampled milk

Culturing and identification of bacteria was done as previously described by (23, 24). The milk was cultured in Nutrient Agar (NA). The culture was incubated for 24 hours at 37⁰C. Morphological and cultural characteristics were used to identify the bacteria using a light microscope. Sub-culturing was carried out using Mannitol Salt Agar (MSA) and Eosin Methylene Blue (EMB) to isolate *Staphylococcus* spp. and *E. coli* respectively. Bacterial identification of isolated bacteria was done based on colony morphological features and gram staining reactions on pure cultures and biochemical tests such as indole, catalase and oxidase tests.

2.5 Questionnaire survey

Both open-ended and closed structured questionnaires were administered to the 40 farmers. The questionnaires composed of a series of questions which were administered to dairy farmers at farm level to determine the bio data, dairy herd size, mastitis related questions and the risk factors of mastitis including previous cases of mastitis in the farm, age, parity, body condition, and stage of lactation. The milk yield per cow was also included. Through direct observations, the udder and leg hygiene was categorized as; highly dirty, moderately dirty, slightly dirty and clean.

2.6 Data analysis

The collected data was entered into Ms Excel (Microsoft 2013, USA) and thereafter exported to a statistical package SPSS [Statistical Package for the Social Sciences, Microsoft 2013, USA) for the statistical analysis. A Chi-square test was used to evaluate relationship between risk factors and occurrence of sub-clinical mastitis ($p<0.05$).

3.0 RESULTS

3.1 Characteristics of farmers and cattle sampled

A total of 40 farmers from Githunguri and Ngewa wards of Githunguri Sub- County participated in the study. The demographic characteristics of the farmers and the sampled cattle are shown in Table 1. More women than men participated in the study and their occupation was majorly farming (85%). Majority (87.5%) of the sampled farmers had undergone formal of education. Exotic breeds were the predominant (66%) cattle breed while crosses accounted for 33.3%. The average number of cows per farm were five and were kept in farms of half acreage.

Table 1: Characteristics of respondents in Githunguri Sub-County, Kenya

Factors	Categories	Frequency	Percentage
Gender	Male	18	45.0
	Female	22	55.0
Occupation of	Business	2	5.0
	Engineer	1	2.5
	Farming	34	85.0
	Pharmacist	1	2.5
	Teacher	1	2.5
	Veterinarian	1	2.5
	Education	Primary	7
	Secondary	16	40.0
	Tertiary	12	30.0
	Informal	5	12.5

3.2 Prevalence of mastitis and identification of bacteria

California Mastitis Test was used to examine the prevalence of mastitis in the study area. The prevalence at cow level was 50.5% while the prevalence at udder level was 12.6%. Cows sampled from Githunguri Ward had a higher (29/50, 58%) prevalence compared to those from Ngewa Ward (17/41, 41.5%). The prevalence of *Staphylococcus* spp. and *E. coli* was 30.8% and 12.1%, respectively. The total number of *Staphylococcus* spp. and *E. coli* isolates were 29 and 11, respectively.

3.3 Relationship between prevalence and risk factors of mastitis

The results of CMT were used to evaluate the relationship between risk factors and prevalence of mastitis. The prevalence of mastitis was significantly ($p < 0.05$) associated with hygiene levels,

being highest (69.2%) in the cows whose udder and leg hygiene were assessed as dirty. The highest prevalence of mastitis was found in cows who were not milked regularly (52.4%) compared to those who were milked regularly (50%) ($p < 0.05$). Cows whose udder were cleaned using tap water had higher prevalence (53.3%) of mastitis compared to those cleaned using borehole water (45.2%). Cows that produced between 21-30 liters of milk had significantly ($p < 0.05$) higher prevalence (66.7%) of mastitis compared to other categories of production. Thus, udder and leg hygiene, milk yield, water source and frequency of milking were significantly ($p < 0.05$) related to the prevalence of mastitis. However, the type of floor, frequency and use of teat dips and towels, culling of chronically infected mastitis cows, presence of drainage system, provision of bedding material, production system, removal of manure and cleaning of the house (cow shed) were not significantly related to the prevalence of mastitis ($p > 0.05$).

Table 2. Relationship between risk factors and prevalence of mastitis in cows kept by small-holder farmers Githunguri Sub- County, Kenya (N=Number of sampled cattle)

Variable		N	Proportion positive	Prevalence (%)	P value
Age	1 year	10	0/10	0.0	0.500
	2-5 years	40	26/40	65	
	More than 5years	41	20/41	48.9	
Lactation stage	Early	15	6/15	40	0.794
	Mid	30	12/30	40	
	Late	46	28/46	60	
Parity	1-3	13	5/13	38.5	0.548
	4-7	42	18/42	42.9	
	7 and above	36	23/36	63.9	
Washing hands during milking	Before milking	40	19/40	47.5	0.162
	After milking	21	15/21	71.4	
	Before and after milking	30	12/30	40	
Pendulous teats	Yes	56	29/56	51.8	0.587
	No	35	17/35	48.6	
Udder leg hygiene	Clean	15	8/15	53.3	0.027
	Slightly dirty	35	18/35	51.2	
	Moderately dirty	28	13/28	46.4	
	Highly dirty	13	7/13	53.8	
Milk yield	Less than 5 liters	10	4/10	40	0.018
	5-10 liters	30	13/30	43.3	
	11-20 liters	8	5/8	62.5	

	21-30 liters	30	20/30	66.7	
	31-40 liters	4	1/4	25	
	41-50 liters	5	2/5	40	
	Above 50 liters	4	1/4	25	
Type of Floor	Concrete	72	31/72	43.1	0.215
	Muddy soil	17	14/17	82.4	
	Both	2	1/2	50	
Use of teat dips and towel	Yes	45	20/45	44.456.5	0.975
	No	46	26/46		
Frequency of using teat dips and towels	Before milking	50	30/50	60	0.696
	After milking	30	14/30	46.7	
	Before and after Milking	11	2/11	18.2	
Frequency of milking	Twice	40	20/40	50	0.030
	Thrice	30	15/30	50	
	No regular schedule	21	11/21	52.4	
Water source	Borehole	60	32/60	53.3	0.015
	Tap	31	14/31	45.2	
Provision of beddings	Yes	51	30/51	58.8	0.305
	No	40	16/40	40	
Production system	Zero grazing	70	40/70	57.1	0.773
	Open grazing	21	6/21	28.9	
House cleaning	Yes	70	38/70	54.3	0.773
	No	21	8/21	38.1	
Removal of manure	Daily	68	29/68	42.6	0.830
	Weekly	23	17/23	73.9	
Drainage System	Yes	66	26/66	39.4	0.108
	No	25	20/25	80	
Culling of chronic mastitis cows	Yes	20	5/20	25	0.484
	No	71	41/71	57.7	

3.5 Antibiotic sensitivity

Antibiotic sensitivity was tested against the isolated *Staphylococcus* spp. and *E. coli* (Table 3). *Staphylococcus* spp. isolates showed varied resistance to all the tested antibiotics with the highest resistance being against Oxytetracycline (57.7%), Tetracycline (38.5%), and Kanamycin (23.1%). Nevertheless, majority of the *S. aureus* were highly sensitive to Vancomycin, (100%), Ampicillin (100%), Chloramphenicol (96.2%), Penicillin (92.3%) and Gentamycin (92.3%). *E. coli* were shown to be more sensitive to Chloramphenicol (100%), Kanamycin (100%),

Oxytetracycline (100%), Gentamycin (100%), Oxytetracycline (100%), Tetracycline (88.9%) Streptomycin (88.9%), and Penicillin G (77.8 %) while being highly resistant against Ampicillin (88.9%) and Vancomycin (88.9%).

Table 3: Antibiotic susceptibility patterns of bacterial isolates from milk of cows kept by farmers in Githunguri Sub- County, Kenya

Antibiotics	Resistant <i>Staphylococcus</i> spp. (%)	Intermediate Resistant <i>Staphylococcus</i> spp. (%)	Susceptible <i>Staphylococcus</i> spp. (%)	Resistance <i>E. coli</i> (%)	Intermediate <i>E. coli</i> (%)	Susceptible <i>E. coli</i> (%)
Ampicillin	0(0%)	0(0%)	26 (100%)	8 (88.9%)	1 (11.1%)	0 (0%)
Chloramphenicol	0(0%)	1(3.8%)	25(96.2%)	0 (0%)	0 (0%)	9(100%)
Kanamycin	6(23.1%)	1 (3.8%)	19 (73.1%)	0 (0%)	0 (0%)	9(100%)
Penicillin G	0(0%)	2 (7.7%)	24(92.3%)	2(22.2%)	0 (0%)	7(77.8%)
Gentamycin	0 (0%)	2(7.7%)	24(92.3%)	0 (0%)	0 (0%)	9(100%)
Streptomycin	3(11.5%)	2(7.7%)	21 (80.8%)	0(0%)	1(0%)	8 (88.9%)
Oxytetracycline	15(57.7%)	1(3.8. %)	10(38.5%)	0(0%)	0(0%)	9 (100%)
Vancomycin	0 (0%)	0(0%)	26(100%)	8 (88.9%)	1(11.1%)	0 (0%)
Tetracycline	10(38.5%)	6 (23.1%)	10(38.5%)	1 (11.1%)	0 (0%)	8(88.9%)

4.0 DISCUSSION

The current study is a continuation of the ongoing studies investigating the occurrence of mastitis and AMR in milk of dairy cows kept by smallholder farmers in Kenya. In the present study, the overall prevalence of mastitis was close to that reported in Rwanda (51.8%) [25] but was lower compared to that observed in cows from other parts of Kenya (64-87.4%) [26], Ethiopia 74.7% [27] and Uganda 86.2% [28]. However, this study showed a higher prevalence of mastitis compared to those done in some parts of Kenya (36% -37%) [29,30] and Zimbabwe (21.1%) [31]. The variations in prevalence in these studies could be attributed to differences in management, environmental, epidemiological and breed factors.

In our study the burden of mastitis was found more common in cows kept and milked under poor hygienic conditions. Poor udder and leg hygiene often leads to higher microbial loads leading to

occurrence of mastitis [32]. The higher prevalence of mastitis in cows where borehole water in cleaning cow's udder could be because of water contamination by environmental bacteria such as *E. coli* [33]. The current study showed that mastitis cases were higher in late lactation stage compared to other stages and this could be attributed to prolonged exposure during the protracted lactation period [6]. As reported by others [34], cows with higher parities were associated with increased prevalence of mastitis. In our study, the peak in prevalence of mastitis was observed in cows having 21-30 litres but thereafter declined. Some studies have showed that occurrence of mastitis increases as milk yield increases [35]. In the current study, the decline in prevalence in cows having over 30 litres of milk could be due to improved management by farmers whose cows produced this amount of milk.

In our study, both *Staphylococcus* spp. and *E. coli* were isolated from the milk. However, it should be noted that our study focused on isolation of these two bacteria and there could have been other species of bacteria in the milk. *Staphylococcus aureus* is a contagious bacterium which can spread from one cow to the other through contact during unhygienic milking procedures observed in the current study. On the other hand, *E. coli* infections could be due to fecal contamination on the udder and teats [36]. *Escherichia coli* is a ubiquitous organism found in the environment surrounding dairy cow, such as bedding and borehole water that was commonly used for cleaning the udder in the study area. The bacterium invades the udder through teat, proliferates and initiates inflammatory response in dairy cows leading to mastitis [33].

The bacteria that were isolated from the milk samples were generally mostly sensitive to Gentamicin and Chloramphenicol. However, *E. coli* and *Staphylococcus* spp. showed high resistance to Vancomycin and Oxytetracycline, respectively. Similar profiles of resistance have been reported elsewhere in Uganda [37], Kenya [4,38] and Tanzania [39]. The susceptibility of *Staphylococcus* spp. and *E. coli* to Chloramphenicol and Gentamicin could be due to their infrequent use in mastitis treatment as most farmers in the study area use Penicillin (Veterinary officer, Guthunguri Sub-County personal communication). The resistance of bacteria to Penicillin may be attributed to production of beta-lactamase enzyme that inactivates Penicillin and other related antibiotics [40]. According to studies done in Tanzania [41] and Uganda [42], *S. aureus* showed high resistance to Oxytetracycline and Tetracycline. In the current study,

resistance of *Staphylococcus* spp. to Oxytetracycline and Tetracycline accounted for most of the cases which is similar to a study undertaken in Uganda [37].

Further, the present study showed that some *Staphylococcus* spp. isolates exhibited multidrug resistance pattern especially to Oxytetracycline and Tetracycline. *Escherichia coli* isolates exhibited resistance pattern to Ampicillin and Vancomycin which is similar to findings of previous studies[43, 44]. This could be attributed to the erratic and extensive use of antibacterial drugs without prior antimicrobial susceptibility testing.

5.0 CONCLUSIONS

Half of the sampled cows in the current study had mastitis whose risk factors was associated with poor hygiene and irregular milking of the cows. The bacteria isolated from the milk included *Staphylococcus* spp. and *E. coli*. To stem the occurrence of the mastitis, extension service workers should train farmers on improved management of the disease. *Staphylococcus* spp. isolates showed varied resistance to all the tested antibiotics with the highest resistance being against Oxytetracycline, Tetracycline and Kanamycin. *Escherichia coli* were shown to be highly resistant against Ampicillin and Vancomycin. Local veterinarians should consider using prudent measures to manage the occurrence of AMR especially on using the antibiotics which were shown to be active against the isolated bacteria.

REFERENCES

1. Mwanyumba PM. Pastoralist livelihoods, livestock herd dynamics and trade in Garissa County, Kenya, Ph. D Thesis, University of Nairobi, 2014.
2. Gachango FG, Andersen L M, Pedersen SM. Adoption of milk cooling technology among smallholder dairy farmers in Kenya. *Tropical Animal Health and Production*. 2014; 46(1): 179-184.
3. Muriuki HG. Dairy development in Kenya. Food and Agricultural Organization, Rome, 2011.
4. Kagira JM, Ngotho M, Mugo E, Kiplimo M, Maina N. Occurrence of antibiotic resistance in bacteria isolated from milk of dairy cows in small-holder farms in Juja Sub-County, Kenya. *Asian Journal of Research in Animal and Veterinary Sciences*. 2022; 9(3), 36-45.

5. Mureithi D, Njuguna M. Prevalence of subclinical mastitis and associated risk factors in dairy farms in urban and peri-urban areas of Thika Sub County, Kenya. *Livestock Research for Rural Development*. 2016; 28: 13.
6. Toroitich KC, Gitau GK, Kitala PM, Gitao GC. The prevalence and causes of mastitis in lactating traditionally managed one-humped camels (*Camelus dromedarius*) in West Pokot County, Kenya. *Livestock Research for Rural Development*. 2017; 29(4), 62.
7. Mbindyo CM, Gitao CG, Bebora L. A cross-sectional study on the prevalence of subclinical mastitis and antimicrobial susceptibility patterns of the bacterial isolates in milk samples of smallholder dairy goats in Kenya. *American Journal of Research and Communication*. 2014; 2(8): 30-51.
8. Nkoroi JM, Maitho T. Prevalence of mastitis and effectiveness of mastitis control in dairy cattle in Mathira Constituency, Nyeri County, Kenya. *Kenya Veterinarian*. 2014; 38(1): 24-31.
9. Mungube EO, Tenhagen BA, Kassa T, Regassa F, Kyule MN, Greiner M, Baumann MPO. Risk factors for dairy cow mastitis in the central highlands of Ethiopia. *Tropical Animal Health and Production*. 2004; 36(5): 463-472.
10. FAO. Impact of mastitis in small scale dairy production systems. *Animal Production and Health Working Paper*. No. 13, 2014.
11. Ruegg PL. A 100-Year Review: Mastitis detection, management, and prevention. *Journal of Dairy Science*. 2017; 100(12): 10381-10397.
12. Oliver SP, Murinda SE. Antimicrobial resistance of mastitis pathogens. *Veterinary Clinics: Food Animal Practice*. 2012; 28(2): 165-185.
13. Dione MM, Amia WC, Ejobi F, Ouma EA, Wieland B. Supply chain and delivery of antimicrobial drugs in smallholder livestock production systems in Uganda. *Frontiers in Veterinary Science*. 2021; 8: 611076.
14. Adekanye UO, Ekiri AB, Galipó E, Muhammad AB, Mateus A, La Ragion RM, Cook AJ. Knowledge, attitudes and practices of veterinarians towards antimicrobial resistance and stewardship in Nigeria. *Antibiotics*. 2020; 9(8): 453.
15. Mahlangu P, Maina N, Kagira J. Prevalence, risk factors, and antibiogram of bacteria isolated from milk of goats with subclinical mastitis in Thika East Subcounty, Kenya.

Journal of Veterinary Medicine, vol. 2018, Article ID 3801479, 8 pages, 2018.
<https://doi.org/10.1155/2018/3801479>.

16. Alaru, P. A. O., Shitandi, A. A., Mahungu, S. M., & Muia, J. M. K. (2022). Dairy Farming Conditions and Utilization Levels of Liquid Brewers' Yeast in Kenya. *Open Journal of Animal Sciences*, 13(1), 1-19.
17. KNBS, V. I. (2019). Population by county and sub-county. *Kenya National Bureau of Statistics*.
18. FAO. Dairy development in Kenya. Rome: Food and Agriculture Organization of the United Nations, 2011.
19. Wambugu S, Kiriimi L, Opiyo J. Productivity trends and performance of dairy farming in Kenya (No. 680-2016-46762), 2011.
20. Kiiza A. Scaling up climate change mitigation practices in smallholder dairy value chains: A case study of Githunguri Dairy Farmer Cooperative Society Ltd, Kiambu County, Kenya. MSc. Thesis. Van Hall Larenstein University of Applied Sciences, Velp. The Netherlands, 2018.
21. Thrusfield, 2007 M. Thrusfield (Ed.), *Veterinary Epidemiology* (3rd ed.), Blackwell Science, Oxford (2007), p. 624.
22. Amaral T, Ruegg P. Association between results of PortaSCC, the CMT and isolation of mastitis pathogens. Natl. Mastitis Counc. Ann. Mtg. Proc., Orlando, FL. Natl. Mastitis Counc. Inc., Madison, WI, 289-290, 2005.
23. Chesbrough M. *Distinct laboratory practice in tropical countries, part 2*. Cambridge University Press, UK; 2006:62-69.
24. Oyeleke SB, Manga SB. *Essentials of laboratory practicals in microbiology*. Minna, Nigeria: Tobest Publishers, 36-75, 2008.
25. Iraguha B, Hamudikuwanda H, Mushonga B. Bovine mastitis prevalence and associated risk factors in dairy cows in Nyagatare District, Rwanda. *Journal of the South African Veterinary Association*. 2015; 86(1): 1-6.
26. Mbindyo CM, Gitao GC, Mulei CM. Prevalence, etiology, and risk factors of mastitis in dairy cattle in Embu and Kajiado Counties, Kenya. *Veterinary Medicine International*. 2020: doi: 10.1155/2020/8831172.

27. Abebe R, Hatiya H, Abera M, Megersa B, Asmare K. Bovine mastitis: prevalence, risk factors and isolation of *Staphylococcus aureus* in dairy herds at Hawassa milk shed, South Ethiopia. *BMC Veterinary Research*. 2016; 12(1): 1-11.
28. Abrahmsén M, Persson Y, Kanyima BM, Båge R, Prevalence of subclinical mastitis in dairy farms in urban and peri-urban areas of Kampala, Uganda. *Tropical Animal Health and Production*. 2014; 46(1): 99-105.
29. Ndirangu PN, Siamba D, Wesonga HO, Mungube EO, Maichomo MW, Mugambi JM. Prevalence of bovine mastitis and multi-antibiotic resistant *Staphylococcus* and *Streptococcus* species in a research centre farm at Naivasha, Kenya. *Bulletin of Animal Health and Production in Africa*. 2017; 65(2): 291-303.
30. Gitau GK, Bundi RM, Vanleeuwen J, Mulei CM, Mastitogenic bacteria isolated from dairy cows in Kenya and their antimicrobial sensitivity. *Journal of the South African Veterinary Association*. 2014; 85(1), 01-08.
31. Katsande S, Matope G, Ndengu M, Pfukenyi DM. Prevalence of mastitis in dairy cows from smallholder farms in Zimbabwe. *Onderstepoort Journal of Veterinary Research*. 2013; 80(1): 1-7.
32. Schreiner DA, Ruegg PL. Relationship between udder and leg hygiene scores and subclinical mastitis. *Journal of Dairy Science*. 2003; 86(11): 3460-3465.
33. Cheng WN, Han SG. Bovine mastitis: risk factors, therapeutic strategies, and alternative treatments-A review. *Asian-Australasian Journal of Animal Sciences*. 2020; 33(11): 1699-1713.
34. Moosavi M, Mirzaei A, Ghavami M, Tamadon A. Relationship between season, lactation number and incidence of clinical mastitis in different stages of lactation in a Holstein dairy farm. *Veterinary Research Forum*. 2014; 5 (1): .13,
35. Siddiquee NU, Tripura TK, Islam MT,. Bhuiyan SA, Rahman AKMA Bhuiyan AKFH. Prevalence of sub-clinical mastitis in high yielding crossbred cows using Draminski Mastitis Detector. *Bangladesh Journal of Veterinary Medicine*. 2013; 11(1): 37-41.
36. Kasa G, Tegegne B, Tadesse B. Isolation and identification of major pathogenic bacteria from clinical mastitic cows in Asella Town, Ethiopia. *Veterinary Medicine International*. 2020. <https://doi.org/10.1155/2020/6656755>

37. Kabiswa W, Nanteza A, Tumwine G, Majalija S. Phylogenetic groups and antimicrobial susceptibility patterns of *Escherichia coli* from healthy chicken in Eastern and Central Uganda. *Journal of Veterinary Medicine*. 2018. doi: 10.1155/2018/9126467.
38. Ngotho M., Kagira J, Nkoiboni D, Njoroge J, Maina N. Risk factors associated with sub-clinical mastitis and antibacterial resistance in small-holder dairy farms of Kajiado North Sub-County, Kenya. *Journal of Veterinary Physiology and Pathology*. 2022; 1(3): 49-55.
39. Seyoum B, Kefyalew H, Abera B, Abdela N. Prevalence, risk factors and antimicrobial susceptibility test of *Staphylococcus aureus* in bovine cross breed mastitic milk in and around Asella town, Oromia regional state, southern Ethiopia. *Acta Tropica*. 2018; 177: 32-36.
40. Waage S, Bjorland J, Caugant DA, Oppegaard H, Tollersrud T, Mørk T, Aarestrup FM. Spread of *Staphylococcus aureus* resistant to penicillin and tetracycline within and between dairy herds. *Epidemiology and Infection*. 2002; 29(1): 193-202.
41. Caudell MA, QuinlanSubbiah MB, Call MDR, Roulette CJ, Roulette JW, Quinlan RJ. Antimicrobial use and veterinary care among agro-pastoralists in Northern Tanzania. *PloS one*, 12(1), e0170328, 2017.
42. Okubo T, Yossapol M, Maruyama F, Wampande EM, Kakooza S, Ohya K, Ushida K. Phenotypic and genotypic analyses of antimicrobial resistant bacteria in livestock in Uganda. *Transboundary and Emerging Diseases*. 2019; 66(1): 317-326.
43. Tadesse DA, Zhao S, Tong E, Ayers S, Singh A, Bartholomew MJ, McDermott PF. Antimicrobial drug resistance in *Escherichia coli* from humans and food animals, United States, 1950–2002. *Emerging Infectious Diseases*. 2012; 18(5): 741.
44. Li P, Wu D, Liu K, Suolang S, He T, Liu X, Lin D. Investigation of antimicrobial resistance in *Escherichia coli* and enterococci isolated from Tibetan pigs. *PLoS One*, 9(4), 2014; e95623.