

# Rice Farmers' Knowledge on Greenhouse Gas Emission (GHG) Mitigation Strategies; New Evidence from Nigeria

## Abstract

In Nigeria, rice remains a major staple food source for the rapidly growing population of an estimated 210 million persons. However, rice production in Nigeria is vulnerable to climate change and associated with greenhouse gas (GHG) emissions, mainly anthropogenic methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O). Understanding farmers' knowledge of GHG emission mitigation strategies would be crucial in reducing emissions from rice fields, and producing rice in a cleaner environment. However, there is a dearth of empirical evidence about the current debate. Although a few scholars have made a substantial contribution, incidentally, their studies did not pay close attention to rice farmers' knowledge level of GHG emission mitigation strategies. As a result, there is a research gap that makes it more important than ever to conduct a thorough investigation.

**Objectives:** The specific objectives of the study were to; i. describe the socio-economic characteristic of the rice farmers in the area and ii. ascertain rice farmer's knowledge level on GHG mitigation strategies

**Materials and Methods:**

The primary method used to gather data was a questionnaire and focused group discussion (FGD). Descriptive statistics and mean score analysis were used to evaluate the data.

## Results:

On average, the age of the farmer was 45 years. Most (72%) were male. Only 31% of the farmers had contacted with extension agents and were visited twice per calendar year. Among various GHG emission mitigation strategies identified in the area, the result revealed that farmers had high knowledge of the use of Aerobic Rice Varieties (ARV) ( $\bar{x} = 3.61$ ;  $\sigma = 0.87$ ) but had low knowledge of the use of Alternate Wetting and Drying (AWD) ( $\bar{x} = 1.88$ ;  $\sigma = 0.45$ ). Findings confirmed the incidence of GHG emissions in rice fields in the area and rice farmers have started practicing some of the strategies to reduce emissions. However, inadequate technical expertise is negatively challenging farmers' capacity to fully implement and scale up GHG emissions mitigation strategies.

## Conclusion:

The study recommends that the government should strengthen and support agricultural extension service systems to enable them to visit farmer's farms regularly to transmit recent GHG emission mitigation innovations. Keywords: Rice; GHG emission, CH<sub>4</sub>, N<sub>2</sub>O, GHG Mitigation Strategies, Knowledge Level, Nigeria.

## INTRODUCTION

Millions of families throughout the world depend on rice (*Oryza spp.*), the second-most popular food crop (behind wheat) to meet their dietary calorie needs [1]. To projections, the demand for rice would increase by 56% by the end of 2050 compared to the production level of 25.1 million tons in 2001 as a result of the population growth rate [2]. A large increase in rice production is required to supply this need on a worldwide scale. In Nigeria, rice is a primary staple grain and is consumed in large quantities by all households including the affluent and poor [3]. The structural rise in rice consumption over the years, with consumption spreading across all socioeconomic strata, including the poor, appears to have been caused by a confluence of many variables [4]. The rise in demand could be a result of rising income levels and population increase as well as the food's convenience in terms of preparation, storage, and calorie availability. Rice is critical in Nigeria from separate vantage points: first, in terms of the number of calories (2.06kg) it provides per person and day; 24.80 kg of calories per annum [5] and second, based on the value of income it generates through its various local production value chains. Meeting the increased demand for rice consumption in Nigeria requires increased production. Also in Nigeria, the growing demand for rice exceeds the supply, resulting in a rice deficit, with smuggled rice filling

the expanding gap between domestic production and consumption, with about 2.0 million metric tons smuggled into Nigeria by the end of 2022 [5; 6]. The efforts of the most populous nation in Africa to provide food security will be jeopardized by any reduction in rice output brought on by climate change impact and global warming. Therefore, describing how climate change affects rice production, GHG emissions mitigation strategies and farmers' knowledge are the kernels of this study. The Southeast region of Nigeria, the most populous nation in Africa, produces a significant amount of rice. The majority of the households in the area rely on rice farming as their main source of nutrition and agricultural revenue to survive [7]. Despite the fact that rice is crucial to the rural economy of Southeast Nigeria's, GHG emission from rice fields is endangering its production, health and environment. Currently, due to the present Federal Government of Nigeria Objective on diversification of the economy, reducing food insecurity and import restriction, rice is grown in almost 36 States in Nigeria including Federal Capital Territory (FCT) under diverse production systems and agro-climatic conditions [8;6]. Rice is traditionally produced in soggy paddy soils. Scientific communities have been quite concerned about rice paddies because they create the dangerous and persistent GHGs, primarily CH<sub>4</sub> and N<sub>2</sub>O [9]. Rice fields release around 30% and 11% of the world's agricultural emissions of CH<sub>4</sub> and N<sub>2</sub>O, respectively [10]. CH<sub>4</sub> global warming potential (GWP) is 34 while N<sub>2</sub>O is 298 which is more than carbon dioxide (CO<sub>2</sub>). In addition, anthropogenic N<sub>2</sub>O is thought to be the main cause of acid rain [11]. Understanding farmers' knowledge of GHG emission and mitigation strategies is among the crucial starting points for deciding what steps should be made to mitigate emissions from rice fields and produce rice in a cleaner environment. Some of the GHG emission mitigation strategies in rice field includes; alternate wetting and drying (AWD); system of rice intensification (SRI); changing tillage operations (CTO); Nitrogen Fertilizer Management (NFM); residue management (RM) and aerobic rice varieties (ARV) [12; 13]. Substantial empirical evidence [9; 11; 13] exists on measuring and mitigating GHG emissions in rice fields in developed and developing countries and different researchers have analyzed the likely effects on rice production with various parameters. Reducing GHG emissions and water use in rice fields is critical for combating climate change, and increasing yield, income and the standard of living of the farmers. A key factor in deciding what steps should be made to mitigate the adverse impacts of climate change on rice productivity and the environment is the amount of awareness that rice farmers have regarding GHG emissions. Studies on the link between GHG emissions in rice fields are increasing. However, there remains a dearth of empirical evidence on the farmers' knowledge of GHG emissions and the different mitigation strategies they practice. Therefore, the absence of this study creates a void in research and makes it increasingly pertinent that the study is systematically undertaken.

### **Objectives of the Study:**

The specific objectives of the study were to;

- i. describe the socio-economic characteristic of the rice farmers in the area and;
- ii. ascertain rice farmers knowledge level on GHG mitigation strategies

### **1. Material and Method**


The study was carried out in the Southeast agricultural zone of Nigeria from January through March 2023. The zone is made up of five States, namely, Abia, Anambra, Ebonyi, Enugu and Imo. It has an estimated land mass of 32,610 km<sup>2</sup> and a population of 22,583,076 [15]. The area

lies between longitude  $2^{\circ}61^1$  and  $6^{\circ}.32^1$  East and latitudes  $6^{\circ}74^1$  and  $8^{\circ}15^1$  North of the Equator with the mean annual temperature ranges from  $21.6^{\circ}\text{C}$  to  $32.4^{\circ}\text{C}$  while the annual rainfall ranging from 720 mm to 1440 mm in the rainforest region [16;17]. The State has good climatic conditions suitable for rice farming and a good proportion of the population are essentially rice farmers. In the selection of respondents who are rice farmers, the study used multistage and purposive random sampling techniques. Selecting farmers who are primarily engaged in rice cultivation in the region was done through purposive sampling. One hundred and fifty-two (150) rice growers made up the samples. The sample proportion and the map of the study are shown in Table 1 and Fig. 1 respectively.:-

**Table 1:Sampling Proportion for the Rice Farmers**

Southeast Agricultural Zones of Nigeria	Total Number of Government	Total Number of Local Communities	Total number of Villages	Total Number of Farmers	Total number of farmers per zone
Ebonyi	5	5	10	5	50
Imo	5	5	10	5	50
Anambra	5	5	10	5	50
<b>Total</b>	<b>15</b>	<b>15</b>	<b>30</b>	<b>30</b>	<b>150</b>

Source: Field Survey Data, 2023



**Figure 1: Map of Southeast Nigeria showing the five various States [18]**

In addition, data collected were analyzed using descriptive statistics such as mean, percentage and mean score of Likert scale type. The study used descriptive statistics to describe the socio-economic characteristics of rice farmers and the mean score of Likert scale type to evaluate rice farmer's knowledge level on GHG emission strategies in rice fields. The primary instrument

used to gather data was a structured questionnaire. The Likert scale type rating method was used to examine the data. The weighted mean calculation methodology was given as follows below:

$$X_w = \frac{\sum_{j=1}^5 ni(5i)}{n}$$

Where  $X_w$  = Weighted Mean Score

$n$  = Number of rice farmers selected

$\Sigma$  = Summation

The various attributes (GHG emission strategies) were rated using a mean score of a 4-point Likert scale type rating model for rice farmer's perceived level of knowledge on various GHG emission strategies and then divided by the number of scales to obtain the discriminating index, for example,  $(4+3+2+1)/4 = 2.50$  cut-off point. It was stated as follows;

SA = Strongly agreed (4)

A = Agreed (3)

D = Disagreed (2)

SD = Strongly Disagreed (1)

**Decision Rule;**

0.1-1.99= No Knowledge;

2.0-2.49= Moderate Knowledge;

2.50 and above = High Knowledge

The level of Significance is 0.05%

## Results and Discussion

### Socio-economic Characteristics of Rice Farmers

The socio-economic characteristics of the rice farmers in the area are shown in **Table 2**. The farmers were 45 years old on average. This suggests that the region's rice farmers were still relatively young and in the prime of their lives, giving them more chances to apply various GHG emission tactics to increase their rice output, as seen in Table 2. Age is particularly important in rice farming since it significantly affects farmers' access to and acceptance of new ideas their need for energy, and their overall production objectives [7]. Greater proportions 0.72 (72%) of the farmers were males. This implies that both sexes (male and female) are involved in rice farming but males are more in number. Nigeria practices a more paternalistic culture, making it easier for males than for women to obtain and possess agricultural productive inputs including lands, financing facilities, better seedlings, and labour. Additionally, male farmers could be able to withstand the stress and strains involved in rice farmers, and the negative impact of climate change and are able to practice more GHG mitigation strategies than their female counterparts are. The average household size in the area was 8 persons, suggesting that the rice producers had quite high households, some of which may have included direct family, relatives, or extended dependents who surely could help with rice production and practice more GHG mitigation strategies in the area. The result is in consonance with the study of [Olugbenga et al.\[19\]](#) who found that household size could be a proxy for labour to increase and enhance their farming activities, expansion of farm income and standard of living. The average length of time spent in school was 12 years, which suggests that the rice farmers in the region had at least a secondary education. Education could be very vital in positively influencing rice farmer's GHG mitigation strategies in increasing rice yield and reducing emissions in rice fields in the area. The finding is

consistent with the result of [Esiobu et al.,\[12\]](#)who asserted that higher education influences farmer’s **decisio-making** positively, acceptance of innovation, better access and utilization of productive input in reducing GHG emission and climate change impact in rice fields. The extension values of 0.31 (31%) and 2.11 revealed that about 51% of the rice farmers accessed extension services and were visited at least 2 times per year by the extension agents in the area. This is quite poor and could negatively affect rice farmers' adaptation to climate change and mitigation of GHG **emissions** in rice fields in the area. Improved and steady contact of farmers with extension agents is critical in improving farmers' access to input, innovation, high yield and income. The result supports the findings of [Esiobu et al\[12\]](#)who reported that extension contact improves farmers' access to recent information and expertise of contemporary farming techniques to raise their yield, income and standard of living. The average experience in farming was 16 years. This is a strong indication that rice farmers have reasonable years of experience in rice farming and may have been adapting to climate change and mitigating GHG **emissions** in rice fields in the area. A unit increase in farming experience is anticipated to boost farmers' practical expertise in resolving difficulties specific to rice farming and their ability to handle both internal and external obstacles influencing rice yield particularly related to climate change and GHG **emissions**. The mean farm size of the rice farmers in the area was 1.53 hectares. This is characteristic of rural farmlands, which are frequently dispersed, fragmented, and small in size. This small farm size could also be attributed to the system of land tenure and urbanization predominant in the area. Land is one of the important productive inputs in rice farming and when farmers have a limited size of it, increased farm production and mitigating GHG emission may be unattainable. The finding tallies with the studies of [Abdulwahab et al. and Tobi&Edgeweblime\[20;21\]](#) who found that large farm size increases yield, farmers adaptation to climate change, mitigating GHG emission and income of the farmers. Additionally, research shows that 0.75 (or 75%) of the rice farmers belonged to farmer clubs or associations. This suggests that rice farmers might have access to current research on climate change adaptation and GHG mitigation in rice fields through the association. Cooperative membership is seen to be a helpful tool for members to manage risks, exchange knowledge, labour, and pool their limited resources to boost yield, income and standard of living [22]. Finally, **the** finding from Table 2 indicates that the mean annual farm income from rice production was ₦480,000 (\$1,041.45). This is an indication that farmers have a relatively high annual farm income, which could positively affect farmer’s adaptation to climate change and mitigationof GHG emission in the area. Most GHG **emissions** strategies are costly [12] and farmers with sizable and sustainable farm income would be able to adapt and mitigate adequately.

**Table 2: Socio-economic Characteristics of Rice Farmers**

S/No	Variables	Mean ( $\bar{x}$ )/Percentage (%)
1	Age (years)	45.00
2	Sex (percentage of male)	0.72
3	Household size (number of persons)	8.00
4	Education (years spent in school)	12.00
5	Extension contact (percentage of access)	0.31
6	Number of extension visits (number of visits per season)	2.11
7	Farming experience (years)	16.00
8	Farm size (hectares)	1.63
9	Membership of farmer groups (percentage of members)	0.75
10	Annual Farm income (Nigerian Naira)	₦480,000 (\$1,041.45)

Source: Field Survey Data, 2023

## 1. Rice Farmer's Knowledge Level on GHG Mitigation Strategies

The outcome of rice farmers' distribution based on knowledge level of GHG mitigation strategies in the area is shown in Table 3. The various attributes were rated on a 4-point Likert type scale rating questions of Strongly Agreed (4); Agreed (3); Disagreed (2) and Strongly Disagreed (1). The values of standard deviation ( $\sigma$ ) denote the degree of variation in the responses of the rice farmers on their knowledge level in mitigating GHG emission in rice fields in the area. Reducing GHG emissions from paddy rice production is a critical target for many African countries (of which Nigeria is included) in order to comply with their climate policy commitments [12]. However, understanding farmers' knowledge of GHG mitigation strategies is key to achieving community, regional, and global climate policy commitments. Therefore, results revealed that rice farmers had high knowledge in the use of Aerobic Rice Varieties (ARV) ( $\bar{x} = 3.61$ ;  $\sigma = 0.87$ ); Residue Management (RM) ( $\bar{x} = 3.42$ ;  $\sigma = 0.60$ ); and Changing Tillage Operations (CTO) ( $\bar{x} = 3.60$ ;  $\sigma = 0.85$ ). The incorporation of rice residues contributes toward long-term nutrient cycling but may be due to high carbon-to-nitrogen ratio (C/N ratios), causing short-term Nitrogen (N) immobilization and thus affecting N availability for subsequent crops [23]. Meanwhile, aerobic rice varieties (ARV) is a production system in which specially developed "aerobic rice" varieties are grown in well-drained soils that are neither waterlogged nor puddled [24]. The technique targets a yield of at least 4-6 tons per hectare with careful management [25;26;27; 28;29]. Rice farmers' high knowledge of these strategies is expected as they may not require so much technical know-how and capital for their overall implementation. Similarly, rice farmers stated that they have moderate knowledge of the use of Nitrogen Fertilizer Management (NFM) ( $\bar{x} = 2.38$ ;  $\sigma = 0.60$ ) in mitigating GHG emissions in rice fields in the area. Differentiated emissions are produced as a result of nitrogen fertilizers' major influence on  $N_2O$  emissions [31]. Rational nitrogen fertilizer use may both encourage good rice yields and lower greenhouse gas emissions [32]. Therefore, to ensure yield and minimize  $N_2O$  loss due to nitrification and denitrification of excess nitrogen in the soil, nitrogen reduction and appropriate application can significantly increase the nitrogen usage efficiency of a rice. The timing and kind of fertilization have a significant impact on the amount of  $CH_4$  and  $N_2O$  emissions in paddy fields [33]. Ultimately, the finding shows that the rice farmers have low knowledge on the use of Alternate Wetting and Drying (AWD) ( $\bar{x} = 1.88$ ;  $\sigma = 0.45$ ); and System of Rice Intensification (SRI) ( $\bar{x} = 1.85$ ;  $\sigma = 0.48$ ) in mitigating GHG emission in a rice field in the area. The low knowledge of farmers on AWD and SRI GHG emission mitigation strategies could be associated with poor extension contact earlier found in the study. Extension agents are charged with transmitting recent innovations to farmers in their farmland to help increase their yield, income and standard of living. However, when the extension system is not well strengthened, it creates a gap between extension agents and farmers. The finding is in line with the study of Esiobu et al [12] who reported that extension contact improves farmers' access to recent information and expertise of contemporary farming techniques to raise their yield, income and standard of living. Also, the low use of AWD and SRI could also be associated with Nigeria's unsustainable rice production practices and high import dependent on rice commodities to meet demand deficient. Although the recent Federal Government of Nigeria intervention has banned rice import to boost local production (FMARD, 2022), this ban has not yet achieved the desired result. Nigeria is still not yet self-sufficient in rice. Meanwhile, the SRI is a comprehensive strategy for sustainable rice farming. Instead of planting them by the fistful and bunched closely together, it called for planting a single seedling with more space between them [34]. Additionally, it calls for spot irrigation and the tolerance of dry spells as opposed to constant flooding and the use of organic input. The SRI has the potential to reduce energy use,

GHG emissions and global warming potential (GWP) in rice-growing areas of Nigeria. The AWD was developed by IRRI in 2002 and it is one of the most significant GHG emission mitigation strategies used by farmers across Southeast Asia [35]. Farmers can use the water-saving technique of AWD to decrease irrigation water usage in rice fields without lowering production. AWD applies irrigation water a few days after the ponded water has disappeared. As a result, the field is periodically inundated and unflooded. For irrigated rice, alternate wetting and drying (AWD) is a management technique that has been developed to minimize water input [27]. AWD incorporates a number of dry spells during the course of the rice growth cycle as opposed to continually flooding the fields while the rice is being grown [36]. However, field water is maintained at a level that allows rice plants to get enough water and avoid experiencing water stress and the use of "safe" or moderate AWD has no negative effects on grain output [37; 38]. From the aggregate mean ( $\bar{x} = 2.79$ ;  $\sigma = 0.64$ ) which is above the discriminatory score ( $\bar{x} \geq 2.50$ ), it shows that the rice farmers perceived rightly their knowledge level on GHG mitigation strategies and the result was therefore accepted. Finally, there is no doubt that the relative level of knowledge of rice farmers is responsible for the low practices of GHG emission mitigation strategies in the area. Addressing these challenges will be critical in improving rice farmer's knowledge of GHG emission mitigation strategies, GHG mitigate emission from rice fields and producing rice in a cleaner environment.

**Table 3: Rice Farmers Knowledge Level on GHG Mitigation Strategies****n =150**

S/No	Items	SA	A	D	SD	Mean ( $\bar{x}$ )	SD ( $\sigma$ )	Decision
1	Changing Tillage Operations (CTO)	102 (68.00)	40 (26.67)	6 (4.00)	2 (1.33)	3.60	0.85	High Knowledge
2	System of Rice Intensification (SRI)	3 (2.00)	13 (8.67)	92 (61.33)	42 (28.00)	1.85	0.48	Low Knowledge
3	Aerobic Rice Varieties (ARV)	99 (66.00)	44 (29.33)	6 (4.00)	1 (0.67)	3.61	0.87	High Knowledge
4	Residue Management (RM)	92 (61.33)	33 (22.00)	22 (14.67)	3 (2.00)	3.42	0.60	High Knowledge
5	Alternate Wetting and Drying (AWD)	3 (2.00)	9 (6.00)	106 (70.67)	32 (21.33)	1.88	0.45	Low Knowledge
6	Nitrogen Fertilizer Management (NFM)	16 (10.67)	28 (18.67)	104 (69.33)	2 (1.33)	2.38	0.60	Moderate Knowledge
<b>Aggregate Mean Score</b>						<b>2.79</b>	<b>0.64</b>	<b>Accepted</b>

Key; SA: Strongly Agreed; A: Agreed; SD: Strongly Disagreed; D: Disagreed; SD ( $\sigma$ ); Standard Deviation; Discriminatory index: Cut off point  $\bar{x}$  Key; SA: Strongly Agreed; A: Agreed; SD: Strongly Disagreed; D: Disagreed; SD ( $\sigma$ ); Standard Deviation; Discriminatory index: Cut off point  $\bar{x} \geq 2.50$  Accepted; 0.1-1.99= Low Knowledge; 2.0-2.49= Moderate Knowledge; 2.50 and above = High Knowledge; \*Figures in parenthesis are percentage; Field Survey Data, 2023

Rice is critical to Nigeria from separate vantage points: first, in terms of the number of calories it provides per person and day; and second, based on the value of income it generates through its various local production value chains. Sustainable rice production offers promising opportunities for ensuring safe food, high yield, and income. Most of the farmers have limited knowledge of AWD. AWD is one of the major GHG mitigation strategies developed by IRRI in 2002, and has been adequately studied and propagated globally by various scholars and institutions. AWD is a promising technique that involves intermittently flooding and drying rice fields. When compared to continuous flooding, it has the potential to cut CH<sub>4</sub> emissions by an average of 48%. The study also found that rice farmers complained of limited capital. Since members of cooperatives, they are encouraged to take advantage of their strength and shared common purpose to collectively project a common demand in obtaining funds and other necessary inputs for the implementation and scaling-up of GHG emission mitigation strategies in the area. From the limitation of the study, only one hundred fifty (150) rice farmers from the study region had to be chosen for the study due to time and resource constraints on the researcher. Therefore, the findings are generally relevant to other parts of the State that were not chosen. Also, the questionnaire and in-person interviews used to acquire the data require the rice farmers to recollect past activities related to farming and GHG emissions. Therefore, the results might be affected by respondents' memory errors. Ultimately, although the GHG emission mitigation techniques farmers utilized varied from one responder to the next, they were all quantified and used to arrive at the study's conclusive findings. Therefore, the results might be inconsistent in different regions where the same situation is present. Further studies should look at the cost-benefit of utilization of each of the rice farmer's GHG Emission mitigation options. It is possible that farmers are utilising these GHG Emission strategies because of the economic gain they receive for using it than the aim of reducing GHG emissions in rice fields.

### **Conclusion and Recommendations**

GHG emissions from rice fields contribute significantly to global warming and climate change. Therefore, mitigating emissions from rice fields is very critical to limit global warming, increasing yield, and income, ensuring food security and improving the standard of living for farmers and all. Understanding farmers' knowledge of GHG emissions and mitigation strategies is among the crucial starting points for deciding what steps should be made to mitigate emissions from rice fields and produce rice in a cleaner environment. Some of the GHG emission mitigation strategies in rice fields identified in the study include alternate wetting and drying (AWD); system of rice intensification (SRI); changing tillage operations (CTO); Nitrogen Fertilizer Management (NFM); residue management (RM) and aerobic rice varieties (ARV). The mean age of the farmers was 45 years. Most were male with an average farm size of 1.53 hectares. The land is one of the essential productive inputs in rice farming and when farmers have a limited size of it, increased farm production and mitigating GHG emissions may be unattainable. Most of the rice farmers were visited at least twice per year by the extension agents in the area. This is quite poor and could negatively affect rice farmers' mitigation efforts to GHG emission in the area. Extension contact improves farmers' access to recent information, expertise and knowledge on contemporary farming techniques to raise their yield, income and standard of living. The low knowledge of farmers on the use of AWD and SRI GHG emission mitigation strategies could be associated with poor extension contact earlier found in the study. AWD is a promising technique that involves intermittently flooding and drying rice fields. When compared to continuous flooding, it has the potential to cut CH<sub>4</sub> emissions by an average of 48%. Extension

agents are charged with transmitting recent innovations to farmers in their farmland to help achieve farmer's production objectives. However, when the extension service system is not well strengthened, it creates a gap between extension agents and farmers. Rice farmers are therefore encouraged to constantly seek information on GHG emission mitigation strategies before embarking on rice production, this will without doubt position them to overcome any adverse effect of global warming. Additionally, the government should assist farmers in implementing climate change and GHG policies to reduce the challenges they face and increase rice production in a clean environment. The study recommends that the government should strengthen and support agricultural extension service systems so as to enable extension personnel to visit farmer's farms regularly to transmit recent innovations related to GHG emission mitigation strategies as these would increase yield, income and standard of living of the farmers. Ultimately, since rice farmers are members of cooperatives, they are encouraged to take advantage of their strength and shared a common purpose to collectively project a common demand in obtaining funds and other necessary inputs for the implementation and scaling-up of GHG emission mitigation strategies in the area.

### **Significance Statement:**

Rice is critical to Nigeria from separate vantage points: first, in terms of the quantity of calories it provides per person and per day; and second, based on the value of income it generates through its various local production value chains. However, rice production in Nigeria is vulnerable to climate change and associated with greenhouse gas (GHG) emissions, mainly anthropogenic methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O). Understanding farmers' knowledge on GHG emission mitigation strategies would be crucial in reducing emission from rice fields, producing rice in a cleaner environment. An essential first step in evaluating what steps should be implemented to combat the detrimental impacts of climate change on rice output and the environment is assessing the degree of GHG emission knowledge among rice farmers. Most rice farmers in the region have limited knowledge on AWD. AWD is one of the major GHG mitigation strategies developed by IRRI in 2002, has been adequately studied and propagated globally by various scholars and institutions. AWD is a promising technique that involves intermittently flooding and drying rice fields. When compared to continuous flooding, it has the potential to cut CH<sub>4</sub> emissions by an average of 48%. The study recommends that the government should strengthen and support agricultural extension service systems so as to enable extension personnel to visit farmer's farms regularly to transmit recent innovations related to GHG emission mitigation strategies as these would increase yield, income and standard of living of the farmers. The study also found that rice farmers complained of limited capital. Therefore, as members of cooperatives, farmers are encouraged to take advantage of their strength and shared a common purpose to collectively project a common demand in obtaining funds and other necessary inputs for the implementation and scaling-up of GHG emission mitigation strategies in the area. Finally, further studies should look at the cost-benefit of utilising each of the rice farmer's GHG Emission Mitigation options. It is possible that rice farmers are utilizing these GHG Emission strategies because of the economic gain they receive for using them rather than the aim of reducing GHG emissions in rice fields

## REFERENCES:

1. Food and Agricultural Organization (FAO) (2023). *Crop Prospects and Food Situation* – Quarterly Global Report No. 1, March 2023. Rome. <https://doi.org/10.4060/cc4665en>
2. Li, D., He, H., Zhou, G., He, Q., & Yang, S. (2023). Rice Yield and Greenhouse Gas Emissions Due to Biochar and Straw Application under Optimal Reduced N Fertilizers in a Double Season Rice Cropping System. *Agronomy*. 2023; 13(4), 1-13. <https://doi.org/10.3390/agronomy13041023>
3. Esiobu, N. S. (2020). Does the Incidence of COVID-19 Pandemic Affect Rice Yield? Lessons from Southeast Nigeria. *Journal of Biology, Agriculture and Healthcare*, 10(15), 38-56
4. Ojo, T.O., Ogundeji, A.A., Babu, S.C., & Alimi, T. (2020). Estimating financing gaps in rice production in Southwestern Nigeria. *Economic Structures*9(12), <https://doi.org/10.1186/s40008-020-0190-y>
5. Libby George (2020). A growing problem: Nigerian rice farmers fall short after borders close, accessed from <https://www.reuters.com/article/ozatp-uk-nigeria-economy-rice-idAFKBN1ZM10B-OZATP> on 12-03-2023
6. Federal Ministry of Agriculture and Rural Development (FMARD), (2022). National Agriculture Technology and Innovation Policy (NATIP), retrieved on 16<sup>nd</sup> February, 2022 from <https://fmard.gov.ng/wp-content/uploads/2022/07/National-Agricultural-Technology-and-Innovation-Policy-2022-2027-e-copy.pdf>
7. Gupta, K., Kumar, R., & Baruah, K.K (2021). Greenhouse gas emission from rice fields: a review from Indian context. *Environmental Science and Pollution Research*28(8), 30551–30572, <https://doi.org/10.1007/s11356-021-13935-1>
8. Babatunde, P.E. (2023). Rice Production, Imports And Economic Growth In Nigeria: An Application Of Autoregressive Distributed Lag,, *International Journal of Advanced Economics*, 5(2),48-56, DOI: 10.51594/ijae.v5i2.449
9. Murugaiyan, V., Ali, J., Frei, M., Zeibig, F., Pandey, A., Wairich, A., Wu, L-B., Murugaiyan, J., & Li, Z. (2021). Identification of Promising Genotypes Through Systematic Evaluation for Arsenic Tolerance and Exclusion in Rice (*Oryza sativa* L.). *Frontiers in Plant Science*, 12(753063).1-16, doi: 10.3389/fpls.2021.753063
10. Chandrasekaran, D., Tabassum-Abbasi, T., & Abbasi, S.A. (2022). Assessment of Methane Emission and the Factors That Influence It, from Three Rice Varieties

Commonly Cultivated in the State of Puducherry. *Atmosphere*, 13(11):1-12, <https://doi.org/10.3390/atmos13111811>

11. Karki, S., Adviento-Borbe, M.A., Massey, J.H., & Reba, M.L. (2021). Assessing Seasonal Methane and Nitrous Oxide Emissions from Furrow-Irrigated Rice with Cover Crops. *Agriculture*, 11,(3),1-15 <https://doi.org/10.3390/agriculture11030261>
12. Esiobu, N.S, Onubuogu, G.C., Njoku, M.S., &Nwachukwu, N.B. (2021). Sustainability and Determinate of Farmers' Mitigation Strategies to Greenhouse Gases Emission: A Case in Rice Agric-Food System of Nigeria. *IntechOpen*. doi: 10.5772/intechopen.93188
13. Jiang, M., Li, X., Xin, L., Tan, M.,&Zhang, W. (2023). Impacts of Rice Cropping System Changes on Paddy Methane Emissions in Southern China. *Land* 2023, 12, 270. <https://doi.org/10.3390/land12020270>
14. Ali, J., Anumalla, M., Murugaiyan, V., & Li, Z. (2021). Green Super Rice (GSR) Traits: Breeding and Genetics for Multiple Biotic and Abiotic Stress Tolerance in Rice. In: Ali, J., Wani, S.H. (eds) *Rice Improvement*. Springer, Cham. [https://doi.org/10.1007/978-3-030-66530-2\\_3](https://doi.org/10.1007/978-3-030-66530-2_3)
15. Nigeria Population Commission (NPC) (2006). Nigeria Population Commission, *Nigeria Federal Government Initiative of individual head count by gender. Spread, State by State*, In:MOFINEWS; (accessed 28 February, 2016).Jan-Feb,2007,<https://nigeria.opendataforafrica.org/ifpbxbd/state-population-2006>.
16. Nigerian Meteorological Agency (NiMET) (2012). Drought and Flood Monitoring in South East Bulletin. Retrieved from [https://www.un-spider.org/sites/default/files/Session2-WG2-JamesAdamu-NIMET\\_0.pdf](https://www.un-spider.org/sites/default/files/Session2-WG2-JamesAdamu-NIMET_0.pdf)25 October, 2018
17. National Bureau of Statistics (NBS) (2007). National Bureau of Statistics Official Gazette (FGP 71/52007/2,500(OL24): National Population Estimates *Legal Notice on Publication of the Details of the Breakdown of the National and State Provisional Totals, Census*. file:///C:/Users/USER/Downloads/Population%20Forecasts.pdf(accessed 28 February, 2016)
18. Nigerian Geological Survey Agency, (NGSA) (2020). Nigeria Geological Maps Retrieved 28th March, 2021, from <https://www.scribd.com/document/81855420/Geological-Map-of-Nigeria#>
19. Olugbenga, O.A., Godbless, F.S., & Jeremiah, S.A. (2023). Cost efficiency and profitability analysis of rice (*Oryza sativa*) production among smallholder farmers in Federal Capital Territory, Nigeria,*Australian Journal of Science and Technology*, 7(1);1-10

20. Abdulwahab, A., Lateef, L. A., Obafemi, B. O., Ayanda, V. O., Adeniyi, K. I., & Raza, H. A. (2023). Traditional and Modern Perspectives in Agricultural Extension Practice in Nigeria. *Journal of Social Sciences Review*, 3(1), 552-561. <https://doi.org/10.54183/jssr.v3i1.196>
21. Tobi, K., & Edgeweblimel, K. (2023). Land Tenure Security and Climate Change Adaptation Strategies in Togo, 3(1), 1-11, doi: <https://assets.researchsquare.com/files/rs-2587883/v1/25df51ac-0150-4ee9-8000-aa532aa70243.pdf?c=1676501566>
22. Esiobu, N. S., Theresa, O. U., Akande, S. N., Udunwa, N. B., Emeruwa, A. M. (2023). Do Farmers Derive Returns From Cassava Production? Lessons from Imo State, Nigeria. *Adv Dairy Sci Res*, 1(1), 11-22, <https://www.opastpublishers.com/open-access-articles/do-farmers-derive-returns-from-cassava-production-lessons-from-imo-state-nigeria.pdf>
23. Sander, B. O., Wassmann, R., Palao, L. K., & Nelson, A. (2017). Climate-based suitability assessment for alternate wetting and drying water management in the Philippines: a novel approach for mapping methane mitigation potential in rice production. *Carbon Manage.* 8, 331–342. doi: 10.1080/17583004.2017.1362945
- 24.** Anapalli, S. S., Pinnamaneni, S. R., Reddy, K. N., Wagle, P., & Ashworth, A. J. (2023). Eddy covariance assessment of alternate wetting and drying floodwater management on rice methane emissions. *Heliyon*, 9(4), e14696. <https://doi.org/10.1016/j.heliyon.2023.e14696>
25. Chirinda, N., Arenas, L., Katto, M., Loaiza, S., Correa, F., Isthitani, M., Loboguerrero, A.M., Martinez-Baron, D., Graterol, E., Jaramillo, S., Torres, C.F., Arango, M., Guzman, M., Avila, I., Hube, S., Kurtz, D.B., Zorrilla, G., Terra, J., Irisarri, P., Tarlera, S., LaHue, G., Scivittaro, W.B., Noguera, A., & Bayer, C. (2018). Sustainable and Low Greenhouse Gas Emitting Rice Production in Latin America and the Caribbean: A Review on the Transition from Ideality to Reality. *Sustainability*, 10(3):671-683, doi: <https://doi.org/10.3390/su10030671>
26. Lai L.R., & Sander, B.O. (2021). Policy Framework for Myanmar Rice Production and In-Depth Study on Greenhouse Gas Emissions. *Journal of Earth and Environmental Science Research*. SRC/JEESR-185. DOI: [doi.org/10.47363](https://doi.org/10.47363), 3(6), 12-13
27. Sander, O.B., Schneider, P., Romasanta R., Samoy-Pascual, K., Evangelina, B. S., Constancio A. A., & Reiner W. (2020). Potential of Alternate Wetting and Drying Irrigation Practices for the Mitigation of GHG Emissions from Rice Fields: Two Cases in Central Luzon (Philippines), *Agriculture* 2020, 10, 0350; doi:10.3390/agriculture10080350

28. Islam, S. M. M., Gaihre, Y. K., Islam, M. R., Ahmed, M. N., Akter, M., Singh, U., & Sander, B. O. (2022). Mitigating greenhouse gas emissions from irrigated rice cultivation through improved fertilizer and water management. *Journal of environmental management*, 307, 114520. <https://doi.org/10.1016/j.jenvman.2022.114520>
29. Kaur, M., Malik, D.P., & Malhi, G.S. (2022). Rice residue management in the Indo-Gangetic Plains for climate and food security. A review. *Agronomy for Sustainable Development*, 42(92). 1-12. <https://doi.org/10.1007/s13593-022-00817-0>
30. Reddy, M. P., Chaudhary, S. K., Kurdekar, A. K., Sandeep, S. N., & Kumar, M. V. (2022). Effect of Brown Manuring and Different Levels of Nutrients on Growth and Yield Attributes of Aerobic Rice (*Oryza sativa* L.). *International Journal of Environment and Climate Change*, 12(12), 171–177. <https://doi.org/10.9734/ijecc/2022/v12i121450>
31. Menegat, S., Ledo, A. & Tirado, R. (2022). Greenhouse gas emissions from global production and use of nitrogen synthetic fertilisers in agriculture. *Scientific reports* 12(3), 14490, <https://doi.org/10.1038/s41598-022-18773-w>
32. Gao, Y., & Serrenho, A. (2023). Greenhouse gas emissions from nitrogen fertilizers could be reduced by up to one-fifth of current levels by 2050 with combined interventions. *Nature Food* 4, 170–178 (2023). <https://doi.org/10.1038/s43016-023-00698-w>
33. Hassan, M.U., Aamer, M., Mahmood, A., Awan, M.I., Barbanti, L., Seleiman, M.F., Bakhsh, G., Alkharabsheh, H.M., Babur, E., Shao, J., Rasheed, A., & Huang, G. (2022). Management Strategies to Mitigate N<sub>2</sub>O Emissions in Agriculture. *Life*, 12(3):439. <https://doi.org/10.3390/life12030439>
34. Nirmala, B., Tuti, M.D., Mahender R.K., Amtul, W., Muthuraman, P., Brajendra, P., & Vidhan, S. (2021). Integrated assessment of system of rice intensification vs. conventional method of transplanting for economic benefit, energy efficiency and lower global warming potential in India, Agroecology and Sustainable Food Systems, <https://doi.org/10.1080/21683565.2020.1868648>
35. International Rice Research Institute (IRRI) (2022). Saving Water with Alternate Wetting Drying (AWD), access from <http://www.knowledgebank.irri.org/training/factsheets/water-management/saving-water-alternate-wetting-drying-awd> on 22-03-2023
36. Allen, J. M., & Sander, B. O. (2019). The Diverse Benefits of Alternate Wetting and Drying (AWD). Available online at: <https://hdl.handle.net/10568/101399> (accessed February 18, 2021).
37. Enriquez, Y., Yadav, S., Evangelista, G.K., Villanueva, D., Burac, M.A., & Pede, V. (2021). Disentangling Challenges to Scaling Alternate Wetting and Drying Technology for Rice Cultivation: Distilling Lessons From 20 Years of Experience in the Philippines.

Front. Sustain. Food Syst. 5:675818. doi: 10.3389/fsufs.2021.675818 (accessed April 5, 2022).

38. Ali, J., Xu, J.L., Gao, Y.M., Ma, X.F., Meng, L.J., Wang, Y., Pang, Y.L., Guan, Y.S., Xu, M.R., &Revilleza, J.E. (2017) Harnessing the hidden genetic diversity for improving multiple abiotic stress tolerance in rice (*Oryza sativa* L.). PLoS ONE 12:e017251

UNDER PEER REVIEW