

Original Research Article

Effect of Sulphur and FYM application on performance of Urdbean [*Vigna Mungo* (L.) Hepper] and Soil properties.

ABSTRACT

Formatted: Font: 14 pt

A field experiment was conducted at the Crop Research Centre of Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut (U.P.), India, during the *kharif* season of 2020 to assess the effect of farmyard manure (FYM) and sulfur on the performance of urdbean [*Vigna mungo* (L.)] and soil properties. The experiment was designed as a Randomized Block Design (RBD) with seven treatments, consisting of different doses of FYM and sulfur, replicated three times. The test variety used was Shekhar-2. The results revealed that the combined application of FYM and sulphur exhibited significant influence on the growth, yield attributes and yields of urdbean as compared to control during the year of experimentation. Significant improvement in viz. Plant height (cm), Number of Trifoliolate Leaves, Dry matter accumulation, Test weight (g), Grain yield (q ha^{-1}), Number of Root nodule, Protein content (%), Protein yield (kg ha^{-1}) and soil properties was noticed with the application of RDF + FYM @ 2.5 t ha^{-1} + S @ 10.0 kg ha^{-1} , which established its superiority over rest of the FYM and sulphur treatments. Increase of 91.61% in grains was recorded with the application of RDF + FYM @ 2.5 t ha^{-1} + S @ 10.0 kg ha^{-1} over control. The application of RDF + FYM @ 2.5 t ha^{-1} + S @ 10.0 kg ha^{-1} resulted in higher protein yield (274.58 kg ha^{-1}). Based on these findings, it can be concluded that the application of RDF + FYM @ 2.5 t ha^{-1} + S @ 10.0 kg ha^{-1} is recommended for sustainable urdbean production.

Keywords – Sulphur, FYM, Growth parameters, Yield attributes and Urdbean.

Formatted: Font: Italic

Introduction

Formatted: Font: 14 pt

Pulses are the major constituent of Indian diet for providing protein rich food to vegetarian masses. Apart from this, pulses provide high quality feed and fodder to livestock and improve the soil health. The pulses are important source of dietary protein and have unique property of maintaining and restoring soil fertility through biological nitrogen fixation as well as conserving and improving physical properties of soil by virtue of their deep root system and leaf fall.

Urdbean [*Vigna mungo* (L.) Hepper] is one of the most important pulse crops of rainfed areas grown throughout the country. This crop is grown in different cropping system as a mixed crop, catch crop, sequential crop in the country. Urdbean seed contains 25-26% proteins, 60% carbohydrates, 1.5% fat, and minerals combination, amino acid, and essential vitamins *etc.* In India urdbean is very popularly grown in Andhra Pradesh, Bihar, Madhya Pradesh, Maharashtra, U.P., West Bengal, Punjab, Haryana, and Karnataka. It is used as nutritive fodder specially for milch cattle. It is also used as a green manuring crop. Black gram (urdbean) is native of India and originated from *Phaseolus sublobatus*, a wild plant. In India total Urdbean production was estimated 1.90 million tonnes from 3.11 million hectare area with a productivity of 642 kg/ha (Kumar *et al.*, 2019).

Formatted: Font: Italic

Sulphur is considered to be the fourth major plant nutrient and hence important as nitrogen, phosphorus and potassium. With the use of high analysis fertilizer and ~~continuous~~continuous depletion of ~~sulphur~~Sulphur from soil by crop removal and leaching, S is becoming a limiting factor in crop growth. In India, sulphur responses have been obtained from pastures, pulses and oilseed. In recent years, the deficiency of ~~sulphur~~Sulphur has increased in the country mainly due to declining organic matter in soil which is a source of ~~sulphur~~Sulphur. Sulphur ~~stimulate~~stimulates cell division, photosynthetic process as well as formation of chlorophyll and also promotes the root nodules in legumes. It is involved in protein synthesis and is considered indispensable for the synthesis of cysteine, cystine and methionine. It resulted in higher plant height and number of branches per plant and ultimately helped in realization of higher grain and straw yield (Singh & Yadav, 2004, Mitra et al., 2006 and Kadam et al., 2006). With the immense benefits of ~~sulphur~~Sulphur on pulse production and lower level of S in Indian soil it is important to supplement S in plant nutrition. The application of ~~sulphur~~Sulphur in integrated approach may be beneficial therefore present investigation was carried out.

Materials and method

Formatted: Font: 14 pt

The field experiment was conducted at the Crop Research Centre (CRC) of Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut (UP) during *kharif* season of 2020. The experimental soil was sandy loam in texture, alkaline in reaction having low in nitrogen, ~~sulphur~~Sulphur medium in phosphorous and potassium. The experiment was laid out in a randomized block design with seven treatments and three replication: T₁ (Control), T₂ (RDF), T₃ (RDF + FYM @ 2.5 t ha⁻¹), T₄ (RDF + S @ 10.0 kg ha⁻¹), T₅ (RDF + FYM @ 5.0 t ha⁻¹), T₆ (RDF + S @ 20 kg ha⁻¹), T₇ (RDF + FYM @ 2.5 t ha⁻¹ + S @ 10 kg ha⁻¹). Recommended dose of NPK (20:60:00) in the experiment was applied through urea, DAP, FYM and elemental S was applied as per treatment. For preparing experimental field one deep ploughing and two harrowing were done. Treated ~~urd~~Urd seed @ 15 kg/ha was sown in plant geometry 30 cm × 10 cm row to row and plant to plant. Soil was analysed for soil pH (1:2) soil EC (dSm⁻¹) at 25°C of 1:2 for soil water solution, organic carbon by wet oxidation method (Walkley and Black method, 1934); available nitrogen by alkaline potassium permanganate method (Subbiah and Asija, 1956); available phosphorus by colorimetric method (Olsen et al., 1954); available potassium by 1N NH₄OAC extraction (Hanway and Heidal, 1952) and available sulphur by turbidimetric method (Chesnin and Yien, 1950).

Result and Discussion

Formatted: Font: 14 pt

Effect on FYM and Sulphur on growth and yield of urdbean

Formatted: Font: 16 pt

Formatted: Font: 14 pt, Not Italic

The effect of different treatments on plant height (Table 1) was assessed, and it was observed that the application of recommended dose of fertilizer (RDF) did not result in a significant difference compared to the control group. However, when sulfur (S) or farmyard manure (FYM) was supplemented along with RDF, a significant improvement in plant height was observed. Among the two supplements, the application of S along with RDF showed a better effect compared to FYM. The application of either level of S or FYM over RDF did not have a significant impact. However, when both S (10 kg/ha) and FYM (2.5 ton/ha) were applied together over RDF, a significant improvement in plant height was noticed. The better effect of S compared to FYM could be due to

Formatted: Font: Bold

the specific nutrient composition or mode of action of sulfur in promoting plant height. Additionally, the conjoint application of S and FYM at specific levels may have synergistic effects, resulting in a significant improvement in plant height. (Dubey *et al.*, (2020))

Formatted: Font: Not Bold

Number of Trifoliolate Leaves (Table 1) at harvest stage, the maximum number of trifoliolate leaf plant⁻¹ was noticed in T₇ (RDF + FYM @ 2.5 t ha⁻¹ + S @ 10.0 kg ha⁻¹) treatment which remained statistically at par T₆ (RDF + S @ 20.0 kg ha⁻¹) and T₅ (RDF + FYM @ 5.0 t ha⁻¹) but significantly higher than rest of the treatments. Significantly lower number of trifoliolate leaf plant⁻¹ (5.95 and 4.90 plant⁻¹) was recorded with control treatment. The presence of FYM and S may have provided essential nutrients, promoting leaf development and resulting in a higher number of trifoliolate leaves. In contrast, the lower number of trifoliolate leaves in the control treatment could be due to nutrient deficiencies or suboptimal conditions for leaf growth. (Barakzai *et al.*, 2020).

Formatted: Font: Not Bold

Dry matter accumulation (Table 1) per plant differ significantly under different treatments. Fertilization over control resulted a significant effect on dry matter accumulation. Application of lower level of S and FYM over RDF could not brought any remarkable variation but a significant variation in dry matter accumulation was noticed with the application of higher level of S and FYM over RDF. The lower level of S and FYM may not have provided a substantial additional nutrient supply, resulting in no remarkable variation in dry matter accumulation compared to RDF alone. However, the higher levels of S and FYM likely supplied an optimal nutrient balance, leading to a significant increase in dry matter accumulation per plant. (Singh *et al.*, 2016)

Formatted: Font: Bold

Test weight (Table 1) remained unaffected by the application of RDF compared to the control group. However, the conjoint application of S and FYM over RDF resulted in a significant improvement in test weight. the significant improvement in test weight with the conjoint application of sulfur (S) and farmyard manure (FYM) over RDF suggests that these supplements may have provided additional nutrients or beneficial effects that positively impacted the test weight parameter. (Dubey *et al.*, 2020).

Formatted: Font: Bold

Formatted: Font: Bold

Grain yield (Table 1) which differ significantly under different treatment ranged from 5.98 to 11.46 q ha⁻¹. Application of RDF over control resulted 39 percent increase in grain yield while application of S @ 10kg ha⁻¹, FYM @ 2.5 ton ha⁻¹, S @ 20kg ha⁻¹, FYM @ 5.00 ton ha⁻¹ over RDF resulted a significant increase of 14.4, 17.1, 27.3 and 31.9 percent respectively. Conjoint application of sulphur and FYM over RDF improved the grain yield by 37.9 percent. Number of trifoliolate leaves increased significantly with the application of higher level of S or FYM over RDF. Among the S and FYM, application of S exerted much effect on trifoliolate leaves than FYM. The application of S sulfur (S) and farmyard manure (FYM) at specific levels over RDF led to further improvements in grain yield, with S showing a greater effect on grain yield and the number of trifoliolate leaves compared to FYM. The conjoint application of S and FYM over RDF further enhanced grain yield, likely due to the combined effect of improved nutrient availability and plant growth stimulation. (Ahamad *et al.*, 2016).

Formatted: Font: Not Bold

Formatted: Font: Bold

Formatted: Font: Not Bold

Root nodules (Table 1) which are sites of nitrogen fixation, also showed significant improvement with the application of higher levels of S and FYM over RDF. Both S and FYM levels differed significantly in terms of the number of root nodules per plant, with higher levels resulting in significantly more root nodules than lower levels. Higher levels of S and FYM

Formatted: Font: Bold

likely provided an enriched nutrient environment, promoting the development of more root nodules per plant (Barakzai *et al.*, 2020).

Formatted: Font: Not Bold

Table 1: Effect of FYM and Sulphur on Plant height, number of trifoliolate leaves, Dry matter accumulation, Test weight, Grain Yield, Number of root nodules per plant

Formatted: Font: Bold

Treatment	Plant height (cm)	Number of Trifoliolate Leaves	Dry matter accumulation	Test weight (g)	Grain yield (q ha ⁻¹)	Number of Root nodule Plant ⁻¹
T ₁	44.54	4.90	9.09	28.34	5.98	21.3
T ₂	49.15	6.43	12.02	31.28	8.31	28.0
RDF						
T ₃	50.88	6.82	11.77	31.67	9.51	29.7
RDF + FYM @ 2.5 t ha ⁻¹						
T ₄	52.13	7.01	12.35	32.18	9.73	30.8
RDF + S @ 10.0 kg ha ⁻¹						
T ₅	53.18	7.78	13.08	33.81	10.58	33.3
RDF + FYM @ 5.0 t ha ⁻¹						
T ₆	53.95	7.87	13.10	33.99	10.96	34.2
RDF + S @ 20.0 kg ha ⁻¹						
T ₇	55.10	8.06	13.88	34.34	11.46	36.3
RDF + FYM @ 2.5 t ha ⁻¹ + S @ 10.0 kg ha ⁻¹						
SEM	1.6	0.23	0.38	1.04	0.30	1.02
CD (P=0.05)	5.1	0.68	1.05	3.04	0.88	3.04

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Effect of FYM and S on pH, EC and organic carbon content of soil

Formatted: Font: Not Italic

The results reveal that (Table 2) the application of FYM and S sulphur significantly affected organic carbon, and non-significantly soil pH and EC. Highest organic carbon and EC (0.52% and 0.34 dSm⁻¹) were found in T₇ (RDF + FYM @ 2.5 t ha⁻¹ + S @ 10.0 kg ha⁻¹) while,

Formatted: Font: Bold

lowest (0.43% and 0.26 dSm⁻¹) ~~under~~ under control (T1). These results are obvious because more availability of nutrients, will favor vigorous growth of roots and plants and will add organic matter to the soil. After the decomposition of organic matter produced organic acids will dissolve the insoluble mineral in the systems and consequently increases the EC. FYM and S sulphur have an acidifying effect on soil pH. The results obtained in the present investigation are inconformity with the findings of Basumatary (2018) and Kantva *et al.* (2019).

Table 2. Effect of FYM and S On pH, EC and organic carbon content of soil

	Treatments	Soil pH	EC (dS m ⁻¹)	Organic Carbon (%)
T ₁	Control	7.80	0.26	0.43
T ₂	RDF	7.77	0.29	0.46
T ₃	RDF + FYM @ 2.5 t ha ⁻¹	7.75	0.29	0.47
T ₄	RDF + S @ 10.0 kg ha ⁻¹	7.72	0.30	0.48
T ₅	RDF + FYM @ 5.0 t ha ⁻¹	7.69	0.31	0.50
T ₆	RDF + S @ 20.0 kg ha ⁻¹	7.63	0.33	0.51
T ₇	RDF + FYM @ 2.5 t ha ⁻¹ + S @ 10.0 kg ha ⁻¹	7.58	0.34	0.52
	SE m (±)	0.2	0.03	0.01
	C.D. (P=0.05)	NS	NS	0.03

Formatted: Font: Not Bold, Not Italic

Formatted: Font: Not Bold

Formatted: Font: Not Bold, Not Italic

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Effect of FYM and S on Available N, P, K and S content of soil

Formatted: Font: Not Italic

Results showed that the availability of nutrients in soil was significantly affected by different treatments of FYM and S sulphur (Table 3). At harvesting stage available nitrogen in soil ranged from 178.4 to 200.2 kg ha⁻¹. Higher available N found in T₇ may be attributed to application of FYM and S sulphur. At harvest available N decline in control and significantly higher N over control was found due to conjoint application of FYM and S sulphur or their higher level along with recommended NPK. Similar results were also found in available phosphorus. At harvest stage available P in soil ranged from 14.1 to 22.2 kg ha⁻¹. At harvest available P decline in control and significant increase over control was found in rest of the treatments. Availability of K increases significantly due to conjoint application of S and FYM over control however rest treatments remain statistically at par. Availability of S increased non

Formatted: Font: Bold

significantly due to fertilization over control. Availability of phosphorus increased significantly due to conjoint application of S and FYM over RDF or higher level of S and FYM. FYM and sulphur produces organic acids, which convert tri-calcium phosphate to di and mono calcium phosphate with the net result of the enhanced availability of phosphorus. Adsorption of PO_4^{-3} with higher level availability resulted in a concurrent desorption of SO_4^{-2} anions from colloidal surface therefore, optimum doses of P fertilizer may result in increased mobility and availability of sulphur-Sulphur in soil. The increment in available nutrient status might be due to favorable conditions for microbial activity as well as chemical activity. These results obtained in the present investigation are in close conformity with the findings of Rambharose *et al.* (2014), Kokani *et al.* (2015), Nagar *et al.* (2016) and Karnavat *et al.* (2018).

Table 3: Effect of FYM and Sulphur on Available nutrient

Treatments		Available nutrients (kg ha ⁻¹)			
		N	P	K	S
T ₁	Control	178.4	14.1	176.5	8.1
T ₂	RDF	190.7	18.6	182.5	8.3
T ₃	RDF + FYM @ 2.5 t ha ⁻¹	191.6	20.5	184.2	8.5
T ₄	RDF + S @ 10.0 kg ha ⁻¹	192.7	19.4	185.5	8.6
T ₅	RDF + FYM @ 5.0 t ha ⁻¹	196.5	21.2	191.3	9.1
T ₆	RDF + S @ 20.0 kg ha ⁻¹	198.6	21.3	192.6	9.3
T ₇	RDF + FYM @ 2.5 t ha ⁻¹ + S @ 10.0 kg ha ⁻¹	200.2	22.2	196.6	9.5
SEm±		5.7	0.7	6.3	0.34
CD (P= 0.05)		16.9	2.1	18.7	NS

Formatted: Font: Not Bold, Not Italic

Formatted: Font: Not Bold

Formatted: Font: Not Bold, Not Italic

Formatted: Font: Not Bold

Formatted: Font: Not Italic

Formatted: Font: Not Bold

Formatted: Font: Not Bold, Not Italic

Formatted: Font: Not Bold

Formatted: Font: Not Italic

Formatted: Font: Not Bold, Not Italic

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Effect of FYM and S on protein content and yield of urdbean

Formatted: Font: 14 pt, Not Italic

The protein content (Table 4) of urdbean increases non significantly due to different FYM and sulphur-Sulphur treatments. Highest protein yield 274.58 kg ha⁻¹ significantly higher than the other treatments with exception of T₆ was found in T₇. Application of S and FYM over RDF improved protein yield significantly. Protein yield increased significantly with the application of higher level of S and FYM than their lower level.

Formatted: Font: Bold

Table: 4 Effect of FYM and Sulphur on Protein content, and protein yield

Treatment		Protein content (%)	Protein yield (kg ha ⁻¹)
T ₁	Control	21.09	126.12
T ₂	RDF	21.90	181.99
T ₃	RDF + FYM @ 2.5 t ha ⁻¹	22.15	210.65
T ₄	RDF + S @ 10.0 kg ha ⁻¹	22.40	217.95
T ₅	RDF + FYM @ 5.0 t ha ⁻¹	23.09	244.06
T ₆	RDF + S @ 20.0 kg ha ⁻¹	23.52	257.78
T ₇	RDF + FYM @ 2.5 t ha ⁻¹ + S @ 10.0 kg ha ⁻¹	23.96	274.58
SEm ±		0.72	7.40
CD (P= 0.05)		NS	21.99

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: 14 pt

Conclusion

In the light of the results summarized as above, it may be concluded that for obtaining higher productivity of urdbean and sustainable soil health the treated seed should be sown with FYM and sulphur and recommended fertilizers. Since the observation is based on one year experimental data, so it is advisable that the experiment should be repeated for few more years to confirm the findings.

References

Rambharose, T Thomas, Sudhir Kumar Singh, Jagdish Prasad Bairwa, Singh Dharmveer (2014) Effect of Sulphur and FYM on physical and chemical properties of sunflower grown soil and nutrient uptake by crop, *Bulletin of environmental and scientific research*, 3(1):1-7.

Ahamad, A. T. I. K., Kumar, N. E. E. R. A. J., & Yadav, D. E. V. I. D. E. E. N. (2017). Integrated nutrient management in pigeonpea-based intercropping systems. *Indian Journal of Agronomy*, 62(4), 125-30.

Barakzai, K. R., Dhar, S., Khalili, A., Rasooli, M., & Tanha, J. (2020). Effect of sources of nutrient and biofertilizers on growth and yield of mungbean (*Vigna radiata* L.). *IJCS*, 8(1), 555-559.

Formatted: Font: 14 pt

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Bold

Formatted: Font: Not Bold

Comment [L1]:

Formatted: Font: Bold

Formatted: Font: Bold

Basumatary, A (2018). Integrated Sulphur Management in Rapeseed (*Brassica Campestris*). Black gram (*Vigna mungo*) Sequence in an Inceptisol of Assam, *Journal of the Indian Society of Soil Science*, 66(4): 425-431

Dubey, D., Rana, N. S., Chaudhari, A., & Dubey, V. (2020). Growth and yield behaviour of urdbean (*Vigna mungo* L.) as influenced by sowing methods and nutrient management practices in north-western plains zone.

Hanway, J. J. and Heidal, H. (1952). Soil analysis methods as used in Iowa State College, Soil Testing Laboratory. *Iowa Agriculture*, 27: 1-13.

Kadam, S.S., Patil, A.V., Mahadkar, U.V. and Gaikwad, C.B. (2006). Effect of potassium and sulphur on the growth and yield of urdbean. *Journal of Maharashtra Agriculture University* 31(3): 382-383.

Karnavat, R., Pavaya, R.P., Malav, J.K., Chaudhary, N., Patel, I.M. and Patel, J.K. (2018). Effect of FYM, phosphorus and PSB on yield, nutrient content and uptake by urdbean (*Vigna mungo* (L.) Wilczek) on loamy sand. *International Journal of Chemical Studies* 6(2): 1026-1029.

Kokani, J. M, K A Shah, BM Tandel, GJ Bhimani (2015). Effect of FYM, phosphorus and sulphur on yield of summer black gram and post-harvest nutrient status of soil. *The Bioscan*, 10(1): 379-383

Kumar, Deepak., Singh, R.P., and Simaiya, Vidya. (2019). Effect of foliar application of nutrients on yield and economics of blackgram (*Vigna mungo* {L.} Hepper) under rainfed Vertisols of Central India. *Journal of Pharmacognosy and Phytochemistry* 8(1): 2373-2376.

Mitra, A.K., Banerjee, K. and Pal, A. K. (2006). Effect of different levels of phosphorus and sulphur on yield attributes, seed yield, protein content of seed and economics of summer urdbean. *Research on Crops*. 7(2):404-405.

Nagar, R. K, V V Goud, Rajesh Kumar and Ravindra Kumar (2016). Effect of organic manures and crop residue management on physical, chemical and biological properties of soil under pigeonpea based intercropping system, *International Journal of Farm Sciences*, 6(1): 101-113.

Olsen, S.R., Cole, R V., Watanabe, F.S. and Lean, L.A (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate. *United States Department of Agriculture Circular*, 19:939.

Singh, O., Kumar, S., Shahi, U. P., Kumar, A., Dwivedi, A., & Kumar, V. (2016). Growth, nodulation, yield and nutrient uptake of urdbean (*Vigna mungo* L. Hepper) as influenced by sulphur and iron fertilization in light textured soil. *International Journal of Bio-Resource & Stress Management*, 7.

Singh, U. and Yadav, D.S 2004. Response of green gram to sulphur and zinc. *Annals of Agriculture* 25(3): 463-464.

Formatted: Font: Not Bold

Formatted: Font: Bold

Comment [L2]: What page no.

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Bold

Formatted: Font: Not Bold

Formatted: Font: Bold

Formatted: Font: Not Bold

Formatted: Font: Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Bold

Formatted: Font: Not Bold

Formatted: Font: Bold, Italic

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Bold

| Subbaiah, B.V. and Asija, G. L. (1956). A rapid procedure for the estimation of available nitrogen in soils. *Current Research Science* 25: 259-260.

Formatted: Font: Not Bold

Formatted: Font: Not Bold

| Walkley, A.J. and Black, J.A. (1934) Estimation of organic carbon by chromic acid titration method. *Soil Science*, 37: 29-38.

Formatted: Font: Not Bold

UNDER PEER REVIEW