

## Original Research Article

# **Evaluation of Cherry Tomato genotypes for yield and quality in polyhouse in Prayagraj agro-climatic condition [*Solanum lycopersicum* (L.) var. *cerasiforme* Mill.]**

## **ABSTRACT**

A field experiment entitled “Evaluation of Cherry tomato genotypes for yield and quality in polyhouse in Prayagraj agro-climatic condition [*Solanum Lycopersicum* (L.) var. *cerasiforme* Mill.]” was conducted from July, 2022 to December, 2022 at the Horticulture Research Farm, Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj. The experiment was laid out in Randomized Block Design (RBD) with three replications. The present investigation was carried out to find out the best cultivar (Yellow Cherry Tomato-round shape, Cherry Red-round shape, Nagmoti, Cherry Red-plum shape, Yellow Cherry Tomato-Pear shape and Pusa Cherry-1 .Pusa Cherry -1 was found the best in overall growth, yield and economic parameters like days of 1<sup>st</sup> flowering, no. of flower per cluster, no. of fruit per cluster, no. of fruits per plant, total yield, high cost benefit ratio.

Among all the varieties highest total yield was found in Pusa cherry-1 (142.9q/ha) whereas lowest in Yellow pear cherry tomato (74.6q/ha). The highest gross return (283883 Rs/ha) and cost benefit ratio (2.133) were obtained from Pusa Cherry-1. Considering the experimental findings variety Pusa Cherry -1 was found to be the most suitable for higher productivity and economic return under Prayagraj agro-climatic conditions.

**Keywords:** Cherry tomato, Growth, Yield and Quality

### **Introduction**

Fruits and vegetables have long been recommended for their high levels of vitamins, minerals, and phytochemicals, and India is the world's second-largest producer of these crops, accounting for 14% of global vegetable production. Despite this impressive output, close to 25% of India's agricultural production goes to waste. However, the country's vegetable production is particularly impressive, contributing 15% of global vegetable production despite utilizing only 3% of its arable land. These figures highlight the potential for India to increase its agricultural output while also reducing waste, which could have a significant impact on global food security and nutrition.

Cherry tomato (*Solanum lycopersicum* var. *cerasiforme*) is a warm season crop and requires long growing periods to reap more harvests, it is the most promising crop under protected structures as a small variety of tomato and generally considered to be similar but not identical to the wild precursor of the domestic tomato. It is characterized by small size fruits, with a bright red colour resembling a cherry, having an excellent taste. Cherry tomato is becoming

popular in the retail chains and marketed at a premium price compared to regular tomato. It is joining the growing market of mini vegetables and is one of the most promising in the line of differentiated products. It is considered as an exotic vegetable, bringing new taste and appearance to dishes.

Cherry tomato is a highly-priced culinary as well as it is an ornamental vegetable. Cherry tomatoes are normally much sweeter than large tomatoes. Cherry tomato has several medicinal values as it promotes gastric secretion, blood purification, intestinal antiseptic, cure cancer of the mouth and sour throat, apart from improving quality of the prepared foods. It is highly nutritious with good amount of vitamins. It is a good appetizer having pleasing test. Tomato juice contains lycopene one of the most powerful antioxidant and vitamin C which are most beneficial to human beings.

It has several medicinal values as it promotes gastric secretion, blood purification, intestinal antiseptic, cures cancer of the mouth and sour throat, apart from improving quality of the prepared foods. It is highly nutritious with good amount of vitamins. It is a good appetizer having pleasing taste. Cherry tomato juice contains lycopene, which is the important antioxidant and vitamin C are most beneficial to human beings.

Presently, cherry tomatoes are widely cultivated in Central America and are distributed in California, Korea, Germany, Mexico and Florida (Renuka *et al.*, 2014). About 24.00 per cent of retail sales of tomatoes in the U.S are contributed by cherry tomato. They are becoming popular in the retail chains and marketed at a premium price compared to regular tomatoes. Cherry tomato adaptation provides high possibilities for inclusion in breeding programs, using their valuable characteristics on genetic diversity for selecting parents, together with their large geographical diversity (Medina and Lobo, 2001).

The cherry tomatoes developed for fresh market and processing should have distinct quality characteristics (Kumar *et al.*, 2014). Therefore, the aim of the present study is to evaluate the different cherry tomato genotypes for yield and quality characters under shade net conditions in order to evaluate their potential for breeding programs.

## **Materials and Method**

Prayagraj is situated at an elevation of 98 meters above sea level at 25.87° N latitudes and 81.150 E longitudes. This region has a sub-tropical climate prevailing in the South-East part of U.P. with both the extremes in temperature, i.e., the winter and the summer. In cold winter months (Dec- Jan), the temperature falls 2-5°C or even low, while in summer months (May-June) it reaches as high as 49°C. The experiment was conducted in the Horticulture Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj, during the year 2022-23. The research experiment was laid out in Randomized Block Design comprising of 6 cherry tomato varieties with 3 replications which makes it a total of 18 plots. The transplanting of seedlings was accomplished on the last week of september, 2022 during rabi season. A total of 6 plants from each variety was sown at a spacing of 90 cm between rows and 60 cm between plants. The unit plot size was 1 m x 1 m. The varieties were allocated randomly to a

unit plot in each of the replication. Regular cultural practices, crop protection measures were taken as per the crop requirement. The crop was watered regularly. Observations were recorded as per the growth, yield and quality parameters and the mean values of data recorded were analyzed statistically by adopting the method suggested by Panse and Sukhatame. The performance of different varieties of cherry tomato was studied and data was collected on the basis of three categories of parameters. First are the growth parameters: Germination percentage, survival percentage, Plant height (cm), Number of branches per plant, Plant spread (cm), leaf area index. Second are the yield related parameters: days to first flowering, number of flowers per cluster, number of fruit per cluster, days to first harvesting, fruit length(cm), fruit diameter(cm), average fruit weight(g), number of fruit per plant(kg), average yield per plant, total yield (t/ha) qualitative characters: Total Soluble Solid (T.S.S) and Asorbic acid, Economic parameters cost of cultivation (Rs), gross return (Rs), Benefit cost Ratio.

## **Result and Discussion**

The result of the investigation based on the various observations viz., Plant growth, Yield and quality of different varieties of cherry tomato are presented and to determine best performing variety of cherry tomato in terms of growth, yield, and quality.

### **GROWTH PARAMETERS**

#### **Germination Percentage**

Significantly the maximum germination percentage 89.9 was recorded in variety Yellow Cherry Tomato (round) followed by variety Red Cherry (plum shape) 85.103 whereas minimum germination percentage 71.1 was recorded in variety Cherry Red (round shape).

#### **Survival Percentage**

Significantly the maximum survival percentage 88.297 was recorded in variety Nagmoti followed by variety Yellow Cherry Tomato (round) 85.803 whereas minimum survival percentage 70.3 was recorded in variety Red Cherry Tomato (plum shape).

#### **Plant Height (cm)**

The height of five randomly selected plants from each plot was measured from ground level to tip of the shoot at 30, 60 90,120 DAT and was recorded. Critical analysis of data displayed in table clearly marked out the obvious difference among the treatments with respect to plant height.

#### **Plant height at 30 days after transplanting:**

Non Significantly the maximum plant height 56.76 cm at 30 days after sowing was recorded in variety Red cherry (plum shape) followed by variety followed by cherry red 54.66 cm whereas minimum plant height 45.03 was recorded in variety Nagmoti.

#### **Plant height at 60 days after transplanting:**

Significantly the maximum plant height 101.93 cm at 60 days after sowing was recorded in variety Pusa cherry 1 followed by Red cherry (plum shape) 98.007 cm whereas minimum plant height 77.75 was recorded in variety Nagmoti.

**Plant height at 90 days after transplanting:**

Significantly the maximum plant height 189.203 cm at 90 days after sowing was recorded in variety red cheery (plum shape) followed by Pusa cheery 1 186 cm whereas minimum plant height 150.1 was recorded in variety Nagmoti.

**Plant height at 120 days after transplanting:**

Significantly the maximum plant height 243.46 cm at 120 days after sowing was recorded in variety Cherry red (round) followed by Yellow cherry tomato (round shape) 231.223 cm whereas minimum plant height 174.87 was recorded in variety Nagmoti.

**Numbers of primary branches per plant at 30 days after transplanting:**

Significantly the maximum number of primary branches 9.18 at 30 days after sowing was recorded in variety Cherry Red (round shape) followed by variety followed by Pusa cherry-1 8.137 whereas minimum number of primary branches 4.69 was recorded in variety Yellow Pear Cherry Tomato.

**Numbers of primary branches per plant at 60 days after transplanting:**

Significantly the maximum number of primary branches 11.64 at 60 days after sowing was recorded in variety Cherry Red (round shape) followed by variety followed by Pusa cherry-1 9.627 whereas minimum number of primary branches 5.25 was recorded in variety Yellow Pear Cherry Tomato.

**Numbers of primary branches per plant at 90 days after transplanting:**

Significantly the maximum number of primary branches 16.153 at 90 days after sowing was recorded in variety Cherry Red (round shape) followed by variety followed by Pusa cherry-1 12.853 whereas minimum number of primary branches 7.19 was recorded in variety Yellow Pear Cherry Tomato.

**Numbers of primary branches per plant at 120 days after transplanting:**

Significantly the maximum number of primary branches 18.42 at 30 days after sowing was recorded in variety Cherry Red (round shape) followed by variety followed by Pusa cherry-1 16.62 whereas, minimum number of primary branches 8.45 was recorded in variety Yellow Pear Cherry Tomato.

**Plant Spread (cm)**

Significantly the maximum plant spread (EW) 75.1 was recorded in variety Yellow Cherry Tomato (round shape) followed by variety Cherry Red (round) 66.7 whereas minimum plant spread 42.297 was recorded in variety Nagmoti.

Significantly the maximum plant spread (SW) 81.503 was recorded in variety Pusa Cherry-1 followed by variety Nagmoti 78.4 whereas minimum plant spread 42.003 was recorded in variety Yellow Cherry Tomato (round shape).

### Leaf Area Index (cm<sup>2</sup>)

Significantly the maximum leaf area index 6.627cm<sup>2</sup> was recorded in variety Yellow Cherry Tomato (round) followed by variety Yellow Cherry Tomato (Pear shape) 6.577 whereas minimum leaf area index 4.65 was recorded in Nagmoti.

**Chart 1 : Varieties and their sources**

SR.NO	NAME OF VARIETY	SOURCE OF VARIETY
1.	Yellow Cherry Tomato- Round Shape	CISH, Lucknow
2.	Cherry Red -Round	CISH, Lucknow
3.	Nagmoti	CISH, Lucknow
4.	Red Cherry- Plum Shape	CISH, Lucknow
5.	Yellow Pear Cherry Tomato	CISH, Lucknow
6.	Pusa Cherry 1	IARI, New Delhi

**Figure 1: Field experiment and variety of fruits.**




Plate 1



Plate 2



## **Yield Parameters**

### **Days of first flowering**

Significantly the maximum number of days taken to first harvesting 89.5 was recorded in variety Nagmoti followed by variety Pusa Cherry-1 84.5 whereas minimum number of days taken to first harvesting 66 was recorded in variety Yellow cherry tomato (pear shape).

### **No. of flowers per cluster**

Significantly the maximum number of flowers per cluster 11.78 was recorded in variety Pusa cherry 1 followed by variety yellow cherry (round) 9.01 whereas minimum number of flowers per cluster 7.713 was recorded in variety Yellow cherry tomato (pear shape).

### **No. of fruits per cluster**

Significantly the maximum number of fruits per cluster 11.47 was recorded in variety Pusa cherry -1 followed by variety yellow cherry (round) 8.56 whereas minimum number of fruits per cluster 3.437 was recorded in variety Yellow cherry tomato (pear shape).

### **Days to first harvesting**

Significantly the maximum number of days taken to first harvesting 89.5 was recorded in variety Nagmoti followed by variety Pusa Cherry-1 84.5 whereas minimum number of days taken to first harvesting 66 was recorded in variety Yellow cherry tomato (pear shape).

### **Fruit length (cm)**

Significantly the maximum fruit length 4.22 cm was recorded in variety Red cherry tomato (plum shape) followed by variety Yellow Cherry Tomato (round) 4.15cm whereas minimum fruits length 1.72 was recorded in variety Nagmoti.

### **Fruit Diameter (cm)**

Significantly the maximum fruit diameter 35 mm was recorded in variety Red Cherry (plum shape) followed by variety Cherry Red (round) 30 whereas minimum number of fruits diameter 15 was recorded in variety Nagmoti.

### **Average fruit weight (g)**

Significantly the maximum average fruit weight 14.59g was recorded in variety Red Cherry (plum shape) followed by variety Cherry Red (round) 14.02 whereas minimum average fruit weight 9.3 was recorded in variety Nagmoti.

### **Numbers of fruit per plant (kg)**

Significantly the maximum number of fruits per plant 63.8 was recorded in variety Pusa cherry-1 followed by variety Red Cherry (plum shape) 47.397 whereas minimum number of fruits per plant 24.6 was recorded in variety Yellow cherry tomato (pear shape).

### **Average yield per plot**

Significantly the maximum average yield per plot 14.29kg was recorded in variety Red Cherry (plum shape) followed by variety Cherry Red (round) 14.03 whereas minimum average yield per plot 7.46 was recorded in variety Yellow cherry tomato (pear shape).

### **Total yield (t/ha)**

Significantly the maximum total yield 142.9 was recorded in variety Pusa cherry-1 followed by variety Red cherry (plum shape) 140.3 whereas minimum total yield 74.6 was recorded in variety Yellow cherry tomato (pear shape).

### **Total soluble solids (°Brix)**

Significantly the maximum TSS 10.72 was recorded in variety Cherry Red (plum shape) followed by variety Yellow cherry tomato (round) 10.54 whereas minimum TSS 7.723 was recorded in variety Nagmoti.

### **Ascorbic acid**

Significantly the maximum ascorbic acid 20.94 was recorded in variety Pusa Cherry-1 followed by variety Yellow Cherry Tomato (round) 19.89 whereas minimum ascorbic acid 16.94 was recorded in variety Cherry Red (round).

**Table 2 - Days of first flowering, No. of flowers per cluster, No. of fruits per cluster, Days to first harvesting, Fruit length (cm), Fruit Diameter (mm), Average fruit weight (g), Numbers of fruit per plant, Average yield per plot(Kg), Total yield (q/ha), Total soluble solids (°Brix), Ascorbic acid (mg/100g)**



### Economics of different genotypes

Economics of all genotypes were calculated according to the expenditure occurred from then nursery till harvesting of fruits viz. Cost of cultivation, gross return, net return and benefit cost ratio has been worked out presented in table 3

Maximum cost benefit ratio was recorded in Pusa Cherry-1 (2.133) and minimum was recorded in Yellow Pear cherry tomato (1.88).

Table 3 Economic parameters

Sr. No.	Genotypes	Cost of cultivation	Gross return	Cost: Benefit ratio
1	<b>Yellow Cherry Tomato- Round Shape</b>	130401	255290	1.957
2	<b>Cherry Red -Round</b>	171402	326043	1.9
3	<b>Nagmoti</b>	149602	290689	1.943
4	<b>Red Cherry- Plum Shape</b>	140802	284699	2.023
5	<b>Yellow Pear Cherry Tomato</b>	163188	307161	1.88
6	<b>Pusa Cherry-1</b>	133089	283883	<b>2.133</b>
	<b>Mean</b>	<b>148080.67</b>	<b>291294.17</b>	<b>1.97</b>
	<b>CV</b>	<b>6.9</b>	<b>7.57</b>	<b>5.19</b>
	<b>SEm</b>	<b>5902.33</b>	<b>12727.92</b>	<b>0.06</b>
	<b>CD at 5%</b>	<b>18598.61</b>	<b>40106.48</b>	<b>0.19</b>
	<b>F Value</b>	<b>S</b>	<b>S</b>	<b>NS</b>

### CONCLUSION

In terms of yield parameters like days to first flowering, no. of flowers per cluster, no. of fruit per cluster, no. of fruit per plant, average yield per plot, total yield variety and in Quality parameters –Ascorbic acid and Benefit cost ratio of Pusa cherry -1 is best over all varieties.

## References

- Ahmad, F., Khan, O., Sarwar, S., Hussain, A. and Ahmad, S., 2007, Performance evaluation of tomato cultivars at high altitude Sarhad J. Agric., 23(3): 581-585.
- Alam, M.S., Sultana, N., Ahmad, S., Hossain, M.M. and Islam, A.K.M.A., 2010, Performance of heat tolerant tomato hybrid lines under hot, humid conditions. Bangladesh. J. Agril. Res., 35(3): 367-373de.
- Ali, A., Hussain, I., Khan, A., Khan, J., Rahman, M.U. and Riaz, A., 2016, Evaluation of various tomato (*Lycopersicon esculentum* Mill.) cultivars for quality, yield and yield components under agro climatic condition of Peshawer. J. Agric. Biolo. Sci., 11(2): 59-62.
- Ali. W., Jilani, M. S., Naeem, N., Waseem, K., Khan, J., Ahmad, M. J. and Ghazanfarullah., 2012., Evaluation of different hybrids of tomato under the climatic condition of Peshawar. Sarad J. Agri., 28(2): 207-212.
- Alley, E. W., Karl, H. N., John, T. W. and Davis, R. M., 1976, Estimation of chlorophyll and carotenoid contents of whole tomato by light absorbance technique. J. Food. Sci., 41(1976): 329-332.
- Bajaj, K. L., Mahajan, R., Kaur, P. P. and Chuma, D. S., 1990, Chemical constituents of processing tomatoes (*Lycopersicon esculentum* Mill.). J. Res. Punjab. Agric. Univ., 27 (2): 226-230.
- Benti, G., Degefa, G., Biri, A. and Tadesse, F., 2017, Performance evaluation of tomato (*Lycopersicon esculentum* Mill.) varieties under supplement irrigation at Erer valley, Babile district, Ethiopia. J. Plant. Sci., 5(1): 1-5.
- Berry, S. Z., Uddin, M.R., Gould, W. A., Bisges., A. D. and Dyer, G. D., 1988, Stability in fruit yield, soluble solids and citric acid of eight machine harvested processing tomato cultivars in Northern Ohio., J. Americ. Soc. Hort. Sci., 113(4):604-608.
- Bhati, V., 2017, Evaluation of tomato genotypes for growth, yield and quality traits under foothills condition of Nagaland, India. Int. J. Curr. Microbiol. App. Sci., 6(3):1645-1649.
- Binalfew, T., Alemu, Y., Geleto, J., Wendine, G. and Hinsermu., 2016, Performance of introduced hybrid tomato (*Solanum lycopersicum* Mill.) cultivar in the Rift valley, Ethiopia. Int. J. Res. Agric. Fores., 3(10): 25-28.
- Biswas, M., Sarkar, D. R., Asif, M. I., Sikder, R. K., Mehraj, H. and Jamaluddin, A. F. M., 2015, Comparison of growth and yield characteristics of BARI tomato varieties. J. Biosci. Agri. Res., 03(01): 01-07.
- Deepa, S. and Thakur, M. C., 2008, Evaluation of diallele progenies for yield and its contributing traits in tomato under mid-hill conditions. Indian J. Hort., 65 (3):297-301.
- Devkota, S., Shrestha, S.L., Dhakal, D.D., Shakya, S.M. and Pandey, A., 2018, Evaluation of tomato hybrids for yield attributes under Khumaltar condition. J. Int. Agil. Anim. Sic., 35:191-196.

- Dhyani, S., Misra, A.C., Panday, V. and Sajwan, P., 2017a, Evaluation of tomato (*Solanum lycopersicomL.*) hybrids for fruit yield characters in Hill region of Uttarakhand. *Int. J. Curr. Micobiol. App. Sci.*, 6 (9): 1622-1633.
- Dhyani, S., Misra, A.C. and Panday, V., 2017b, Assessment of tomato (*Solanum lycopersicomL.*) hybrids for fruit quality and yield characters in Hill region of Uttarakhand. *Int. J. Sci. Res.*, 7(9): 691-695.
- Dundi, K. B. and Mandalageri, B. B., 1991, Heterosis for shelf-life and its components in tomato (*Lycopersicon esculentum Mill.*). *South Indian Hort.*, 39: 353-355.
- Dunsin, O., Agbaje, G., Aboyeji., C.M. and Gbadamosi, A., 2016, Comparison of growth, yield and fruit quality performances of tomato varieties under controlled environment condition of the Southern Guinea Savannah. *J. Agric. Environ. Sci.*, 16(10): 1662-1665.
- Fish, W. W., Perkins, V. P. and Collins, J. K., 2002, Quantitative assay for lycopene that utilizes reduced volumes of organic solvents, *J. Fd. Comp. Anal.*, 15: 309-317.
- Ghosh, K.P., Islsm, A.K.M.A., Mian, M.A.K. and Hossain, M. M., 2010, Variability and character association in F<sub>2</sub> segregating population of different commercial hybrids of tomato (*Solanum lycopersicumL.*). *J. Appl. Sci. Environ. Manage.* 14 (2): 91-95
- Gomathi, S. P. 2008. Development of semi determinate F<sub>1</sub>hybrids in tomato (*Solanum lycopersicum Mill.*) with combined resistance to viral disease and nematodes. M.Sc. Thesis, TNAU, Coimbatore.
- Hasan, M.M., Bari, M.A.A. and Hossain., M.A., 2016, Genetic variability and traits association analysis of tomato (*Lycopersicon esculentumL.*) genotypes for yield and quality attributes. *Uni. J. Plant. Sci.*, 4(3): 23-34.
- Islam, M.S., Mohanta, H.C., Ismail, M.R., Rafii, M.Y. and Malek, M.A., 2012, Genetic variability and trait relationship in cherry tomato (*Solanum lycopersicumvar. cerasiforme (Dunnal) A. Gray*). *Bangladesh J. Bot.*, 41(2):163-167.
- Jasmine, J. A. P. and Ramadass, S., 1994, Qualitative evaluation of tomato hybrids and varieties. *South Indian Hort.*, 42 (1): 26-28.
- Joshi, A. K., Kumar, A. and Sharma, B. k., 1998a, Evaluation tomato genotypes for horticultural traits. *Punjab vegetable grower.* 33: 21-22.
- Kamimura, S., K., Ito, H., Yoshikawa, S., Monma. and Kanna. T., 1985, "Furikoma" -New tomato variety for processing. *Bull. Veg. Orn. Crops Res.*, 5:47.
- Kanneh, S. M., Quee, D.D., Ngegba, P. M. and Musa, P. D., 2017, Evaluation of tomato (*Solanum lycopersicumL.*) genotypes for horticultural characteristics on the Upland in Southern Sierra Leone. *J. Agri. Sci.*, 9(6): 213-220.
- Kaushik, S. K., Tomar, D. S. and Dixit, A. K., 2011, Genetics of fruit yield and it's contributing characters in tomato (*Solanum lycopersicom*). *J. Agric. Bio. and Sustainable Dev.*, 3(10): 209-213. Lekshmi, S. L. and Celine, V. A., 2015., Evaluation of tomato hybrids for fruit, yield and quality traits under polyhouse conditions. *Int. J. App. Pure Sci. Agri.*, 01(7): 58-64.

Mahendrakar, P., Mulge, R. Madalageri, M. B. Patil, M. S., Ravi, B. A. and Chandan, K., 2006, Exploitation of hybrid vigour for growth and yield parameter in tomato. ATSH, 33.

Malavika, O., Indira, P. and Sheela, K.B., 2017, Performance evaluation of cherry tomato genotypes under rain shelter. J. Tro. Agrci., 55(2): 180-183.

Najeema, M.H., Revanappa, H.P., Hadimani. and Birador, I.B., 2018 Evaluation of cherry tomato (*Solanum lycopersicum*L. var *cerasiforme*) genotypes for yield and quality traits. Int. J. Curr. Microbiol. App. Sci., 7(6):2433-2439.

Nitzsche, P., Tietjen, W., Kline. W. and Garrison.S., 2003. Evaluation of grape and cherry tomatoes in North New Jersey. In: Tomato varieties/ trial reports-2003. <https://njaws.rutgers.edu/>.

Omprasad, J., Reddy, S. S. P., Madhumathi, C. and Balakrishan, M., 2018a, Evaluation of cherry tomato under shade Net for growth and yield attributes. Int. J. Curr. Microbiol. App. Sci., 7:700-707.

Omprasad, J., Reddy, S. S. P., Madhumathi, C. and Balakrishan, M., 2018b, Evaluation of cherry tomato for quality characters under shade net. Int. J. Curr. Microbiol. App. Sci., 7:700-707.

Panse, V. G. and Sukhatme, P. V., 1985, Statistical methods for agricultural workers, ICAR, New Delhi.

Parmar, D. K., Thakur, D. R., Jamwal, R. S. and Gurudevsing., 2018, Evaluation of tomato cultivar for yield, profit and quality performance in an organic management system in North Western Himalayas, India. Int. J. Curr. Micro. App. Sci., 7(10): 498-506.

Venkadeswaran, E., Vethamoni, I. P., Arumugam,T., Manivannan,N. and Harish,S., 2018a, Evaluation and selection of cherry tomato (*Solanum lycopersicum*L. var *cerasiforme*) genotypes for growth and yield contribution characters. Int. J. Curr. Micro. App. Sci., 7 (6): 155-1165.

Venkadeswaran, E., Vethamoni, I. P., Arumugam,T., Manivannan,N. and Harish,S., 2018b, Evaluating the yield and quality characters of cherry tomato (*Solanum lycopersicum*L. var *cerasiforme*) genotypes. Int. J. Chem. Stud., 7 (6): 155-1165.