

Cumulative Effect of Fertilizer and Organic Spray (Banana Pseudostem Sap or Vermiwash) on Yield, Quality, Uptake and Economics of Linseed (Linum usitatissimum L.)

ABSTRACT

The study on linseed aimed to investigate the effectiveness of fertilizer and organic spray of banana pseudostem sap or vermiwash on yield, quality, uptake and economics of the crop was conducted at College Farm, N. M. College of Agriculture, Navsari Agricultural University, Navsari during rabi season of 2019-20. The experiment was laid out in Randomized Block Design with four replicates. The experimental field was fairly leveled and uniform. The treatments comprising of T1(50% RDF), T2 (50% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS), T3 (50% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS), T4(75% RDF), T5 (75% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS), T6 (75% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS), T7(100% RDF), T8 (100% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS) and T9(100% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS). The result indicated significant differences on seed yield, oil yield and nutrient uptake in the seed and stover. Treatment T9 outperformed the other treatments by registering maximum gross return of ₹94611 ha⁻¹, net return (₹63679 ha⁻¹) and benefit cost ratio (3.06). It could be concluded that to obtain better yield, quality, uptake and economics of rabi linseed crop, the crop should be applied with 100% RDF (60-30-00 N-P2O5-K2O kg ha⁻¹) and foliar spray of 1% vermiwash or 1% banana pseudostem sap at 30 and 60 DAS.

Formatted: Font: Bold

Formatted: Justified, Line spacing: single

Keywords: Linseed; yield; quality; economics; vermiwash; banana pseudostem sap.

1. INTRODUCTION

India is one of the leading oilseeds growing country in the World and fourth largest vegetable oil economy next to USA, China, and Brazil. The diverse agro-ecological conditions in the country are favorable for growing many annual oilseed crops, which include groundnut, rapeseed, mustard, soybean, sunflower, sesame, safflower, niger, castor and linseed. Linseed also known as flax is one of the oldest oilseed crops grown in almost all the countries of the world. Linseed (Linum usitatissimum L.), one of the oldest oilseed crop cultivated either as oilseed crop or fibre crop, is perhaps one of the most antique plant. Linseed is self-pollinated crop which is widely adapted to temperate climate of the world. Linseed is an annual herbaceous plant and grows to a height of 30 to 120 cm. Fiber types are tall growing and less branched than the seed types. The stem is glabrous greyish green with leaves narrow and alternate. Fruit is a capsule, globular in shape mostly indehiscent. Seeds are compressed shining yellow or light brown. Linseed or flax is one of most important rabi oilseed crop. It contains 35 to 45 % oil.

Formatted: Justified, Line spacing: 1.5 lines

Fertilizer application is an important option to [enhance](#) plant nutrients and improve crop yield. Nitrogen is a structural component of chlorophyll and protein therefore, adequate supply of nitrogen is beneficial for both carbohydrate and protein metabolism as it promotes cell division and cell enlargement, resulting in more leaf area and thus ensuring better growth and development of plant and improve yield. Phosphorus is an important plant nutrient which helps in root development and requires for grain formation to maturity of grain and ultimately improve crop yield of linseed (Parmar et al., [2020](#)).

[Banana Pseudostem Sap](#)

Banana is the heavy nutrient feeder crop [and its](#) pseudostem sap (BPS) has been extracted from the outer sheath of pseudostem of banana. [The sap is](#) colorless clean water immediately after extraction. These are vital for plant growth and development (Islam et al.). [The banana sap](#) contains essential macro and micronutrients, growth promoting substances like cytokinin and Gibberelic Acid ([GA3](#)) and this mixture [when](#) inoculated with different microbes like Rhizobium, Azatobacter, etc.[it plays](#) an important role in enhancing yield.

Formatted: Justified, Line spacing: 1.5 lines

[Vermiwash](#)

Vermiwash is a liquid extract produced from vermicompost in a medium where earthworms are richly populated. Vermiwash [has](#) micronutrients [and composed of](#) beneficial soil microbes and also contain 'plant growth hormones [and](#) enzymes'. [There is very little information on banana pseudostem sap and vermiwash therefore](#); the [study aims to](#) investigate [their effectiveness](#) on [linseed production](#).

Formatted: Justified, Line spacing: 1.5 lines

2. MATERIALS AND METHODS

[Navsari Agricultural](#) University campus is geographically located at 20°57' N latitude and 72° 54' E longitude with an altitude of 10 m above mean sea level. Navsari falls under Agro-ecological [situation-III](#) of South Gujarat Heavy Rainfall Zone which is characterized by fairly warm summer, mild winter and warm humid monsoon with heavy rainfall. The week-wise meteorological data pertaining to the maximum and the minimum temperatures, relative humidity and evaporation during the period of experimentation (November-2019 to March-2020) [was](#) recorded at the meteorological observatory of college farm, N. M. College of Agriculture, Navsari. The soils [of](#) the experimental site was characterized by dark greyish brown type [with](#) medium to poor drainage, flat topography and good water holding capacity [with](#) clayey (65.70 %) in texture, low in available nitrogen (180.24 kg ha⁻¹), medium in available

phosphorus (35.30 kg ha⁻¹) and high in available potassium (281.50 kg ha⁻¹). The experiment was laid out in Randomized Block Design [with](#) nine treatments ([Table 1](#)) [with four replicates](#). [The present experiment was carried out with local variety of linseed. The spacing was kept between row 30 cm and between plant 10 cm. Oil content of seed was determined by Soxhlet apparatus as per the method suggested by Tiwari et al. \(2011\).](#)

Table 1: List of experimental treatments

Treatment	Treatment Description
T1	50% RDF
T2	50% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS
T3	50% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS
T4	75% RDF
T5	75% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS
T6	75% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS
T7	100% RDF
T8	100% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS
T9	100% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS

Formatted Table

Statistical procedures appropriate to the randomized block design [were](#) as described by Panse and Sukhatme (1985) and the [mean](#) differences tested by F- test. Summary table for treatment mean [were](#) furnished with standard error of mean (S.Em.) and co-efficient of variation (%). Critical Difference (C.D.) at 5 % level of probability [was](#) calculated where the treatment effects were found significant.

Formatted: Justified, Line spacing: 1.5 lines

3. RESULTS AND DISCUSSION

3.1 Effect of [banana pseudostem sap and vermiwash](#) on yield

[Results](#) in [Table 2](#) [showed](#) significant [differences on linseed and stover yields in](#) treatment T9(100% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS). However, [treatments](#) T8, T7, T6 and T5 [registered equally as T9](#). High [linseed yield \(???\)](#) was recorded with the application of organic spray over other treatments. [The high seed yield could](#) be attributed to high number of capsules [and seed per](#) plant and adequate supply of nutrient element at the right time from organic spray and inorganic fertilizers which ultimately helps to improve seed yield as well as stover yield of linseed crop. Similar results were reported by Meena et al. (2011), Khajani et al. (2012) and Tanvi patle et al. (2022).

3.2 Effect of banana pseudostem sap and vermiwash on quality

The quality of produce in the treatments was judged by oil content and oil yield in the seed. Results in Table 2 showed no significant differences on oil content. However, T9 (100% RDF + foliar spray of 1% banana pseudostem sap at 30 and 60 DAS) recorded the maximum oil yield (449 kg ha⁻¹). This was followed by treatment T8 (440 kg ha⁻¹), T7 (402 kg ha⁻¹) and T6 (380 kg ha⁻¹). The oil yield correlated with seed yield and oil content in seed. This could be due to high seed yield under corresponding treatments. These results are in accordance with the findings of Esmail et al. (2014) and Nitin Sood et al. (2020).

Formatted: Justified, Line spacing: 1.5 lines

3.3 Effect of NPK content and uptake by crop

The results in Tables 3 and 4 pertaining to nitrogen, phosphorus and potassium content in seed and stover of linseed were not significant. However, treatment (T9) recorded high nitrogen, phosphorus and potassium uptake by the seed of linseed. Treatments (T8) and (T7) recorded figures close to T9. Application of 100% RDF without organic liquid spray recorded 44%, 39.78% and 42.91% nitrogen, phosphorus and potassium respectively compared to 50% RDF which registered 19.45%, 19.66% and 21.64% nitrogen, phosphorus and potassium respectively. 75% RDF registered the lowest figures for nitrogen, phosphorus and potassium. Similar trend was observed in the case of fertilizer levels with foliar spray of 1% vermiwash or foliar spray of 1% banana pseudostem sap at 30 and 60 DAS. The results in Table 4 showed significant differences on the uptake of nitrogen, phosphorus and potassium by linseed particularly in the stovers. The application of 100% RDF + foliar spray of 1% banana pseudostem sap at 30 and 60 DAS (T9) recorded the maximum uptake of nitrogen (15.93 kg ha⁻¹), phosphorus (9.38 kg ha⁻¹) and potassium (17.15 kg ha⁻¹) by stover of linseed. Treatments (T8) and (T7) registered close to T9. This could be due to the increased uptake the macronutrients in seed or stover yield and the application of inorganic fertilizer with organic liquid spray (banana pseudostem sap or vermiwash) increased the availability of nitrogen, phosphorus and potassium to the plant. The results are in conformity with the findings obtained by Zafar et al. (2019) and Nitin Sood et al. (2020).

Formatted: Justified, Line spacing: 1.5 lines

3.4 Effect of NPK, banana pseudostem sap and vermiwash on economics

The regional adaptability of any agronomic practices in any crop is completely based on the highest economic return. The results in Table 5 in respect to cost of cultivation, gross return, net return and benefit cost ratio was not significant. However, the maximum net return (₹63679 ha⁻¹) was recorded in treatment T9 followed by treatment T8 (₹63579 ha⁻¹) and T7 (₹57304 ha⁻¹). In case of BCR, maximum

was 3.06 in treatments T9 and T8. This could be attributed by the high seed and stover yield and comparatively lower cost of cultivation.

4. CONCLUSION

Based on the findings of the experiment, it could the application of fertilizer 60-30-00 N-P₂O₅-K₂O kg ha⁻¹ with 100% RDF and foliar spray of 1% vermiwash or 1% banana pseudostem sap at 30 and 60 DAS is profitable in terms of yield and quality of linseed.

Formatted: Justified, Line spacing: 1.5 lines

REFERENCES

Esmail AO, Yasin HS and Mahmood BJ. (2014). Effect of levels of phosphorus and iron on growth, yield and quality of flax. *Journal of Agriculture and Veterinary Science*. 2014;7(5): 07-11.

Islam MS, Kasim S, Amin AM, Hun TG, Alam K and Hague MA (2022). Banana-pseudostem sap growing media as a novel source of phytochemicals mineral nutrients: influence on seedling growth of sweet corn. ISSN: 0718-5839.

Khajani FP, Irannezhad H, Majidian M and Oraki H (2012). Influence of different levels of nitrogen, phosphorus and potassium on yield and yield components of flax seed oil (*Linum usitatissimum* L.) variety Lirina. *Journal of Medicinal Plants Research*. 2;6(6): 1050-1054.

Meena RL, Singh TK, Kumar R, Singh AK and Hariom (2011). Production performance of linseed (*Linum usitatissimum* L.) to fertility and seed rates in dry land conditions of eastern Uttar Pradesh. *Indian Journal of Soil Conservation*.;39(3): 230-235.

Nitin Sood, Pankaj C, Rana SS and Naveen Datt (2020). Effect of nutrient management practices and varieties on yield, NPK uptake, quality and soil properties in linseed (*Linum usitatissimum* L.). *International Journal of Current Microbiology and Applied science*. 2020;9(12): 250-255.

Panse VG and Sukhtme PV (1985). Statistical method for agricultural workers. *Indian Council of Agricultural Research Publication*.;87-89.

Parmar SK, Thanki JD, Tandel BB and Pankhaniya RM (2020). Effect of nitrogen phosphorus and sulphur application on yield, quality, uptake and economics of linseed (*Linum usitatissimum* L.). *International Journal of Chemical Studies*. 2020;8(5): 1956-1960.

Formatted: Line spacing: 1.5 lines

[Tanvi patle, Chack S, Singh T, Singh AT and Bharti V \(2022\). Integrated nutrient management in linseed \(Linum usitatissimum L.\). The Pharma Innovation Journal. 11\(10\): 874-878.](#)

Tiwari PN, Gamhith PN [and](#), Rajan TS [\(2011\)](#). Rapid and non destructive determination of oil in oilseeds. [Journal of the American oil Chemists Society. 2011;51: 10-49.](#)

Zafar A, Sarwar G, Hussain S, Aftab M, Sarfraz M, Manzoor MZ, Riaz A, Naz A [and](#), Niaz A [\(2019\)](#). Effect of various rates of NPK on concentration, uptake and use efficiency of nutrients in linseed under saline sodic soil environment. [International Journal of Biosciences. 2019;15\(6\): 8-15.](#)

Formatted: Line spacing: 1.5 lines

UNDER PEER REVIEW

Table 2: Impact of nutrient management practices on yield and quality of linseed

Treatment	Yield		Quality	
	Seed yield (kg ha ⁻¹)	Stover yield (kg ha ⁻¹)	Oil Content (%)	Oil Yield (kg ha ⁻¹)
T ₁ - 50% RDF	690	1305	35.54	247
T ₂ - 50% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS	819	1491	36.12	297
T ₃ - 50% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS	806	1513	36.63	296
T ₄ - 75% RDF	891	1679	36.81	328
T ₅ - 75% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS	1006	1819	37.73	379
T ₆ - 75% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS	1012	1872	37.57	380
T ₇ - 100% RDF	1060	1944	37.93	402
T ₈ - 100% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS	1156	2031	38.07	440
T ₉ - 100% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS	1157	2051	38.85	449
SEM±	55.32	85.67	1.15	23.80
CD at 5%	162	250	NS	69
CV%	11.58	9.81	6.20	13.31

Table 3: Impact of nutrient management practices on content and uptake of nutrient by seed of linseed

Treatment	Nutrient content (%)			Nutrient uptake (kg ha ⁻¹)		
	N	P	K	N	P	K
T ₁ - 50% RDF	2.08	0.58	1.37	14.42	3.98	9.39
T ₂ - 50% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS	2.21	0.60	1.38	18.01	4.96	11.16
T ₃ - 50% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS	2.28	0.63	1.43	18.39	4.70	11.33
T ₄ - 75% RDF	2.34	0.60	1.46	20.95	5.31	12.89
T ₅ - 75% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS	2.36	0.63	1.58	23.82	6.36	15.87
T ₆ - 75% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS	2.33	0.65	1.57	23.95	6.60	15.92
T ₇ - 100% RDF	2.45	0.62	1.55	26.01	6.61	16.45
T ₈ - 100% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS	2.47	0.65	1.62	28.60	7.49	18.73
T ₉ - 100% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS	2.49	0.66	1.65	28.67	7.64	19.04
SEm±	0.10	0.02	0.06	1.49	0.42	0.83
CD at 5%	NS	NS	NS	4.36	1.24	2.44
CV%	9.10	9.02	9.11	13.30	14.25	11.54

Table 4: Impact of nutrient management practices on content and uptake of nutrient by stover of linseed

Treatment	Nutrient content (%)			Nutrient uptake (kg ha ⁻¹)		
	N	P	K	N	P	K
T ₁ - 50% RDF	0.71	0.39	0.76	9.17	5.10	9.97
T ₂ - 50% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS	0.73	0.43	0.78	10.82	6.44	11.64
T ₃ - 50% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS	0.73	0.42	0.77	11.02	6.35	11.66
T ₄ - 75% RDF	0.72	0.45	0.78	12.10	7.49	13.11
T ₅ - 75% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS	0.74	0.42	0.79	13.44	7.60	14.31
T ₆ - 75% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS	0.73	0.44	0.82	13.61	8.56	15.27
T ₇ - 100% RDF	0.74	0.45	0.80	14.48	8.99	11.49
T ₈ - 100% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS	0.75	0.44	0.81	14.95	8.85	16.25
T ₉ - 100% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS	0.77	0.46	0.83	15.93	9.38	17.15
SEm±	0.02	0.01	0.03	0.73	0.51	0.92
CD at 5%	NS	NS	NS	2.13	1.50	2.68
CV%	6.39	9.21	10.03	11.41	13.50	13.27

Table 5: Impact of different treatments on economics of linseed

Treatment	Gross return (₹ ha ⁻¹)	Cost of cultivation (₹ ha ⁻¹)	Net return (₹ ha ⁻¹)	B:C ratio
T ₁ - 50% RDF	56505	28363	28142	1.99
T ₂ - 50% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS	67011	29855	37156	2.24
T ₃ - 50% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS	65993	29855	36138	2.21
T ₄ - 75% RDF	72959	28904	44055	2.52
T ₅ - 75% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS	82299	30396	51903	2.71
T ₆ - 75% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS	82832	30396	52436	2.73
T ₇ - 100% RDF	86744	29440	57304	2.95
T ₈ - 100% RDF + Foliar spray of 1% vermiwash at 30 and 60 DAS	94511	30932	63579	3.06
T ₉ - 100% RDF + Foliar spray of 1% banana pseudostem sap at 30 and 60 DAS	94611	30932	63679	3.06

*Selling price: Linseed seed – 80 ₹ kg⁻¹

*Cost of Input: Linseed 120 ₹ kg⁻¹, Urea-5.93 ₹ kg⁻¹ and SSP-7.44 ₹ kg⁻¹