

Standardization of different growing environment conditions for button rose

Abstract

An investigation on Standardization of different growing environment conditions for button rose was carried out at Floriculture Research Station, Thovalai, Tamil Nadu during the year 2017-18 in randomized block design with five treatments of growing condition. The result revealed that, the significantly influence in button rose, the treatment T₃ (35% shade net only as top cover) recorded the highest plant height (53.38 cm), number of shoots/plant (7.00) and number of flowers in a cluster (9.75). The same treatment exhibited the highest number of leaves (77.75), early flowering (18 days) and early bud initiation after previous flowering (14.75 days) which was on par with the treatment T₅ (open condition) which recorded 76.50, 18.75 days and 15.00 days respectively at three months after planting. Among the treatments, the treatment T₂ (50% shade net) recorded the lowest plant height (46.63 cm), number of leaves/plant (70.75), number of shoots/plant (3.50), number of flowers in a cluster (6.50) and delayed flowering (20.75 days) and bud initiation after previous flowering (20.00 days). In case of yield parameters T₅- Open condition recorded maximum individual flower weight (1.06 g), Number of petals per flower (65.23) and Flower yield per plant / year (125.38g)

Keywords: Rose, shade net house, growth, flower yield,

Introduction

Rose (*Rosa indica*) is a tropical and subtropical plant which belongs to family Rosaceae. Rose is one of nature's most beautiful creations and is universally known as 'Queen of flowers' for its shape, different sizes, attractive colours and most delightful fragrance with varied uses. Roses are most popular flowers among many floriculture crops of the world. The major rose producing states in India are Karnataka, Maharashtra, Punjab, Uttar Pradesh, Delhi, Chandigarh, West Bengal, Himachal Pradesh, Rajasthan, Kashmir and Gujarat. The rose word derived from the Latin word "Erose" meaning the "God of love". The rose has 200 species and more than 20,000 varieties (Gauchanet *al.*, (3). Among these the important categories of roses are Hybrid Tea, Floribunda, Hybrid perpetual, Grandifloras, Tea scented china roses, Polyanthas, Miniatures, Cabbage rose, Bourbon rose, French rose alba, Musk rose, Mambler's moss roses, Noisette roses. Rose is a symbol of love, adoration and innocence and it occupies a prominent position in the tradition, religious and social culture of every country in the world. Rose as cut flower has great demand in the internal as well as export

markets. Rose can be successfully cultivated in mild climate with good sunshine. The flower production in rose can be increased by using different growing structure like polyhouse, shade net. As it is difficult to obtain good quality cut flowers under open conditions throughout the year. Therefore, the crops should be cultivated under the cover to get good quality produce (Polyhouse, Shade net), it is large enough to grow crops under partial or fully controlled environmental conditions to obtain optimum growth and quality production. The main advantages of polyhouse and Shade net cultivation are the crops can be cultivated successfully throughout the year, getting high productivity with excellent quality, it is easy to protect the crops against extreme climatic conditions and incidence of pests and disease. Climatic factors play a vital role in the production of quality roses.

Materials and Methods

The present research was carried out at Floriculture Research Station, Thovalai to 'Standardization of different growing environment conditions for button rose'. The present study was carried out during 2019-2020 in randomized block design to study effect of different growing condition on vegetative, flowering and yield parameter of button rose with five treatments viz., T₁-Shade net (35%), T₂-Shade net (50%), T₃-35% shade net only as top cover, T₄-50% shade net only as top cover and T₅-Open condition. The treatments were imposed on new budded plants of button rose and each treatment consisting of four plants was replicated four. Observation was recorded on four randomly selected plants in each treatment i.e., highest plant height, number of shoots/plants, number of leaves, number of flowers in a cluster, early flowering and early bud initiation after previous flowering at three months after planting. The experimental data were analysed statistically by ANOVA (Analysis of Variance) technique (Panse and Sukhatme, (7).

Result and discussion Results and discussion is more appropriate

The field preparation was done and button rose were planted under different growing environments viz., under 35% shade net, 50% shade net, 35% and 50% shade net only as top cover and purely under open conditions.

Growth parameters

The vegetative growth parameter like plant height, maximum number of branches, number of leaves in rose shows significant effect on plant growth. The treatment T₃ *i.e.* shade net house were recorded significantly in button rose, the treatment T₃ (35% shade net only as top cover) recorded the highest plant height (53.38 cm), number of shoots/plant (7.00), the highest number of leaves (77.75), which was followed by T₅-Open condition plant height (52.13 cm), the highest number of leaves (76.50 cm) open condition this might be due to optimum light intensity in combination with warmer environment and higher relative humidity inside the polyhouse which helped in increase in height as compared to open condition and shade environments (Table 1). These results are in agreement to those reported by the Mohanty *et al.* (5). Shade net house this might be due to Reduction of light intensity to optimum condition, increased temperature and relative humidity under polyhouse might have favoured early sprouting of buds and increase in branches as compared to open condition. The present findings are in the same line as reported by Moe (6). Significantly maximum number of leaves (77.75) were also recorded in treatment T₃ *i.e.* shade net house this might be due to the long waves of light which enter into the polyhouse might have converted into short wave, the short waves were trapped into the polyhouse which helps to increase the inner temperature and CO₂ inside the polyhouse which increase the photosynthesis rate of plant and simultaneously increase the number of leaves. (Patil *et al.* 2012).

Flowering parameters

The data revealed that the significantly the treatment T₃ *i.e.* shade net house recorded maximum number of flowers in a cluster (9.75). The same treatment exhibited early flowering (18 days) and early bud initiation after previous flowering (14.75 days) which was on par with the treatment T₅ (open condition) which recorded 76.50, 18.75 days and 15.00 days respectively at three months after planting (Table 1). Among the treatments, the treatment T₂ (50% shade net) recorded the lowest plant height (46.63 cm), number of leaves/plant (70.75), number of shoots/plant (3.50), number of flowers in a cluster (6.50) and delayed flowering (20.75 days) and bud initiation after previous flowering (20.00 days). This might be due suitable growing condition with optimum light intensity, favorable temperature and moist air inside the shade net house might have helped in the faster growth and early appearance of flower buds as observed in the present study. This might have favored accumulation of more carbohydrate than other growing environments. It is the tendency of plant to come to reproductive phase early if it has enough carbohydrate in it (Malhotra and Kumar, (4). Shade net house condition this might be due to, in shade net house plant get maximum sunshine which helps to increase the number of flowers per cluster and early flowering in button rose. Malhotra and Kumar (4) also observed these characters of roses under polyhouse as compared to open condition. The results of the present investigation indicated that number of

flowers per plant were the maximum in open field situation. This might be due to the fact that the total number of leaves were maximum under open field which accumulated more carbohydrates and were directly used for increasing the number of flowers. This might be also due to a greater number of shoots in open field which had produced a greater number of flowers Patilet *al.* (8). From the above results it is noticed that, maximum number of flowers per plant were observed in variety Gladiator. This might be due to the varietal characters of the various varieties. These results are in line with those of results obtained by Tabassumet *al.* (10) in rose cultivars and reported maximum number of flowers plant-1 in rose cv. Daydream among rose cultivars. The variation in number of flowers per plant might be due to the varietal character and another one reason is environmental conditions. This result was confirmed by Ranchanaet *al.* (9).

Yield parameters

The treatment T₅ i.e. shade net house recorded significantly maximum individual flower weight (1.06 g), Number of petals per flower (65.23), Flower yield per plant / year (125.38g) (Table 2) this might be due to the microclimatic conditions in polyhouse which helps to increase the flower yield and more light intensity, warmer environment. The same result was noticed by Talukdaret *al.* (11). However, performance of button rose under shadenet and open condition were statistically comparable with each other. The similar findings were also observed by Mohantayet *al.* (5). Similar findings were observed by Bhattacharjeeet *al.* (1) and Fascella and Zizzo (2).

Table 1. Influence of growing environment in button rose at 3 months after planting

Treatments	Plant height (cm)	No. of leaves / plant	No. of shoots / plant	Days taken for flowering	No. of flowers in a cluster	Days taken for bud initiation after flowering
T ₁	49.13	72.00	4.25	20.00	7.75	19.00
T ₂	46.63	70.75	3.50	20.75	6.50	20.00
T ₃	53.38	77.75	7.00	18.00	9.75	14.75
T ₄	51.13	75.00	5.75	19.25	8.75	17.00
T ₅	52.13	76.50	5.50	18.75	9.00	15.00
Mean	50.48	74.4	5.20	19.35	8.35	17.15
SEd	0.50	0.62	0.51	0.41	0.34	0.65
CD (0.05)	1.10	1.36	1.12	0.90	0.73	1.42

Table.2. Influence of growing environment on growth and yield of button rose

Treatment	Individual flower weight (g)	No. of petals / flower	Flower yield / plant / year (g)
T ₁	0.94	63.18	112.48
T ₂	0.90	58.73	111.10
T ₃	0.95	64.48	117.33
T ₄	0.90	62.43	114.30
T ₅	1.06	65.23	125.38
Mean	0.95	62.81	116.12
SEd	0.0116	0.1057	0.0878
CD (0.05)	0.0253	0.2302	0.1913

Conclusion English should be checked properly with the addition of few lines describing the area of study and the speciality

The button rose, the treatment T₃ (35% shade net only as top cover) recorded the highest plant height (53.38 cm), number of shoots/plant (7.00) number of leaves (77.75), number of flowers in a cluster (9.75), early flowering (18.00 days), early bud initiation after previous flowering (14.75 days), whereas yield parameters T₅- Open condition recorded maximum individual flower weight (1.06 g), Number of petals per flower (65.23) and Flower yield per plant / year (125.38g).

References Reference section must be checked to comply with the format of the journal

1. Bhattacharjee SK, Singh VC, Saxena NK. Studies on vegetative growth, flowering, flower quality and vase life of roses. Singapur J Primary Industries. 1993;21(2):67-71.
2. Fascella G, Zizzo GV. Evaluating the productivity of red rose cultivars in soil less culture. ActaHorti. 2007; 751:138-140.
3. Gauchan DP, Pokhrel AR, Pratap M, Lama P. Current status of cut flower business in Nepal. Kathmandu Uni. J Sci. Eng. Technol. 2009; 5:87-98.
4. Malhotra R, Kumar R. Effect of different treatments overhead shading and polythene covering on rose Cv. Raktagandha. Proceedings of National symposium on Recent

Trends and Future Strategies in Ornamental Horticulture, Uni. Agri. Sci., Dharwad, Karnataka (India) 2000.

5. Moe R. Effect of day length, light intensity and temperature on growth and flowering of roses. *J Amer. Soc. Hort. Sci* 1972;97(6):796-800.
6. Mohanty CR, Mohanty A, Das AB, Kar DS. Studied on Comparative performance of some rose varieties under open and protected environment. *The Asian J of Hort.* 2011;6(2):288-293.
7. Panse VG, Sukhatme PV. Statistical methods for agricultural workers. ICAR Publication, New Delhi. 1978, 359.
8. Patil SJ, Patel NL, Gaikwad SS, Bhalerao PP. Flower production of hybrid tea rose (*Rosa hybrida* L.) cv. 'GLADIATOR' under protected condition through different treatments during winter season. *Asian J of Hort* 2012;7(1):154-159
9. Ranchana P, Selvaraj N, Jawaharlal M, Kumar CSR, Anita B. Cut rose varieties suitable for organic farming system in Nilgiris, Tamil Nadu. *J Bio. Chem. Research.* 2014;31(1):104-108.
10. Tabassum R, Ghaffoor A, Waseem K, Nadeem MA. Evaluation of rose cultivars as cut flower production. *Asian J. Pl. Sci.* 2002; 1:668-669.
11. Talukdar M, Barooah S. *ActaHort* 1987;205:145-148