

Original Research Article

The Prevalence of Physicians' Burnout Syndrome in Ibrahim Malik Teaching Hospital, Khartoum –Sudan ,2020

Abstract:

Background: Burnout is described as exploratory and qualitatively as a state of emotional exhaustion due to excessive demands on energy, strength or resources in workplace. It is characterized by physical symptoms such as exhaustion, fatigue, frequent headache, and gastrointestinal disorders, sleeplessness and behavioral signs include frustration, anger, a suspicious attitude omnipotence or overconfidence, cynicism and signs of depression.

Methods: This descriptive cross-sectional hospital-based study was conducted in Ibrahim Malik teaching hospital in Khartoum, Sudan. This research was conducted in Ibrahim Malik Teaching Hospital, located in Khartoum State, Sudan. It was founded in 1977 and it is a governmental owned hospital providing many and various facilities to the general public. The hospital provides a 24/7 Emergency service, as well as many outpatients' clinic. It also has dental clinics included in its services. It also includes residency training for doctors as well as a full vaccination program. The hospital holds a total of 326 beds with 8 different wings/departments. As of October 2020, the hospital has around 500 doctors in employment.

Result: (72%) of the participant were females and (28%) were males, age ranged between 21 and more than 51 years, age was categorized into three groups 20–30 (73%), 31–40 (20%) and 41–50 (5%) and more than 50 (2%). Participants were distributed in 6 different specialties with Pediatrics as the most represented specialty (18%). In the level of profession most of the participant were house officer (42%) followed by registrar (32%) and the least were specialist (4%). Level of emotional exhaustion, depersonalization, and personal accomplishment are displayed in Table 3 (77%) of the sample had a high level of (EE), (54%) had a high level of (DP), and (14%) had a low level of (PA). (19%) of the residents included in this study had a high level of burnout in all three domains of the index, and (93%) had a high level in at least one of the three. The high level of each burnout component distributed according to the research different demographic

Comment [ATD1]: Incorrect age

variables and the results of the inferential statistical tests. Males had higher levels of EE (82.8%), and DP (55%), and lower levels of PA (14.9%) than females (71.6%), (54%), (12.1%) respectively.

Conclusion: In conclusion, our study showed that burn out is highly prevalent among Ibrahim Malik hospital doctors (68 %), while female doctors had a higher degree burnout than males. Among the different professional levels, the highest percentage of burnout belonged to GPs and the lowest belonged to consultant's. ER doctors suffered the most from burnout, whereas surgeons suffered the least. Several factors attributing to the burnout symptoms of these doctors have been identified.

Key words: Burnout, emotional exhaustion, depersonalization, personal accomplishment

1. Introduction

Burnout is described as exploratory and qualitatively as a state of emotional exhaustion due to excessive demands on energy, strength or resources in the workplace. It is characterized by physical symptoms such as exhaustion, fatigue, frequent headache, and gastrointestinal disorders, sleeplessness and behavioral signs include frustration, anger, suspicious attitude, omnipotence or overconfidence, cynicism and signs of depression [1]. Maslach defines "burnout as a psychological syndrome emerging as a prolonged response to chronic interpersonal stressors on the job" [2]. Burnout syndrome is characterized by three dimensions: emotional exhaustion (EE): (energy depletion, emotional and physical fatigue or depletion), depersonalization (DP): (cynicism, loss of empathy, detachment from the job or subject or responsibility), and personal accomplishment (PA): inefficacy, ineffectiveness and sense of lack of accomplishment or competence. Maslach Burnout Inventory (MBI), is the most widely used self-reported questionnaire to measure the three dimensions of burnout quantitatively [2,3]. The three dimensions of burnout are emotional exhaustion (EE), depersonalization (DP) (cynicism), and personal accomplishment (PA) (low sense of personal accomplishment). The World Health Organization (WHO) defined burnout as an occupational syndrome that results from chronic workplace stress that has not been successfully managed. It is characterized by three dimensions: feelings of energy depletion or exhaustion, increased mental distance from one's job, or feelings of negativism or cynicism related to one's job; and reduced professional efficacy. Burnout is recognized in the edition of International Classification of Disease (ICD-11- the Official Compendium of Diseases 11th edition), WHO. Burnout was not classified as a medical condition,

also recognized workplace burnout as the diagnosable condition (Diagnostic Code QD85) resulting from chronic workplace stress and encompassing a constellation of exhaustion, cynicism, and reduced efficacy [4]. Burnout appears as a global phenomenon affecting physicians all over the world, burnout affects physician, patient, and health care organization. Burnout affects physical health, mental health, and psychological health and personal life of the physicians, leading to occupational consequences [5-9]. Burnout consequences negatively affect the quality of patient healthcare, patient's safety and satisfaction, including medical errors, malpractice, sub-optimal health care services [10,11]. Burnout consequences overburden the health care organizations with a high economic cost due to high rate of physician turnover, early retirement, low productivity, paying compensation of malpractice and medical errors, patient poor satisfaction with the quality of health service [12-15].

2. Methods:

2.1. Study design and area:

This descriptive cross-sectional hospital-based study was conducted in Ibrahim Malik teaching hospital in Khartoum, Sudan.

This research was conducted in Ibrahim Malik Teaching Hospital, located in Khartoum State, Sudan. It was founded in 1977 and it is a governmental owned hospital providing many and various facilities to the general public. The hospital provides a 24/7 Emergency service, as well as many outpatient's clinics. It also has dental clinics included in its services. It also includes residency training for doctors as well as a full vaccination program. The hospital holds a total of 326 beds with 8 different wings/departments. As of October 2020, the hospital has around 500 doctors in employment.

2.2. Data collection and management:

The data was collected from 100 Physicians by a questionnaire specially prepared for the purposes of this study. Filling out the questionnaires was distributed by the research team in person. The questionnaire consisted of 27 questions, with 9 questions attributed to each of the 3 risk factors (Depersonalization, Emotional Exhaustion, and Personal Accomplishment). Data was analyzed by using SPSS program version 26 and the result was represented in tables and figures. Burnout was assessed as mild, moderate, serious or severe according to the following table [16]

Table(1):TheParameterstobeused formeasuringburnoutsymptoms

Degree	Percentage	Comment
None-Mild	0%-25%	This Physician is unlikely to be suffering from Burnout, and they are likely content with their professional and personal life.
Moderate	25.1%-50%	This Physician is likely to be suffering from a moderate degree of Burnout syndrome, and is recommended to seek out information on methods of self-care or increase engaging activities to increase mental and personal wellbeing
Serious	50.1%-75%	This physician is likely suffering from a serious degree of Burnout Syndrome and should seek assistance from a trusted mental health professional
Severe	75.1%-100%	This physician is likely suffering from a severe degree of Burnout Syndrome, and should seek immediate help from a trusted mental health professional

3. Results

The final sample size was (100), Table 2 represents the distribution of residents according to demographic and work-related variables. (72%) of the participants were females and (28%) were males, aged between 2 and more than 51 years, and age was categorized into three groups 20–30 (73%), 31–40 (20%) and 41–50 (5%) and more than 50 (2%). Participants were distributed in 6 different specialties with Pediatrics as the most represented specialty (18%). In the level of profession most of the participants were house officer (42%) followed by registrar (32%) and the least were specialist (4%).

Table 2 Distribution of the sample according to the demographic and work-related variables

Variables		Percentage	Frequency
Gender	Females	72%	72
	Males	28%	28
Age group	20–30	73%	73
	31–40	20%	20
	41–50	5%	5
	51 and more	2%	2
Level of profession	House officer	42	42 %
	Registrar	32	32 %
	Specialist	4	4 %
	Consultant	7	7 %
	GP	15	15 %
Specialty	Surgery	10 %	10
	Internal Medicine	14%	14
	Pediatrics	18 %	18
	OB/GYN	19 %	19
	ER	21%	21
	Orthopedic	5%	5

Gp= general physician, OB / GYN = obstetrics and gynecology, ER = emergency

Comment [ATD2]: In all table 2 and 3 as the frequency and percentage is same, data can be shown in one column instead of two.

Levels of emotional exhaustion, depersonalization, and personal accomplishment are displayed in Table 3 (77 %) of the sample had a high level of (EE), (54 %) had a high level of (DP), and (14 %) had a low level of (PA). (19 %) of the residents included in this study had a high level of burnout in all three domains of the index, and (93 %) had a high level in at least one of the three.

Table 3 Levels of burnout for the three dimensions of the index

	EE		DP		PA	
	Percent	Frequency	Percent	Frequency	Percent	Frequency
Low level	7%	7	16.0%	16	65%	65
Moderate	16%	16	30%	30	21%	21
High level	77%	77	54%	54	14%	14
High level of burnout in all domains of the index					19%	19
High level of burnout in at least one domain of the index					93%	93

EE = emotional exhaustion, DP = depersonalization, PA = personal accomplishment

Table 4 represents the percentage of the high level of each burnout component distributed according to the research different demographic variables and the results of the inferential statistical tests. Males had higher levels of EE (82.8%), and DP (55%), and lower levels of PA (14.9%) than females (71.6%), (54%), (12.1%) respectively. Significant relation was only found between gender and level of EE using T-test ($p=0.004$).

Table 4 Research variables and statistical test

Research variables		Burnout variables			Statistical tests			
Variable		Percent EE	High level of fDP	High level of PA	Statistical test	P value		
Gender	Females 44.10%	71.6%	54%	12.1%	t-test	0.004 for EE		
	Males 55.90%	82.8%	55%	14.9%				
Age group	21–25	34%	73.6%	56.1%	ANOVA	0.037 for EE		
	26–30	59.50%	79.3%	53.2%				
Specialty	31–35	6.50%	86.3%	59%	ANOVA	0.066 no significant relation		
	General Surgery	32.6%	81.4%	58%				
	Internal Medicine	76.2%	54.5%	14.6%				
	Pediatric	24.2%	9.2%	80%			45%	15%
	Obstetrics	5.7%	76.9%	60%			11.5%	
Orthopedic	5.7%	64.6%	45%	5.8%				
Emergency Medicine	89%	34%	12.3%					

4. Discussion:

The findings of this study indicate that the collapse pattern has an intimidating high frequency in the population it was made on, which could indicate the need for farther assessments and interventions for the 900 registered croakers in Sudan [17]. This supported utmost of the former studies made in Africa [18] [19] [20]. Contradicting the “Burnout in Australasian Youngish Fellows” study [21] and “Physician well-being frequency of collapse and associated threat factors in a tertiary sanitarium, Riyadh, Saudi Arabia” exploration results, surgeons in Ibrahim Malik Teaching Hospital had the alternate smallest probable threat of collapse, which, still, was only fairly lower than another thing. Collapse within the Internal Medicine department, had- fairly- the smallest frequencies, which was against the “Physician well-being frequencies of

collapse and associated threat factors in a tertiary sanitarium, Riyadh, Saudi Arabia” exploration results [21]. Despite the diversity of the relative differences between fortes, this difference may not exactly contradict the findings of those studies, due to the overall high frequency of probable collapse among all fortes, with the frequency ranging from 51 to 81. Slightly analogous to the “Collapse among gynecological resides in Lahore, Pakistan A cross-sectionalcheck”, the frequency of collapse in gynecologists was nearly 60 [22]. Nonetheless, this attendant chance included all specialty situations of Obstetrics and Gynecology; hence the comparison to that study could be fairly invalid since it only included Gynecological resides. Alarmingly, the most affected croakers.Were those at the early stage of their medical career. Supporting the “Physician well- being frequency of collapse and associated threat factors in a tertiary sanitarium, Riyadh, Saudi Arabia” exploration, the frequency was advanced in resides than advisers [22]. The results also supported the analogous pattern of advanced collapse in inferior croakers. in opposition to a dropped threat with high specialization situations shown in the “Collapse in anesthesiologyand ferocious care is there a problem in Germany?” Exploration [23]. Both could be linked to the allegedly dropped workload and hours of advisers in discrepancy to the prolonged GPs’and resides’ workload and work hours. Different to the “Collapse and sources of stress among medical resides at Hammad Medical Corporation, Qatar” exploration, women scored worse in depersonalization [24] unlike their counterparts which scored much better [24]. A slight difference from some of the former studies made in Africa, in which the results showed that depersonalization was fairly more affected than emotional prostration when comparing situations. particular accomplishment was the least affected factor in opposition to the result of one of the former studies [25], which could be due to the developing nature of the country with, an allegedly, increased number of people with low socioeconomic status presenting to governmental sanitarium to seek medical help due to lower prices), which could increase the feeling of accomplishment in croakers. analogous to The 2013 Medscape lifestyle Report (which was grounded on the check of over,000 croakers .in the US), “Burnout in Australasian YoungishFellows” study, and “ Gender differences in the effect of grief responses and collapse on emotional torture among clinical oncologists” exploration, collapse in womanishcroakers was advanced than their manly associates [26] [27], with all womanish actors likely suffering from collapse in comparison to 98 of manly actors. This could be linked to the hormonal changes women suffer during ovulation, period, gestation and menopause or indeed due to the increased

artistic burden regarding women's liabilities in the Sudanese society, hence, pouring to the increased threat of collapse. still, this is a threat that wasn't studied in this exploration.

5. Conclusion:

In conclusion, our study showed that burn out is highly prevalent among Ibrahim Malik hospitaldoctors (68 %), while female doctors had a higher degree burnout than males. Among the differentprofessional levels, the highest percentage of burnout belonged to GPs and the lowest belonged toconsultant's.ERdoctors sufferedthemostfrom burnout,whereassurgeonssufferedtheleast. Several factors attributing to the burnout symptoms of these doctors have been identified. The firstfactor being personal accomplishment, which was the most affected factor in general practitionersand the least affected in consultants. Emotional exhaustion was the highest in specialist and the leastinhouseofficers. Depersonalizationwasthehighestinconsultantsand theleastinspecialists. In terms of the riskfactors accrediting toburnout in terms of gender,our results concluded thatmales gota higherpercentage of lowpersonal accomplishmentandemotional exhaustion thanfemales. Meanwhile, femalesscored a higherpercentageofdepersonalizationthan males. In regards to the risk factors affecting the different specialties, results have shown that emotionalexhaustion was the most affected in ICU doctors and least affected in orthopedic doctors.

LowpersonalaccomplishmentwasthehighestinorthopedicsandthelowestinOB/GYN. Depersonalization,wasthe highestinOB/GYN andthelowestinICU. Finally, the prevalence of symptoms of Burnout Syndrome among doctors employed at IbrahimMalik TeachingHospitalinOctober2020wasfoundtobeveryhigh, constitutingapressingchallenge thatneedstobemetbyorganizations, individualsandsocietyatlarge.

Reference

1. Freudenberger HJ. "Staff burnout". *Journal of Social Issues* 30 (1974): 159-165.
2. Maslach C, Jackson SE. The measurement of experienced burnout. *Journal of organizational behavior*. 1981 Apr;2(2):99-113.
3. Maslach C and Leiter MV. "Understanding the burnout experience: Recent research and its implications for psychiatry". *World Psychiatry*15 (2016): 103-111.
4. World Health Organization. Burn-out an "occupational phenomenon": International Classification of Diseases. Geneva: World HealthOrganization (2019).

5. Salvagioni DA, Melanda FN, Mesas AE, González AD, Gabani FL, Andrade SM. Physical, psychological and occupational consequences of job burnout: A systematic review of prospective studies. *PloS one*. 2017 Oct 4;12(10):e0185781.
6. Garcia C de L., *et al.* "Influence of Burnout on Patient Safety: Systematic Review and Meta-Analysis". *Medicina* 55.9 (2019): 553.
7. Udemezue O Imo. "Burnout and psychiatric morbidity among doctors in the UK: a systematic literature review of prevalence and associated factors". *BJ Psych Bulletin* 41 (2017): 197-204.
8. Patel RS., *et al.* "Factors related to physician burnout and its consequence: A review". *Behavioral Sciences* 8 (2018): 98.
9. Shanafelt TD, West CP, Sinsky C, Trockel M, Tutty M, Satele DV, Carlasare LE, Dyrbye LN. Changes in burnout and satisfaction with work-life integration in physicians and the general US working population between 2011 and 2017. In *Mayo Clinic Proceedings* 2019 Sep 1 (Vol. 94, No. 9, pp. 1681-1694). Elsevier.
10. Panagioti M, Geraghty K, Johnson J, Zhou A, Panagopoulou E, Chew-Graham C, Peters D, Hodkinson A, Riley R, Esmail A. Association between physician burnout and patient safety, professionalism, and patient satisfaction: a systematic review and meta-analysis. *JAMA internal medicine*. 2018 Oct 1;178(10):1317-31.
11. Tawfik DS, Profit J, Morgenthaler TI, Satele DV, Sinsky CA, Dyrbye LN, Tutty MA, West CP, Shanafelt TD. Physician burnout, well-being, and work unit safety grades in relationship to reported medical errors. In *Mayo Clinic Proceedings* 2018 Nov 1 (Vol. 93, No. 11, pp. 1571-1580). Elsevier.
12. Dewa CS, Loong D, Bonato S, Trojanowski L, Rea M. The relationship between resident burnout and safety-related and acceptability-related quality of healthcare: a systematic literature review. *BMC medical education*. 2017 Dec;17(1):1-6.
13. Tawfik DS, Scheid A, Profit J, Shanafelt T, Trockel M, Adair KC, Sexton JB, Ioannidis JP. Evidence relating health care provider burnout and quality of care: a systematic review and meta-analysis. *Annals of internal medicine*. 2019 Oct 15;171(8):555-67.
14. Han S, Shanafelt TD, Sinsky CA, Awad KM, Dyrbye LN, Fiscus LC, Trockel M, Goh J. Estimating the attributable cost of physician burnout in the United States. *Annals of internal medicine*. 2019 Jun 4;170(11):784-90.

15. Owoc J, Mańczak M, Jabłońska M, Tombarkiewicz M, Olszewski R. Association between physician burnout and self-reported errors: meta-analysis. *Journal of Patient Safety*. 2022 Jan 1;18(1):e180-8.
16. Panagioti M, Geraghty K, Johnson J, Zhou A, Panagopoulou E, Chew-Graham C, Peters D, Hodkinson A, Riley R, Esmail A. Association between physician burnout and patient safety, professionalism, and patient satisfaction: a systematic review and meta-analysis. *JAMA internal medicine*. 2018 Oct 1;178(10):1317-31.
17. MAM I & Mahgoub M A. Physician Burnout Syndrome. *Sudan JMS Vol. 4, No. 4, Dec2009:327-331*
18. Abdo SA, El-Sallamy RM, El-Sherbiny AA, Kabbash IA. Burnout among physicians and nursing staff working in the emergency hospital of Tanta University, Egypt. *East Mediterr Health J*. 2016 Mar 15;21(12):906-15.
19. Liebenberg A, Coetzee J, Conradie H, Coetzee J. Burnout among rural hospital doctors in the Western Cape: comparison with previous South African studies. *Afr J Prim Health Care Fam Med*. 2018;10(1):e1–e7.
20. Ogundipe OA, Olagunju AT, Lasebikan VO, Coker AO. Burnout among doctors in residency training in a tertiary hospital. *Asian J Psychiatr*. 2014;10:27–32.
21. Aldrees TM, Aleissa S, Zamakhshary M, Badri M, Sadat-Ali M. Physician well-being: prevalence of burnout and associated risk factors in a tertiary hospital, Riyadh, Saudi Arabia. *Annual Saudi Medicine*. 2013 Sep-Oct;33(5):451-6.
22. Opoku ST, Apenteng BA. Career satisfaction and burnout among Ghanaian physicians. *Int Health*. 2014;6(1):54–61.
23. Rajan S, Engelbrecht A. A cross-sectional survey of burnout amongst doctors in a cohort of public sector emergency centres in Gauteng, South Africa. *Afr J Emerg Med*. 2018;8(3):95–99.
24. van der Walt N, Scribante J, Perrie H. Burnout among anaesthetists in South Africa. *S Afr J Anaesth Analg*. 2015;21(6):169–172.
25. van der Walt N, Scribante J, Perrie H. Burnout among anaesthetists in South Africa. *S Afr J Anaesth Analg*. 2015;21(6):169–172.
26. Stodel JM, Stewart-Smith A. The influence of burnout on skills retention of junior

Comment [ATD3]: Follow the format of bibliography

doctors

atRedCrossWarMemorialChildren'sHospital:acasestudy.SAfrMedJ.2011Feb;101(2):11
5-8

27. Kim MH, Mazenga AC, Simon K, Yu X, Ahmed S, Nyasulu P, Kazembe PN, Ngoma S, Abrams EJ. Burnout and self-reported suboptimal patient care amongst health care workers providing HIV care in Malawi. PLoS One. 2018 Feb 21; 13(2):e0192983.

Comment [ATD4]: Follow any one bibliography format. Either Vancouver/ APA/Oxford. Or format given by the journal

UNDER PEER REVIEW