

Incidence and management of cystic calculi in dogs in and around Meerut

Abstract:

A total of 54 animals of either sex with cystic calculi were prospectively studied. The Mmost commonly affected dogs were ~~often the~~ mature adult dogs aged ~~between~~ 3 to 10 years, with ~~were affected and the~~ juveniles (<1 year) faring were less well commonly affected. The Pprevalence was highest during ~~maximal in~~ extreme winter and summer seasons. Cystic calculi were diagnosed either by radiography or ultrasonography. All the dogs were given urinary alkalizer and calculolytic drugs to dissolve the calculi, and the dogs with urinary obstruction were treated surgically by cystostomy. The Mmajority of the cases showed uneventful recovery.

Keywords: Cystic calculi, cystostomy, calculolytic drugs, dogs, incidence, Meerut.

Introduction:

Cystic calculi (urinary bladder stones) are a common condition responsible for the retention and obstruction of urine in dogs (Dvorska *et al.*, 2016). The formation of cystic calculi is generally associated with the precipitation and crystal formation of a variety of minerals (Stiller *et al.*, 2014), though it is a multifactorial disease (Ackerman, 2016). The sequence of events that triggers stone formation is not fully understood. In association with highly concentrated urine, less water and a high dietary intake of minerals and protein may contribute to increased saturation of salts in the urine ~~Less water and high dietary intake of minerals and protein in association with highly concentrated urine may contribute to increased saturation of salts in the~~ urine (Parivar *et al.*, 1996). Disease conditions such as bacterial infections in the urinary tract can also increase urine salt concentration (Weese *et al.*, 2021). The understanding of incidence is

important for the treatment and prevention of urinary stones in dogs. Surgery is the primary treatment of obstructive urolithiasis (Larson, 1996). ~~Surgical procedure of c~~Cystostomy together with medical dissolution of calculi is considered an effective technique for resolution of calculi in small animals (Ewoldt *et al.*, 2006). Animals with prolonged obstruction have high morbidity due to subsequent uremia. Surgical management of such patients should be done very cautiously. In this study, ~~the~~ incidence of cystic calculi, their diagnosis, and surgical and medical management in 54 cases was reported.

Material and Methods:

All the cases were treated and examined as per standard treatment and examination protocols, so ~~the~~ ethical approval was not necessary for this study. A total of 54 dogs of either sex with complaints of dribbling of urine, urinary retention, and urinary obstruction were prospectively studied (**Table.1**). These cases were presented ~~to Veterinary Clinical Complex, College of Veterinary and Animal Sciences, SVPUA&T, from~~ Meerut ~~from and~~ the adjoining areas between the years 2018-2021. ~~A P~~physical examination was done to check the status of the urinary bladder. Further, radiography or ultrasonography was performed to establish the diagnosis (**Fig.1 A and B**). All the animals with urinary obstruction were treated surgically by cystotomy technique, and animals with dribbling of urine were treated ~~by with~~ urinary alkalizers and calculolytic drugs.

After the confirmation of calculi in ~~the~~ urinary bladder, all the animals were aseptically prepared for surgery and administered general anesthesia. The surgical interventions were done by keeping ~~the~~ dogs in dorsal recumbency. ~~An I~~ntravenous infusion of normal saline was maintained during surgery. An ~~approximately about~~ 5-6 c.m. long mid-line (in females) and right paramedian skin incision lateral to penis was given (in males). ~~The P~~penis, along with ~~the~~ incised

skin, was retracted to the left lateral side till the midline was visualized, followed by a routine laparotomy in the caudal abdomen. The distended urinary bladder was located and exteriorized (Fig. 2A). The urine in the bladder was removed by suction through a sterile syringe. Then an incision was ~~given~~ made on the dorsal aspect of the urinary bladder to reach the lumen. The lumen was searched for calculi/concretions and then the calculi were removed (Fig. 2 B) and the lumen of the urinary bladder was flushed with normal saline. A sterile polyethylene catheter was passed from the urinary bladder through the urethra to check the urethral patency. If, the urethra was found obstructed, a urethrotomy was also performed to remove concretions/calculi. A sterile polyethylene catheter (infant feeding tube) was passed from the external urethral orifice to the urinary bladder. The external end of the catheter was kept fixed with to the preputial skin by stay sutures. Cystotomy, urethrotomy, and laparotomy wounds were closed, dressed, and protected in a routine manner.

Postoperatively, an amoxicillin-sulbactam antibiotic combination (15 mg/kg, IM) for 5 days, analgesic meloxicam (0.2 mg/kg, IM) for 3-5 days, and antacid pantoprazole (1 mg/kg, PO) for 5 days were administered. The urethral catheter was removed on 8-12 days and skin sutures were removed after healing of the wounds, generally on the 12th day.

In animals with dribbling of urine and diagnosed with very fine concretions in the urinary bladder, a sterile polyethylene catheter (infant feeding tube) was passed to the urinary bladder from the urethra to dislodge and push back any concretions. The catheter was secured to the preputial skin and urinary flow was maintained. After this maneuver, all the affected animals were given urinary alkalizer disodium hydrogen citrate (Syp. Alkasol) and the calculolytic drug Tab. Cystone till the cystic concretions were not seen ultrasonographically, and the catheter was removed accordingly.

Result and Discussion:

Cases of cystic calculi were more prevalent in the extreme winter and summer. Mature adult dogs (3-10 years) (48%) were most affected, followed by young adult dogs (1-3 years) 28% and juvenile dogs (<1 year) 11% were least affected, followed by geriatric dogs (>10 years) 13% (Fig. 3). Anesthesia, cystotomy, and urethrotomy procedures, along with catheterization of the urinary bladder and fixation of the external end of the catheter with preputial skin, were achieved without difficulties. The majority of the dogs showed uneventful recovery except for a few minor incidences of infection and pus formation at surgical sites.

Cystic calculi are a common problem in dogs (Ling *et al.*, 1998) and are seen mostly in extremely cold-winter and hot climates-summer (Singh *et al.*, 2011). This may be associated with less water intake and more water losses during winter and summer, respectively. Diagnosis of cystic calculi can be made either by radiography or ultrasonography. Sometimes, the diagnosis of the cystic calculi by radiography becomes difficult, if they are radiolucent (Larson, 2009) or very fine. By ultrasonography, radiolucent and very fine calculi/concretions can be diagnosed easily as hyperechoic structures showing acoustic shadowing below them. Acoustic shadowing distal to floating calculi/concretions was not seen. Similar findings were also reported by Verma *et al.* (2006) and Dehmiwal *et al.* (2016). Surgery is the primary treatment for obstructive cystic calculi, except for very fine calculi/concretions which can be managed by urinary alkalizers and calculolytic drugs. Nutritional management may reduce the occurrence of calculi formation significantly in the dogs.

Conclusion:

In this study, the incidence of cystic calculi in dogs in and around Meerut was recorded. Such cases were managed effectively by surgery and medicinal treatment ~~effectively~~ after diagnosis~~ing~~ either by radiography or ultrasonography. Further, elaborative research is required on nutritional management to prevent the occurrence of ~~the~~ urolithiasis in dogs.

References:

- Ackerman, N., (2016). Nutritional management of canine urolithiasis. *The Veterinary Nurse*, 7(1), pp.12-17.
- Dehmiwal, D., Behl, S.M., Singh, P., Tayal, R., Pal, M., Saharan, S. and Chandolia, R.K., (2016). Diagnosis and surgical management of cystolith in dogs. *Haryana Vet*, 55(1), pp.103-105.
- Dvorska, J. and Saganuwan, S.A., 2015. A review on urolithiasis in dogs and cats. *Bulgarian Journal of Veterinary Medicine*, 18(1).
- Ewoldt JM, Anderson DE, Miesner MD, Saville WJ (2006). Short- and long-term outcome and factors predicting survival after surgical tube cystostomy for treatment of obstructive urolithiasis in small ruminants. *Veterinary Surgery*, 35: 417-422
- Larson BL (1996). Identifying, treating, and preventing bovine urolithiasis. *Veterinary Medicine* 91, 366-377.
- Larson, M.M. (2009). The Kidneys and Ureters. In: BSAVA Manual of Canine and Feline Abdominal Imaging. O'Brien, R. and Barr, F. (eds.). British Small Animal Veterinary Association, Gloucester.

- Ling, G.V., Ruby, A.L., Johnson, D.L., Thurmond, M. and Franti, C.E. (1998). Renal calculi in dogs and cats: prevalence, mineral type, breed, age and gender interrelationships (1981-93). *Internal Med.* 12: 11-21.
- Parivar, F., Low, R.K. and Stoller, M.L., 1996. The influence of diet on urinary stone disease. *The Journal of urology*, 155(2), pp.432-440.
- Stiller, A.T., Lulich, J.P. and Furrow, E., 2014. Urethral plugs in dogs. *Journal of veterinary internal medicine*, 28(2), pp.324-330.
- Singh, T., Kinjavdekar, P., Aithal, H.P., Pawde, A.M., Pratap, K. and Mukherjee, R., (2011). Obstructive urolithiasis in domestic animals: A study on pattern of occurrence and etiology.
- Verma, P., Mohindroo, J., Singh, S.S. and Singh, C.B. (2006). Sonographic findings in affections of urinary system in dogs. *Indian J. Vet. Surg.* 27(2): 104-107.
- Weese, J.S., Blondeau, J., Boothe, D., Guardabassi, L.G., Gumley, N., Papich, M., Jesseni, L.R., Lappin, M., Rankin, S., Westropp, J.L. and Sykes, J. (2021). International Society for Companion Animal Infectious Diseases (ISCAID) guidelines for the diagnosis and management of bacterial urinary tract infections in dogs and cats. *Journal of Japanese Association of Veterinary Nephrology and Urology*, 13(1), pp.46-63.

Table: 1. Incidence of cystic calculi as per the age groups.

S.No.	Age group	Group code	Number of cases
1.	Juvenile (<1 year)	J	06
2.	Young adult (1-3 years)	Y	15
3.	Mature adult (3-10 years)	M	26
4.	Geriatric (>10 years)	G	07
Total			54

Legends of the figures:



Figure: 1. Radiographic (A) and ultrasonographic (B) diagnosis of cystic calculi in dogs.




Figure: 2. Intra-operative view of exteriorization of urinary bladder (A) and removed calculi (B).

Figure: 3. Incidence of cystic calculi in different age groups.

UNDER REVIEW