

PERCEPTIONS AND EXPERIENCES OF PREGNANT WOMEN ABOUT ROUTINE HIV COUNSELLING AND TESTING IN TERTIARY HOSPITALS IN RIVERS STATE

ABSTRACT

The study is aimed at investigating Perceptions and Experiences of Pregnant Women about Routine HIV Counselling and Testing in Tertiary Hospitals in Rivers State. Qualitative and quantitative research designs were used for the study. Three objectives and three research questions guided the study. A sample size of 200 pregnant women was used for the quantitative study while depth interviews were held with 16 pregnant women who tested for HIV at antenatal clinics (ANC), as well as four (4) healthcare workers involved in routine HIV counselling and testing. The instrument was subjected to a reliability test and a reliability coefficient of 0.85 was obtained. Data were analysed using the content analysis approach and simple percentages. The qualitative section of the study was divided into attitude of husbands, attitude of healthcare workers, fear of death for being HIV positive and suicidal tendency subthemes. The findings of the study revealed that 5% of the respondents were below between 18 - 25years, 55% were between 26-30 years, 35% were between 31-36years while 5% were 37years and above. The result further revealed that HIV counselling and testing is well received and acceptable for most pregnant women. Most women were aware of routine antenatal HIV testing before registering for ANC services. The women indicated that they had learnt about HIV testing and its benefits through various community education activities, mass media and during their previous pregnancies. During ANC services, most health care workers were over stressed by the large numbers of women waiting for HIV testing. This was noticed to have downplayed or reduced the quality of counselling services, thereby affecting the quality of PMTCT services. Also, routine HIV counselling and testing in tertiary hospitals in River State is very important and necessary but should not be made compulsory. The study concluded that routine counselling and testing will enable individuals to make informed choices regarding the HIV test and take appropriate action. Therefore, the study recommended that routine counselling and testing facilities should be made available in all antenatal units of the hospitals.

Keywords: Perceptions, Experiences, Pregnant Women, Routine HIV Counselling.

1.0 INTRODUCTION

Over the years, human immunodeficiency virus (HIV) has continued to be a global public health problem with too much burden on women and children (Source). This virus has destroyed many innocent lives and exposed them to poor quality of life. According to (UNAIDS 2020; Alemu et al. Ambaw & Wilder Smith, 2017), there are many ways people

Comment [TM1]: Write in full.

Comment [TM2]: Is it below or between?

Comment [TM3]: Write in full.

can be infected with HIV, but the one that is paramount and remains a public health challenge, especially in developing countries, remains the mother-to-child transmission. According to the World Health Organization (2010). This virus can be transmitted from an infected pregnant mother to her child during pregnancy, childbirth and breastfeeding. Without preventive interventions, the possibility of a baby getting HIV from an infected mother is in the range between 15% and 45%, which in most cases occurs during pregnancy, childbirth or breastfeeding (Federal Ministry of Health Ethiopia 2015). In Asia, the percentage of HIV prevalence rate among pregnant women is between 1 to 2 (Sinha et al., 2008). Likewise, in India, the percentage of women that have done HIV testing is about 96 (Pai et al., 2006), also in Vietnam, the percentage of HIV prevalence among Pregnant women are reported to be 0.37 (UNGASS 2008). Pregnant women infected with the virus (HIV) are on the increase in sub-Saharan Africa with prevalence ranging from 5 to 37% (Nuwagaba-Biribonwoha, 2017) of mother-to-child transmission (MTCT) which must be treated through routine HIV counselling and tests (HCT).

Globally, the percentages of all HIV/AIDS cases are more of young people who are between the ages of 10 –24 years over 50. Irrespective of high vulnerability to HIV infection, VCT uptake by young people is significantly very lower (WHO 2020). According to a survey from sub-Saharan Africa (2005–2010), only 10% of males and 15% female of 15–24 years of age knew their HIV status, this means that the majority of young people in this age group are undiagnosed leading to HIV epidemic, thereby exposing them to high risk of either acquiring or transmitting the disease (WHO 2018).

In Tanzania, irrespective of the fact that awareness of VCT in facilities is high, 65.8% of males and 46.3% of females of 15–24 years are still not aware of their HIV status and unfortunately, this is where the HIV epidemic is concentrated and very high (National Bureau of Statistics 2019). In Arusha, 69.8% of males and 30.3% of females of 15–24 years had never been tested (National Bureau of Statistics 2019). Only 50% of females and 39% of males in secondary or higher institutions in Tanzania know their HIV status, giving an average of 45% VCT uptake among them (WHO 2018). The low response to VCT among young people is said to be associated with different factors, which range from fear of the unknown and knowing their HIV status to the limiting factors towards the service and social issues influencing the attitudes and behaviours of the service providers (National Bureau of Statistics 2019). However, an approximately 50% increase in HIV-related deaths among adolescents between 2005 and 2012 was found to be contributed to inadequate friendly VCT

services, poor prioritization of adolescent issues, inadequate treatment, and lack of support for young people.

Also, according to the global summary report of 2019 figure, "31.6–44.6 million people were living with HIV and 1.2–2.2 million people acquired HIV infection. More than half of the people newly infected with HIV live in Sub-Saharan Africa" World Health Organization (2021). In a very close look into a decade's trend and monitoring the progress from 2010 to 2019, one would see that the HIV/AIDS-acquired immunodeficiency syndrome epidemic is on the increase in Eastern Europe and Central Asia, with the number of people acquiring HIV increasing by more than 70% [\(Source\)](#). Similarly, the decade quantitative assessment reveals an increase of more than 20% in North Africa, Latin America, and the Middle East World Health Organization (2021).

UN General Assembly agrees that ending AIDS as a public health threat by 2030 requires 4 milestones by 2020 and urgent action [agreed](#). These milestones include reducing the number of new HIV infections to less than 500,000 annually worldwide and reducing the number of deaths from AIDS-related causes to less than 500,000 annually worldwide, This includes eliminating HIV-related stigma and discrimination. Although the number of deaths from AIDS-related causes has fallen steadily by nearly a third and annual incidence rates are at their lowest since 1989, these global goals have not been met and [it](#) remains a big challenge, United Nations(2021).

The epidemiological rationale for HIV programs focuses on key populations. These key populations continue to bear a significant HIV burden and influence the dynamics of the HIV epidemic. In many African countries, certain groups are particularly vulnerable to HIV infection, including migrant workers, refugees, truck drivers, military personnel, miners, children, [adolescents](#), adolescents, and young women. These populations are not uniformly at risk or equally affected in different countries and epidemic settings, WHO(2016).

Established by United Nations Member States, Universal Health Coverage (UHC) aims to address the determinants of infectious diseases and their comorbidities, including HIV/AIDS, through multisectoral action. World Health Organization (2020). The Sustainable Development Goals (SDGs) are set for 2030 and include the empowerment of women and girls, including access to quality essential health services. In September 2019, UN Member States reaffirmed their commitment to this goal at the UN General Assembly High-Level Meeting on UHC. They also set a new goal of 1 billion more people having access to quality

essential health services by 2023- Zhenmin (2020). But today, nearly half of the people living with HIV/AIDS are women.

Despite high awareness and knowledge of VCT services, the response to HIV testing by young people has been reported to be low (National Bureau of Statistics 2019). This study aimed at determining the factors that influence young people to uptake VCT to influence early HIV detection to avoid increased AIDS cases and risk behaviors and influence support and care to HIV victims. Furthermore, facilitate the designing of appropriate strategies by the government and policymakers geared toward increasing VCT uptake among young people. Since young people are an important force for development, increasing VCT uptake would benefit not only the young people themselves but also determine the health of the future generation and the sustainable economic development of the nation.

HIV/AIDS counselling gives individuals psychological support or emotional treatment for preventing the transmission of the virus, controlling its spread and arresting the impact of the virus (Idogho, 2010). The testing involves a blood sample of the individual to know his/her status. HIV/AIDS counselling and testing help those that are negative and also, enable sick people to understand their body system and plan possible ways to manage life challenges and at large, support the growth of society (UNAIDS 2018). HIV counselling and testing (HCT) assume an important role in the treatment and prevention of HIV/AIDS, antiretroviral drugs are also very effective and are used for treatment, and HIV-related ailments such as tuberculosis (TB) control and psychological support (Idoho, 2010). Counselling in HIV/AIDS is in three phases, which are; pre-test counselling, post-test counselling and follow-up sessions with adherence counsellors (Yahaya et al., 2010). There are two (2) basic types of HIV testing which include the “opt-in” or voluntary counselling and testing VCT; and “opt-out” or routine HIV counselling and testing. There has been constantly increasing international support for "routine" or "opt-out" testing, where all patients are counselled and tested unless they decline. Routine testing occurs in selected healthcare settings, such as pre-natal clinics or medical wards with a high prevalence of HIV, Tuberculosis, or sexually transmitted infections. This approach has successfully identified many patients in need of treatment and care (WHO 2010). Routine HCT during pregnancy is a starting point for instituting a prevention program. This strategy promotes adequate treatment for HIV-positive women and has a positive impact on the child HIV transmission rate. For HIV-negative women, it provides opportunities for education and behavioural changes (De Cock et al., 2009). But experience to date in many countries shows great variation in willingness to make

use of the services that are available and very low acceptance of routine HTC services (UNAIDS, 2010). Given the importance of this HCT, the utilisation of routine voluntary counselling and testing (VCT) especially among pregnant women in Nigeria is still poor (Olajide et al., 2015). It, therefore, becomes important in this present study to look at the perception and experiences of pregnant women in their utilisation of routine HIV counselling and testing.

Perception is a concept that involves the dynamic psychological process responsible for attending to, organizing and interpreting sensory data. Perception is the process of recognizing, organizing and interpreting sensory information, it deals with the human senses and generates signals from the environment through the five sense organs: sight, hearing, touch, smell and taste (Human Perception and Information Processing, 2015). Pregnant mothers who have a positive perception toward Mother to Child Transmission of HIV (MTCT) can assess their susceptibility correctly; understand the severity of MTCT of HIV, and the benefit of testing (Jones et al 2015). A situation where pregnant women have a poor perception towards MTCT could constitute a significant barrier to the uptake of HIV routine testing, as mothers may not see the need for testing or inaccurately assess the risk associated with non-engagement in HIV testing and counselling. It is assumed that the perception that pregnant women hold may go a long way to determining whether they will engage in routine HIV testing and counselling or not.

Experiences of pregnant women may play a role in routine HIV testing and counselling. Pregnant women face various experiences in a bid to get medical attention and these health experiences are also similar when trying to be tested for HIV. These experiences sometimes are so frustrating and hence discourage many pregnant women to be tested for HIV. Some of the reasons why pregnant women do not test for HIV include the inaccessibility of healthcare facilities (Gunn et al 2016) perceived lack of confidence, stigma, and discrimination (Mumtaz et al 2013), also the cost of treatment, as well as illiteracy and inability to secure husband's permission. Others include attitude, skills of health workers and inadequate resources (Kwapong 2014). All these experiences may also have serious negative implications for pregnant women not wanting to go for routine HIV counselling and testing. Based on this premise the study examines the perception and experiences of pregnant women about routine HIV counselling and testing in tertiary hospitals in River State.

1.1 Aim and Objective of the Study

The aim of this study is to examine the perceptions and experiences of pregnant women about routine HIV counselling and testing in tertiary hospitals. The specific objectives of this study are to;

1. Ascertain the perception of pregnant women about routine HIV counselling and testing in tertiary hospitals in River State.
2. Explore the experiences of pregnant women about routine HIV counselling and testing in tertiary hospitals.

1.2 Hypothesis

The following hypothesis was formulated for the study;

Ho1: Educational qualification does not significantly determine the perception of pregnant women about Routine HIV counselling and testing in tertiary hospitals in River State.

2.0 METHODOLOGY

A mixed methods study design comprising of quantitative and qualitative data collection was employed. The study population consist of 500 pregnant women who are attending antenatal in two (2) tertiary hospitals in Rivers states. The sample size of 220 women attending antenatal were selected from the total population of 500. Simple random sampling techniques was used to select the sample. A total of 10 pregnant women and 10 healthcare workers working in the ante-natal unit were be interviewed. The study made use of both descriptive and inferential statistic in analysing the data collected. ~~The study made use of both descriptive and inferential statistics in analysing the data collected.~~ Inferential statistics helped to provide explanations for a condition or event. Descriptive statistics that merely summarize the data that has actually been measured are fundamentally different from it in that they allowed for drawing conclusions based on extrapolations (Mishra, et al. 2020; Siedlecki, 2020).

Formatted: Indent: Left: 0", First line: 0"

The study also made use of thematic analysis in handling data transcribed from the recorded interview sections during the data collection process. Tables and graphs were used to further illustrate the data collected through questionnaire. While, categorizing enormous data sets into codes, subthemes and themes, thematic analysis gave one a great deal of flexibility in how one interpreted the data and made it easier to work with large data sets (Esfehani and Walters, 2018), hence the rationale for adopting thematic analysis. Also saturation of the data reached when there was no new information from the participants. However, SPSS analytical tools were used to analyse the data collected through questionnaire among pregnant women in tertiary hospitals in River State.

3.0 RESULT AND DISCUSSION

RESULTS

Characteristics of study participants

The participants' age ranges from 15 to 37 years and above and the majority of them had completed primary education. Also, most of the women were multiparas and housewives. All of them were married and living with a partner.

4.1.2 Socio-Demographic Characteristics




Fig. 1: Age of respondents

Fig.1 shows that 5% of the respondents were below 18 - 25years, 55% were between 26-30 years, 35% are between 31-36years while 5% were 37years and above.




Fig. 2: Educational Qualifications of Respondents

Fig 2 Showed the findings of the result which revealed that 10% of the respondents had no formal education, 35% had primary education, 45% had secondary education while 10% had tertiary education.




Fig. 3 revealed that 30% of the respondents had 1 child, 60% had 2 -4 children, while 10% had 5 – 7 children.

Research Question 1: What are the perceptions of pregnant women about Routine HIV counselling and testing in tertiary hospitals in River State?

Table 1. Perception of pregnant women about Routine HIV counselling and testing in tertiary hospitals in River State.

Comment [TM4]: From what I see here, likert scale was used but never mentioned anywhere in the study

| S/N | Items | SA | S | D | SD |
|-----|-------------------------------------------------------------|---------|---------|---------|----------|
| | HIV testing and counselling is scaring | (62.9%) | (26.9%) | (5.7%) | (4.6%) |
| | HIV testing is a waste of resources | (59.2%) | (27.9%) | (8.9%) | (4.0%) |
| | there is a negative feeling against routine HIV testing | (51.4%) | (29.1%) | (12.6%) | (6.9%) |
| | HIV testing is important during antenatal | (55.4%) | (21.7%) | (16.6%) | (6.3%) |
| | Routine HIV testing often brings lack of trust in the home. | (64%) | (9.1%) | (16%) | 10.85714 |
| | the advantages of routine HIV testing outweighs its | (62.9%) | (26.9%) | (5.7%) | (4.6%) |

disadvantage

Routine HIV testing aid early detection and prevention plan against the virus. (51.4%) (29.1%) (12.6%) (6.9%)

Routine HIV testing also facilitate sexual health improvement and that of the baby as well (73.7%) (13.7%) (12%) (0.6%)

Table 1 revealed that 62.9% of the respondents strongly agreed that they are always afraid of HIV testing and counselling, 59.2% revealed that routine HIV testing as a waste of resources, 51.4% strongly agreed that they often have a negative feeling against routine HIV testing, 55.4% of the respondents strongly agreed that HIV testing is important during antenatal care, while, 64% strongly agreed that routine HIV testing often brings about lack of trust in the home. 62.9% strongly agreed that the advantages of routine HIV testing outweighs its disadvantages, 51.4% of the respondents believed that routine HIV testing aids early detection and prevention plans against the virus while 73.7% strongly agreed that routine HIV testing also facilitates sexual health improvement and that of the baby as well.

Comment [TM5]: These results need to be discussed and compared to previous studies

Interview responses

HIV testing during pregnancy is beneficial

Throughout the interview sessions, the majority of the women expressed the perception of HIV testing during pregnancy. Two major perceptions of routine HIV testing that were brought up continuously during the interviews were 'knowing one's status and protection of unborn babies from HIV infection'. Most women described that knowing a pregnant woman's HIV status is essential to protect her unborn baby as there is access to treatment.

It is very good to get blood tested for HIV. It is good to know your status. A woman will take care of her child if she knows her HIV status. (Participant 1).

Participants reported that they knew the benefits of HTC during pregnancy from different sources including community education and mass media.

They taught us during antenatal and I also heard from a television that if a pregnant woman gets tested for HIV, it is possible to prevent the transmission of the

virus to the baby as the woman is seen by health workers and get treatment.... (Participant 3).

ANC visits for previous pregnancies were also mentioned as one of the sources of women's awareness that HIV testing is performed at the health institution and their knowledge on the benefits of HIV testing. As elaborated by one woman.

Whenever you come for follow up (ANC), they (health workers) first test your blood for HIV. In case I am infected with HIV, they have something to do (give treatment)... (Participant 4).

According to the current HIV testing guidelines, women must be explicitly informed of their right either accept or to refuse testing. The guidelines, stated that, clients should receive pre-test information in a group or individually on HIV/AIDS and PMTCT.

In this study, Our in-depth interviews among pregnant women found that the pre-test counselling accompanying the test was limited. HIV testing was not however perceived as a choice, but rather as a compulsory service for all pregnant women. Some participants stated that pregnant women are tested for HIV as part of ANC, along with other routine examining procedures.

Blood is taken from my hand (showing her finger) for HIV testing. The HIV testing is performed anytime we are pregnant and come for follow-up... (Participant 5).

Another woman elaborated the perception that routine HIV testing is a government policy;

They (health workers) simply told me to test my blood for HIV. I didn't ask why they needed my blood since I thought that the test is a must... (Participant 2).

This limited pre-test counselling was reflected by some women's lack of knowledge about the possibility of protecting children born to HIV positive women as one woman described below:

If the mother is healthy, the baby will also be healthy. Pregnant woman's knowledge of her HIV status has no benefit to the baby if she has already been infected with the virus... (Participant 7).

Though, health workers acknowledged the importance of providing pre-test information they reported that they provide limited pre-test counselling due to large number of clients.

Counselling is important. If there are few clients we explain to them some introduction about PMTCT and then do HIV testing. Of course, counselling is missed. If we take longer time in providing pre-test counselling, others who wait outside will complain... (Health worker 4).

In addition, another health worker emphasized on the importance of post-test counselling for HIV positive women compared to for those who are HIV negative women. She believed that there is no need to counsel HIV negative women as the goal of HIV testing is to prevent mother to child transmission of HIV.

The goal that needs to be emphasized is to save the baby. If the woman becomes HIV positive, there will be more counselling after the test. (Health worker 3).

When asked if they know that routine HIV testing provided as a part of ANC is a choice, only few knew that HIV testing is optional. Those who knew it was optional indicated their views as follows:

It is based on my will. It is known that HIV testing has a benefit, but it is possible to refuse. All women should be tested but should be on their will. I should be tested for HIV based on my interest... (Participant 8).

Health workers also approved the compulsory nature of the HIV testing as follows:

It is well documented that HIV testing should be voluntary, as stated in the guidelines. But you (health worker) emphasize on the importance of HIV testing when you give counselling, and you must push them to accept... (Health worker 1).

HIV testing perceived as a pre-requisite to other health services

We asked our participants if refusal of HIV testing impacts the services that a woman gets from the health facility. Some women were concerned about their inability to decline the HIV

test. They thought that refusal of HIV testing could result in a denial of other health services including delivery service:

I can't refuse. Refusing HIV testing will create a problem to the woman. If she refuses what they (health care providers) ask her to do, they may not help her when she come back for delivery...(Participant 6)

Research Question 2: What are the experiences of pregnant women about Routine HIV counselling and testing in tertiary hospitals?

Table 2. Experiences of pregnant women about Routine HIV counselling and testing in tertiary hospitals

| S/N | Items | SA | S | D | SD |
|-----|-----------------------------------------------------------------------------------------------|----------|----------|---------|--------|
| | There are routine plans for checking HIV status during pregnancy | (62.9%) | (26.9%) | (5.7%) | (4.8%) |
| | there are routine strategy for checking HIV status though it has nothing to do with pregnancy | (71.50%) | (24.10%) | (1.20%) | (3.2%) |
| | Counselling sessions makes one feel uncomfortable | (51.4%) | (29.1%) | (12.6%) | (6.9%) |
| | Antenatal includes HIV screening test | (62.9%) | (26.9%) | (5.7%) | (4.6%) |
| | Because of stigmatization routine HIV testing is avoided. | (55.4%) | (21.7%) | (16.6%) | (6.3%) |
| | HIV screening test makes one feel uncomfortable | (58.2%) | (18.7%) | (16.6%) | (6.5%) |

Table 2 revealed that 62.7% of the respondents strongly agreed that they do have routine plans on checking HIV status during pregnancy, 71.5% strongly agreed that they do have a routine strategy though it has nothing to do with their pregnancy, 51.4% strongly agreed that

they do not like the counselling session it makes me feel uncomfortable, 62.9% strongly agreed that they do not like the counselling session it makes me feel uncomfortable, 55.4% strongly agreed that they are afraid of been stigmatize so they avoid routine test, 58.2% strongly agreed that they feel uncomfortable whenever they go for HIV screening test.

4.2 Hypothesis

H₀₁: Educational qualification does not significantly determine the perception of pregnant women about Routine HIV counselling and testing in tertiary hospitals in River State.

Comment [TM6]: Under methodology put on the model specification for this regression analysis

Table 3: Educational qualification does not significantly determine the perception of pregnant women about Routine HIV counselling and testing in tertiary hospitals in River State.

| Model | Sum of Squares | df | Mean Square | F | Sig. |
|------------|----------------|-----|-------------|------|-------------------|
| Regression | 899.992 | 1 | 5.992 | .084 | .199 ^b |
| Residual | 2136.758 | 219 | 1.379 | | |
| Total | 3036.75 | 220 | | | |

The table 3 shows that the computed F is 0.084 while the p value = 0.199 therefore, since $P < .05$, i.e. $p = .001$ is statistically significant at the chosen alpha level of 0.05 meaning that educational qualification significantly determine the perception of pregnant women about Routine HIV counselling and testing in tertiary hospitals in River State

4.0 Discussion

In this study, routine HIV counselling and testing was well received and accepted by most pregnant women. This was apparent as most women were aware of the routine prenatal HIV testing before registering for ANC services. The women indicated that they were pre-information about HIV testing and its benefits through various community education activities, mass media and during their previous prenatal visits for the most recent pregnancy. During ANC services, health care workers were over-stressed by the large numbers of women waiting for HIV testing. This was noticed to have downplayed or reduced the quality of counselling services thereby affected the quality of PMTCT services. To this end,

strategies such as, increased and deployment of more staff to ANC days to ensure that, HIV counselling and testing take place in the manner it was meant for.

In this study, women valued routine HIV counselling and testing as this gave them an insight into their own status and also, protect their unborn babies, knowing that, those that are HIV positive could access treatment and suppressed the viral load. Therefore, acceptance of HIV testing in the study setting may be an indication why women are in full agreement to protect their unborn children as well as taking care of their own health. It was also, observed that, some studies in Africa also identified the importance for mothers / women to protect their children (both born and unborn) and to show concern for their own health as a major

indicator for acceptance of HIV counseling and testing during pregnancy (Branson, 2017). Also, some women that are aware of the voluntary option of opt-out HIV testing, were against the routine HIV testing. The option that women should have the right to accept or refuse HIV testing had been documented in other African settings (Chandisarewa et al., 2017). This results were in line with “global guidelines and Ethiopian HIV testing policy that clients’ right to informed consent should be respected” (Nigatu & Woldegebriel, 2016).

In routine provider-initiated HIV counseling and testing, as stipulated in global and national guidelines, clients must receive pre-test information in a group or individually on HIV/AIDS and PMTCT (Branson et al., 2018). In our study, however, majority of pregnant women, along with the data from the interviews with health workers portrayed that pre-test counselling was limited. Similarly, in a study conducted in Addis Ababa, revealed that providers took blood samples for HIV test from clients without pre-test counselling (Angotti et al., 2016). Health care providers mentioned that counselling was particularly important for HIV positive pregnant women, implied that counselling was not critical if a woman was HIV negative. This finding suggested that counselling was not prioritized for majority of people who tested negative, which indicated missed opportunities for primary prevention of HIV. Pre-test counselling was vital for pregnant women to ensure that they understand the implications of negative or positive test results for themselves, their partners and their unborn children (Larsson et al 2016).

In this study, client's viral load was raised as a barrier to pre-test counselling. In the ANC environments, health care providers were overwhelmed by the great numbers of women who attended their clinics for testing. This was observed to have affected the quality of counselling services, thus hampered the delivery of full package of PMTCT services. Similarly, studies in sub-Saharan Africa countries had demonstrated limited pre-test counselling related to the shortage of health workers in these facilities (Perez et al., 2016). To this effect, strategies included increased in the number of health workers and the use of lay counsellors to ensure that provider-initiated HIV testing and counselling took place in the manner it was intended to (Rujumba et al., 2016).

The Ethiopia PMTCT guidelines recommended that providers must explicitly inform the client the right to say "no" (to opt-out), and that this decision by no means affect the services rendered in the health facility (Mugore et al., 2018). In this study, however, some women did not perceive HIV testing as a choice, but rather as a compulsory service for all pregnant women. The perception that HIV testing provided as part of ANC was compulsory, has also been documented in other African settings (Hardon et al., 2017). Some women perceived this approach as a government policy intended to protect children from HIV infection. For such women, the implication was that acceptance of HIV testing could be considered as a compliance with what they perceived as a government law. Health workers' emphases on the benefits of HIV testing during pre-test counselling found in this study, could have downplayed women's perceptions of the possibility of not testing, which was in congruent with a study conducted in Uganda (Ismail & Ali, 2019). Similarly, a study from four African countries revealed that some providers underlay the stance as the moral imperative to protect the unborn child (Perez et al., 2016). Moreover, some participants perceived HIV testing was a prerequisite to receive other health services including delivery services. Similar findings have been documented in other African settings (Njeru et al., 2017). This was a fundamental shortcoming of unclear pre-test counselling which entirely eliminated pre-test counselling or provided insufficient information to the opportunities for informed consent (Steen et al., 2017).

5.0 Conclusions

Routine HIV counselling and testing which was provided as part of antenatal care services in the study setting, was well accepted among the pregnant women which will enabled them to make informed choices regarding the HIV testing and to take appropriate action. In light of this, there was a sense of obligation as women feel HIV counselling and testing was a requirement for delivery services as an evident that participants were able to state the benefits of routine counselling and testing.

Participants' views were that there was a low acceptance of routine antenatal counselling and testing, because, the voluntary routine HIV testing was not fully explained and understood, which may be related to the limited pre-test counselling. The study suggest that, there was need to strengthen pre-testing counselling to ensure that HIV testing is conducted in a way that would ensure pregnant women's autonomy and maximise primary HIV prevention opportunities.

This study also provided useful information about routine antenatal counselling and testing. Constraints which hindered the implementation of routine antenatal counselling and testing were identified, and recommendations to facilitate the application of the principles were made.

5.0 Recommendations

The recommendations of the study are based on the suggestions made by the participants. Stigma and discrimination has been identified as a barrier to the acceptance of routine antenatal counselling and testing. Therefore, there is a serious call for interventions that would deal with violence against women and men over women's sexualities in communities. The men should be educated about the importance of routine antenatal counselling and testing. This would encourage partners to share the burden of HIV status and reduce stigma

and discrimination. This cannot change overnight; therefore it remains a challenge to society at large to change their perceptions.

1. Community education about the importance of routine antenatal counselling and testing and PMTCT should be strengthened. It will encourage community involvement and the reduction of stigma and discrimination in the society. It will also improve the knowledge about the benefits of routine antenatal counselling and testing. Information and education empower individuals to make correct decisions concerning health care.
2. Since the healthcare workers are the important support system for pregnant women with HIV/AIDS, they should keep the information regarding the HIV results confidential and provide care and support to those who are HIV positive.
3. Routine antenatal counselling and testing should be available and accessible at all health facilities offering antenatal services. This will allow all pregnant women to benefit from the available interventions designed to reduce HIV infection among children.
4. Effective communication and cooperation should be encouraged between partners. It will increase the male involvement in the programme and ensure that men know the importance of routine antenatal counselling and testing. Male involvement will allow both partners to accept and share their results (negative or positive).
5. Cultural practices should be discouraged in order to prevent HIV transmission from the mother to the baby.
6. A policy on stigmatization and discrimination should be effectively implemented. It will help to reduce and eliminate the stigma and discrimination attached to HIV/AIDS.
7. Routine antenatal counselling and testing services should be extended to all hospitals, health centers and clinics that are offering antenatal services.

REFERENCES

- Adekeye A. (2015). Attitude of counsellors towards undergoing HIV voluntary counselling and testing in Ado-Odo/Ota, Nigeria. *International Journal of Psychology and Counselling*, 3(3), 33-39.
- Adekeye, O., Ebiai, A., & Adeusi, S. (2011). Psycho-cultural variables predicting attitude of students' towards HIV counselling and testing in selected tertiary institutions in Lagos state, Nigeria. *JORIND* (9)1 431-440
- Aini-kaarin T. (2005). Perceptions and experiences of pregnant women towards HIV voluntary antenatal counselling and testing in Oshakati hospital, Namibia. A-mini-thesis submitted in partial fulfilment of the requirements for the degree of Masters in Public Health, the Faculty of School of Public Health, University of the Western Cape.
- Becker, M. H. (1974). *The Health Belief Model and Personnel Health Behaviour*. San Francisco: Society for Public Health Education, Inc
- Bello, I B. (2016). Personality traits as predictors of psychological health among secondary school adolescents in Ede, Osun State. A Research project submitted to the department of behavioural studies, college of management and social sciences, Redeemer's University, Ede, Osun State, Nigeria
- Branson B. (2017). Current HIV epidemiology and revised recommendations for HIV testing in health care settings. *Public Health Rep*, 122(5):579–83.
- Branson BM, Handsfield HH, Lampe MA, Janssen RS, Taylor AW, Lyss SB, (2018). Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *J Natl Med Assoc*. 100(1):131–47.
- British Medical Association, (1987). WHOQOL Group. Development of the WHOQOL: Rationale and current status. *International Journal Mental Health*, 23, 24–56.
- Carlsson G., (2020). UN (United Nations) Global AIDS Update 2019: Communities at the Centre, Defending Rights, Breaking Barriers and Reaching People with HIV Services . 1st. New York, NY, USA.

- Central Statistical Agency. (2008). The 2007 National Census Preliminary Report for Ethiopia Addis Ababa. Ethiopia: Central Statistical Agency..
- Chandisarewa W, Stranix-Chibanda L, Chirapa E, Miller A, Simoyi M, Mahomva A, (2017). Routine offer of antenatal HIV testing (“opt-out” approach) to prevent mother-to-child transmission of HIV in urban Zimbabwe. *Bull World Health Organ*, 85(11);843–50.
- Clinical Ophthalmology (Auckland, NZ)*, 6, 2023.
- Cohen R, Lynch S, Bygrave H, Eggers E, Vlahakis N, Hilderbrand K. (2019). Antiretroviral treatment outcomes from a nurse-driven, community-supported HIV/AIDS treatment programme in rural Lesotho: observational cohort assessment at two years. *J Int AIDS Soc*. 12(1); 23.
- CONOMIC, F., (1991). IN RIVERS STATE, NIGERIA. *African Study Monographs*, 12(4),
- Corneli A, (2010). Patient and provider perspectives on HIV testing and counselling for TB patients in Kinshasa, Democratic Republic of Congo. Poster presented at 3rd IAS conference on HIV pathogenesis and treatment. Rio de Janeiro, Brazil, 24-27.
- Creek TL, Ntuny R, Seipone K, Smith M, Mogodi M, Smit M, (2017). Successful introduction of routine opt-out HIV testing in antenatal care in Botswana. *J Acquir Immune Deficiency Syndrome*. 45(1);102–7.
- Creswell, J., &Poth, C. (2013). *Qualitative inquiry and research design: Choosing among five approaches* (Fourth ed.). Los Angeles: SAGE Publications.
- CSA . (2021), *Ethiopia Demographic and Health Survey*. Calverton:Central Statistical Authority, Addis Ababa and ORC Macro.
- Dapaah, J. M., &Senah, K. A. (2016). HIV/AIDS clients, privacy and confidentiality; the case of two health centres in the Ashanti Region of Ghana. *BMC medical ethics*, 17(1), 1-10.
- Davenport, K. (2000). Perception of a Chilly Climate: Differences in Traditional and Non-traditional Majors for Women. *Research in Higher Education*..49 256-273
- De Cock KM, Bunnell R, Mermin J. (2016). Unfinished business—expanding HIV testing in developing countries. *N Eng J Med*. 354(5):440–2.

- De Cock, K., Granich R.M., Dye C., & Williams B. G. (2009). HIV/AIDS care. HIV/AIDS, Maternal and Child Health in Africa. *Lancet*, 359:2027-107
- De Veirman, M., Hudders, L., & Nelson, M. R. (2019). What is influencer marketing and how does it target children? A review and direction for future research. *Frontiers in Psychology*, 10, article2685. <https://doi.org/10.3389/fpsyg.2019.02685>
- FMOH .(2020) *Report on progress towards implementation of the UN Declaration of Commitment on HIV/AIDS 2020*. Addis Ababa.
- Federal Democratic Republic of Ethiopia(2018). Federal HIV/AIDS Prevention and Control Office.Federal Ministry of Health. National HIV & AIDS and Reproductive Health Survey 2016. NARHS plus II.Federal Ministry of Health.
- Federal Ministry of Health(2015). Technical Report: 2013 National HIV Seroprevalence Sentinel Survey.
- Galadanci, S., Iiyasu, Z., Tukur, J., Muktar-Yola, M., & Adeleke. I. (2008). Uptake of voluntary counselling and testing for HIV by pregnant women in a prevention-of-mother-to-child-transmission programme at Aminu Kano Teaching Hospital, Nigeria. *African Journal of AIDS Research*, 7(1), 143–148
- Garbin, C., Pacheco, K., Santiago, T., Miyada, S., Garbin, A., &Moimaz, S.(2015). Perception of HIV among pregnant women in the public health system in two municipalities of the state of São Paulo. *Brazillian Journal Oral Science*, 14(4), 282-286
- Gruskin S, Ahmed S, Ferguson L (2018). Provider-initiated HIV testing and counseling in health facilities-what does this mean for the health and human rights of pregnant women? *Dev World Bioeth*. 8(1):23–32.
- Gunn JK., Asaolu IO., Centre KE., Gibson SJ., Wightman P., &Ezeanolue EE. (2016). Antenatal care and uptake of HIV testing among pregnant women in sub-Saharan Africa: a cross-sectional study. *Journal of the Inter- national AIDS Society*, 19(1):20605. <https://doi.org/10.7448/IAS.19.1.20605> PMID: 26787516
- Hardon Anita, Vernooij Eva, Bongololo-Mbera Grace, Cherutich Peter, Desclaux Alice, Kyaddondo David, (2017). Women’s views on consent, counselling and

confidentiality in PMTCT: a mixed-methods study in four African countries. *BMC Public Health*. 12:26.

Hugger, K., Braun, L. M., Noll, C., Nowak, T., Gräßer, L., Zimmermann, D., & Kaspar, K. (2019). Zwischen Authentizität und Inszenierung: Zur medienkritischen Einschätzung information orientierter YouTuber*innen-Videos durch Jugendliche [Between authenticity and staging: On the media-critical evaluation of information oriented YouTuber videos by young people]. In: F. von Gross, & R. Röllecke (Hrsg.), *Instagram und YouTube der (Pre-)Teens – Inspiration, Beeinflussung, Teilhabe* [Instagram and YouTube of (pre-)teens – inspiration, influence, participation] (pp. 29–36). Munich: kopaed.

Ismail H.; Ali A. (2019). Status of ANC-linked HIV counseling and testing as an intervention for PMTCT in public health facilities in Addis Ababa: quality of HIV counseling given to pregnant women for PMTCT. *Ethiop J Health Dev*. 23(3):190–8.

Jones CL, Jensen JD, Scherr CL, Brown NR., Christy K, & Weaver J. (2015) The health belief model as an explanatory framework in communication research: exploring parallel, serial, and moderated mediation. *Health communication*, 30(6), 566–76. <https://doi.org/10.1080/10410236.2013.873363> PMID:25010519

Kendra, H. (2012). The perception and social skill In: Desai, V., & Potter R. B. ed. *The Companion to Development Studies*. London: Hodder Education.

Kwapong GD., Boateng D., Agyei-Baffour P., & Addy EA. (2014). Health service barriers to HIV testing and counselling among pregnant women attending Antenatal Clinic; a cross-sectional study. *BMC health services research*, 14(1):267. <https://doi.org/10.1186/1472-6963-14-267> PMID: 24942820

Larsson EC, Thorson A, Pariyo G, Conrad P, Arinaitwe M, Kemigisa M, (2015). Opt-out HIV testing during antenatal care: experiences of pregnant women in rural Uganda. *Health Policy Plan*. 27(1):69–75.

Lou, C., & Yuan, S. (2019). Influencer marketing: How message value and credibility affect consumer trust of branded content on social media. *Journal of Interactive Advertising*, 19(1), 58–73. <https://doi.org/10.1080/15252019.2018.1533501>

- Martínez, C., & Olsson, T. (2019). Making sense of YouTubers: How Swedish children construct and negotiate the YouTuber Misslisibellas a girl celebrity. *Journal of Children and Media*, 13(1), 36–52. <https://doi.org/10.1080/17482798.2018.1517656>
- Mathingau, L. (2013). Factors influencing acceptance of routine HIV testing and counselling among pregnant women in Kibera informal settlement, Nairobi, Kenya. A research thesis submitted in partial fulfilment of the requirement of degree of master of public health in the school of public health of Kenyatta University
- Ministry Of Health and Social Services (2014). *Guidelines for the Prevention of Mother-to-Child Transmission of HIV*. 1st edition. Windhoek: Namibia.
- Mitiku, I., Addissie, A., & Molla, M (2017). Perceptions and experiences of pregnant women about routine HIV testing and counselling in Ghimbi town, Ethiopia: a qualitative study. *BMC Research Notes*, 10:101
- Msellati P. (2019). Improving mothers' access to PMTCT programs in West Africa: a public health perspective. *Soc Sci Med.*69(6):807–12.
- Mugore L, Engelsmann B, Ngoro T, Dabis F, Perez F. (2018). An assessment of the understanding of the offer of routine HIV testing among pregnant women in rural Zimbabwe. *AIDS Care*. 20(6):660–6.
- Muhenje O., Mbori-Ngacha D., Akun T., DeCock K. (2008). Barriers to acceptance of PMCT services at a Provincial General Hospital, Kisumu, Kenya. Program and abstracts of the 15th International AIDS Conference; July 11-16 2004; Bangkok, Thailand. Abstract ThPeB7090.
- Nwozichi, C., & Farotimi, A. (2014) Knowledge and Acceptance of HIV Counselling And Testing (HCT) among Non-Teaching Staff of A Nigerian Private University. *IOSR Journal of Nursing and Health Science* 3(1), 57-62
- Oladeinde, B. H., Omoregie, R., Olley, M., & Anunibe, J. A. (2011). Prevalence of HIV and anemia among pregnant women. *North American journal of medical sciences*, 3(12), 548.
- Olajide, O.M., Oguntayo, A.O., & Kolawole, A.O. (2015) Acceptability of HIV Testing and Counselling by Antenatal Clients of a Tertiary Institution in Northern Nigeria. *Open*

Comment [TM7]: There is nowhere where this is referenced in this study. Re-check your references.

Comment [TM8]: Same comment as above.

Journal of Obstetrics and Gynecology, 5, 94-102.<http://dx.doi.org/10.4236/ojog.2015.52013>

- Olanrewaju, A., Ola, F., Akintunde, O., Ibrahim, B., &Ibiyemi F. (2017). HIV Voluntary Counselling And Testing Of Pregnant Women In Primary Health Care Centres In Ilesa, Nigeria. *The Internet Journal of Third World Medicine*, 6 (1) 23-45
- Pai NP, Barick R, Tulsy JP, Shivkumar PV, Cohan D, Kalantri S, Pai M, Klein MB, Chhabra S: Impact of round-the-clock, rapid oral fluid HIV testing of women in labor in rural India. *PLoS Medicine*, 5: 5-10.1371/journal.pmed.0050005
- Perez F, Zvandaziva C, Engelsmann B, Dabis F. (2016). Acceptability of routine HIV testing (“opt-out”) in antenatal services in two rural districts of Zimbabwe. *J Acquir Immune Defic Syndr*. 2006;41(4):514–20.
- Perez F., Orne-Gliemann J., Mukotekwa T., Miller A., Glenshaw M., Mahomva A., (2016). Prevention of mother to child transmission of HIV: evaluation of a pilot program in a district hospital in rural Zimbabwe. *BMJ*. 329:1147–50.
pp.167-183.
- Steen TW, Seipone K, Anderson MG, Kejelepula M, Keapoletswe K, Moffat HJ. (2017). Two and a Half years of routine HIV testing in Botswana. *AIDS*. 44(4):484–8.
- Stringer, J. (2003). Nevirapine to Prevent Mother-to-Child Transmission of HIV-1. Among Woman of Unknown Sero status. *African Medical Journal*, 13, 1850-1853.
- Ufornwa EC, OkoroiwuHU(2017). HIV post exposure prophylaxis in a tertiary health care institution in Southeastern Nigeria: Clients’ characteristics and indications for initiation. *Nigeria Health J*.
- UNAIDS (2020). Seizing the moment: Tackling entrenched inequalities to end epidemics. Geneva: Joint United Nations Programme on HIV/AIDS (UNAIDS)
- UNAIDS ‘AIDSinfo (2018). Available at: <http://aidsinfo.unaids.org/> (Accessed:17.12.2018).
- UnitedNations(2021). *Political Declaration on HIV and AIDS: Intensifying Our Efforts to Eliminate HIV and AIDS. Resolution Adopted by the General Assembly, A/RES/65/277* . San Francisco, CA, USA: The United States of America: 95th Plenary Meeting, United Nations.

World Health Organization(2020).*Global Tuberculosis Report 2020* . 1st. Geneva, Switzerland: World Health Organization.

WHO(2016).*Policy Brief: Consolidated Guidelines on HIV Prevention, Diagnosis, Treatment and Care for Key Populations, 2016 Update* . 2nd. Geneva, Switzerland: WHO Library Cataloguing-in-Publication Data, World Health Organization.

UNDER PEER REVIEW