

## ISOLATION AND ANTIBIOTIC RESISTANCE OF *STAPHYLOCOCCUS AUREUS* ISOLATED FROM NOSOCOMIAL SOURCES

### Abstract

Methicillin resistance *Staphylococcus aureus* have been reported worldwide to emerge mostly in developing and developed countries. This study aimed at isolated and antibiotic resistance from nosocomial sources in Dalhatu Araf specialist Hospital, Lafia, Nigeria. A total of (200) samples were collected from February 2021 to May 2021 from different Nosocomial sources such as door handles, seat handles, surgical equipment and stretchers and *Staphylococcus aureus* was isolated and identified using standard microbiological method. The Antibiotic susceptibility test for the isolates were carried out and interpreted in accordance with clinical and laboratory standard institute (CLSI) protocol. The occurrence of *Staphylococcus aureus* from the samples was 50 (25%). The highest occurrence of *Staphylococcus aureus* is from seat handle swab with (32%) and the lowest occurrence is (18%) from surgical equipments. The Antibiotic resistance of *Staphylococcus aureus* showed that the isolates were more resistant to oxacillin.

### Introduction

Staphylococci are Gram-positive *cocci*, usually commensal organisms that are found occurring on the skin and mucosa of humans and animals. *Staphylococcus aureus* (*S. aureus*) is an important pathogen of clinical significance, causing variety of illnesses in both humans and animals worldwide (Chakraborty *et al.*, 2011). It causes superficial skin infections and life-threatening diseases including endocarditis, sepsis and soft tissues, urinary tract, respiratory, intestinal tract, and bloodstream infections (Rallapalli *et al.*, 2008). Close association has been observed to enhance spread of staphylococcal strains among livestock and veterinary care-givers and animal handlers through contact or aerosol (Ajuwape *et al.*, 2001). *S. aureus* is a major food borne pathogen due to its production of enterotoxins that cause serious intoxications (Wu *et al.*, 2011; Liu *et al.*, 2014). Fast identification of *S. aureus* and its toxins in food is crucial to

**Comment [ks1]:** Please put the key words.

May be:  
Hospital Infection Control  
Program, *Staphylococcus aureus*,  
Vancomycin Resistance  
Methicillin-Resistant *Staphylococcus aureus*

**Comment [ks2]:** please put more recent references

look:  
Infection Control  
[Yacob Habboush](#), [Siva Naga S. Yarrarapu](#)<sup>1</sup>, [Nilmarie Guzman](#)<sup>2</sup>  
•Bookshelf ID: [NBK519017](#)

determine microbial risk and assure food quality. According to Bautista Trujillo *et al* (2013) rapid isolation and identification of the *S. aureus* pathogen is a major goal of diagnostic microbiology. A variety of selective or differential culture media have been used to isolate and identify the organism (Thakar *et al.*, 2013). The use of culture media for *S. aureus* isolation in combination with coagulase activity and haemolysis determination as secondary tests have improved the accuracy of identification, and was in consonance with gene sequence analysis compared with the use of the culture media alone (Bautista Trujillo *et al.*, 2013). The management of *S. aureus* infections especially methicillin resistant ones is often difficult because methicillin resistant *S. aureus* (MRSA) is usually resistant to multiple antibiotics. Vancomycin is commonly used to treat such infections and occasionally, Macrolide-Lincosamide Streptogramin B (MLS<sub>B</sub>) family of antibiotics are used as substitute (Adhikari *et al.*, 2017). Due to the rising incidences of methicillin resistance, glycopeptides such as vancomycin have been recommended as therapeutic agents for serious staphylococcal infections (Nunes *et al.*, 2006). However, the extensive use of glycopeptides has decreased the susceptibility of staphylococcal species to these agents (Pfeltz *et al.*, 2000; Finan *et al.*, 2001; Cui *et al.*, 2003; Kim *et al.*, 2012). Inducible vancomycin resistance is due to a sophisticated mechanism that combines synthesis of cell wall peptidoglycan precursors with low affinity for glycopeptides and elimination of the normal target precursors (Foucault *et al.*, 2010). *Staphylococcus aureus* develops resistance to antimicrobials by employing different mechanisms. These mechanisms include limiting uptake of the drug, modification of the drug target, enzymatic inactivation of the drug, and active efflux of the drug. The bacteria may use one or several of these mechanisms depending on the antimicrobial. In particular, the localization of resistance genes on transferable genetic elements such as plasmids and transposons facilitates horizontal transfer of resistance between bacteria (Van Hoek *et al.*, 2011). The development of such resistance does not cause the organism to be more intrinsically virulent than strains of *Staphylococcus aureus* that have no antibiotic resistance, but resistance does make MRSA infections more difficult to treat with standard types of antibiotics and thus more dangerous (Jenson & Lyon, 2009). Methicillin-resistant *Staphylococcus aureus* (MRSA) is a gram-positive bacterium that is resistant to methicillin (a member of the penicillin family) and many other beta-lactam antimicrobials (beta-lactam antimicrobials include penicillins and cephalosporins), and are resistant to macrolides and aminoglycosides. Methicillin-resistant *Staphylococcus aureus* (MRSA) is an isolate of *Staphylococcus aureus* which has acquired genes

**Comment [ks3]:** please put more recent references look:

Device-associated infection rates and bacterial resistance in six academic teaching hospitals of Iran: Findings from the International Nosocomial Infection Control Consortium (INICC)

[Somayeh Jahani-Sherafat](#)<sup>1</sup>, [Maryam Razaqhi](#)<sup>1</sup>, [Victor D Rosenthal](#)<sup>2</sup>, [Elahe Tajeddin](#)<sup>3</sup>, [Simasadat Seyedjavadi](#)<sup>4</sup>, [Marjan Rashidan](#)<sup>1</sup>, [Masoud Alebouyeh](#)<sup>3</sup>, [Maryam Rostampour](#)<sup>4</sup>, [Arezo Haghi](#)<sup>5</sup>, [Masoumeh Sayarbayat](#)<sup>6</sup>, [Somayeh Farazmandian](#)<sup>7</sup>, [Tahere Yarmohammadi](#)<sup>8</sup>, [Fardokht K Arshadi](#)<sup>9</sup>, [Nahid Mansouri](#)<sup>10</sup>, [Mohammad R Sarbazi](#)<sup>11</sup>, [Mariano Vilar](#)<sup>2</sup>, [Mohammad R Zali](#)<sup>12</sup>

Affiliations expand

• PMID: 26027477

• DOI: [10.1016/j.jiph.2015.04.028](https://doi.org/10.1016/j.jiph.2015.04.028)

encoding antibiotic resistance to all penicillins including methicillin. This resistance is mediated by an altered penicillin binding protein (PBP2a) which is encoded by the *MecA* gene that is carried on a large mobile genetic element, the staphylococcal cassette chromosome (Palavecino, 2007; Ahmed *et al.*, 2012).

In Africa, countries show different MRSA prevalence (Bell & Turnidge, 2002). MRSA is one of the major causes of infections in humans, occurring in both the community and the hospital (Ugwu *et al.*, 2016). Akerele *et al.* (2016) reported that acquisition of MRSA has been associated with two different environments; Community-associated MRSA (CA-MRSA) and healthcare-associated MRSA (HA-MRSA). They are usually differentiated by their structural and functional genomic traits (Otto, 2013). MRSA infections in the community can also be caused by livestock-associated methicillin resistant *Staphylococcus aureus* (LA-MRSA). It is initially associated with livestock (Lewis *et al.*, 2008; Layer *et al.*, 2012), and differs from genotypic HA-MRSA and genotypic CA-MRSA in its genomic traits. The risk factors for community acquired infection include intravenous drug use, close contact with persons who have MRSA, men who have sex with men, crowding, poor hygiene, recent antibiotic use, and previous hospitalization (Moran *et al.*, 2006; King *et al.*, 2006; Hota *et al.*, 2007; Boucher & Corey, 2008), while risk factors associated with nosocomial acquired MRSA colonization and infection are advanced age, male gender, previous hospitalization, length of hospitalization, stay in an ICU, chronic medical illness, prior and prolonged antibiotic treatment, presence and size of a wound, exposure to colonized or infected patient, presence of invasive indwelling devices (Kaye *et al.*, 2000).

### Study area

This study was carried out at Dalhatu Araf Specialist Hospital (DASH) in Lafia, Nasarawa State, Nigeria. DASH was established in 2003 by the Nasarawa State Government to cater for the health needs of the people of the state at the tertiary level.

### Ethical Clearance

This study was conducted according to good laboratory conditions and in accordance with the declaration of national institutional standard. Also written consent was obtained from the state

**Comment [ks4]:** it is not necessary to put the name of the hospital  
put only the country and city where the study was carried out.

research ethical committee through the state ministry of health, and also written consent was obtained from the research ethical committee of Dalhatu Araf Specialist Hospital, Lafia.

### Sample Size Determination

A prevalence of 16% was used. This is based on studies carried out in Northern Nigeria (Okon *et al.*, 2011; 2014). The sample was then determined using the formula;

$$n = \frac{pqz^2}{d^2}$$

n = minimum sample size required

p = proportion of the target population estimated to have particular problem

q = 1 – p

z = level of precision (1.96) which corresponds to 95 % confidence level

d = degree of accuracy desired set at 0.05

$$n = \frac{0.16 (0.84) (1.96^2)}{0.05^2} = 200$$

### Sample Collection

A total of two hundred (200) samples were collected from different nosocomial sources such as catheters, bed handles, door handles, dishes, forceps and toilet seats within the hospital. Aseptic procedures was used for the collection, surfaces was swabbed using sterile cotton swabs immersed in normal saline solution.

### Isolation and identification of *Staphylococcus aureus*

The swab-sticks containing the specimen was inoculated in nutrient broth and incubated at 37°C for 18 h. It was then sub-cultured on Mannitol Salt Agar (MSA) by streaking, then incubated at 37°C for 24 h, and the cultural characteristics of colonies on the MSA was observed. Golden-yellow colonies were indicative of *S. aureus* (Owaku *et al.*, 2017). Presumptive *S. aureus* was identified by microscopy (Gram staining), biochemical tests and commercial kit identification.

### Antimicrobial Susceptibility Testing

The antimicrobial susceptibility testing of the isolates was carried out using Kirby-Bauer disk diffusion method as described in Clinical and Laboratory Standards Institute (CLSI) Guidelines (CLSI, 2012).

**Result**

The cultural, morphological and biochemical characteristics of *S.aureus* isolated from fomites are as shown in Table 1

The occurrence of *S. aureus* in the fomites is as shown in Table 2. From the 200 samples, 50 (20.0%) *S. aureus* were isolated, with the highest occurrence (32.0%) from swabs taken from seat handles and the lowest (18.0%) from surgical equipment.

Streptomycin had 80% which was the highest, Penicillin had 76%, Rifampicin had 72%, Ampicilin/sulbactam had 64%, Clindamycin had 64% , Oxacillin had 62%, Ceftriaxone had 58%, Vancomycin had 28%, Levofloxacin had 26% and Gentamycin had 22% which is the lowest as shown in Table 3.

**Table1. Cultural, Morphological and Biochemical characteristics of *Staphylococcus aureus***

Cultural characteristics	Morphological Characteristics	Biochemical Characteristics													Inference	
		C	C	V	A	ON	U	A	M	S	L	A	R	T	M	
		at	oa	p	kp	PG	r	rg	an	u	ac	r	f	r	al	
Golden yellow colonies	+ Cocci in cluster	+	+	+	+	-	+	+	+	+	+	-	-	+	+	<i>S. aureus</i>

---

on MSA

---

MSA= Mannitol Salt Agar; Cat= Catalase; Coa= Coagulase; Vp= Voges-Proskauer; Akp= Alkaline phosphatase; ONPG= Ortho-nitrophenyl- $\beta$ -galactoside; Ur= Urease; Arg= Arginine Utilization; Man= Mannitol; Su= Sucrose; Lac= Lactose; Ar= Arabinose; Rf= Raffinose; Tr= Trehalose; Mal= Maltose; + =Positive; +w= Positive to weak reaction; - =Negative

**Table2. Isolation rates of *Staphylococcus aureus* in relation to the fomites.**

---

Source	No. of Samples	No. (%) of <i>S. Aureus</i>
Door handles	50	12 (24.0)
Seat handles	50	16 (32.0)
Surgical equipment	50	9 (18.0)
Stretchers	50	13 (26.0)
<b>Total</b>	<b>200</b>	<b>50 (25.0)</b>

---

**Table3. Antibiotic Resistance Profile of *Staphylococcus aureus* isolated from fomites**

---

Antibiotics	Disc Content ( $\mu$ g)	No. (%) of Resistance (n=50)
Rifampicin	5	26 (72.0)
Clindamycin	2	22 (64.0)
Vancomycin	15	14 (28.0)
Levofloxacin	5	13 (26.0)
Oxacillin	1	31 (62.0)
Ceftriaxone	30	29 (58.0)
Ampicilin/sulbactam	25	32 (64.0)
Streptomycin	25	40 (80.0)

---

Gentamycin	30	11 (22.0)
Penicillin	5	38 (76.0)

## Discussion

*Staphylococcus aureus* has been identified as one of the most common pathogens associated with both hospital and community-acquired infections worldwide, with various infections that are devastating and life-threatening (Peterson *et al.*, 2013; Yaw *et al.*, 2014). The propensity for staphylococci to develop antimicrobial resistance is a cause for great concern in both human medicine (Kaur & Chate, 2015). Antimicrobial resistance was high-lightened as an urgency issue (Acar & Moulin, 2012). Decades now, multidrug resistance in *S. aureus* has spread throughout the world, evident with many studies across the world. However, the occurrence rate of multidrug resistant *Staphylococcus aureus* infections can vary from country to country and between hospitals; and it also varies between different units of the same hospital and also varies in prevalence depending on geographical area and the socio-demographic characteristics of the populations (Kim *et al.*, 2018).

From this study, we observed that the occurrence of *S. aureus* isolated from selected fomites was 20.0% and less than 33.6% reported by Onanuga and Awhowho (2016) isolated from fomites in Yenagoa. The occurrence of *S. aureus* isolates from some formite samples in this study were observed to be higher 32.0%, 26.0% and 24.0% from Seat handles, Stretchers and door handles swabs respectively than 1.8%, 21.1%, 0.9% for the different specimens respectively as reported by Oyepola *et al.* (2015) from fomites in Southwest Nigeria. The isolation of *S. aureus* from the fomite samples of door handles, stretchers and seat handles swabs of fomites suggested that the organism may likely be responsible for most hospital acquired infections, since *S. aureus* has been reported as one of the bacteria associated with hospital infections (Kasper *et al.*, 2015). However, the occurrence of *S. aureus* 20.0% from fomite swabs was in close agreement with 22.1% from studies reported by Oyepola *et al.* (2015) and also in agreement that *S. aureus* is also a pathogen associated with wounds infections (Kasper *et al.*, 2015).

The antimicrobial susceptibility testing of the isolates was carried out using Kirby-Bauer disk diffusion method as described in Clinical and Laboratory Standards Institute (CLSI) Guidelines (CLSI, 2012). The following antibiotic resistance was recorded; Streptomycin had 80% which was the highest, Penicillin had 76%, Rifampicin had 72%, Ampicillin/sulbactam had 64%, Clindamycin had 64%, Oxacillin had 62%, Ceftriaxone had 58%, Vancomycin had 28%, Levofloxacin had 26% and Gentamycin had 22% which is the lowest.

### Conclusion

The occurrence of *S.aureus* isolates in the selected fomites in this study was high. The *S.aureus* isolates were more resistant to antibiotics such as streptomycin, clindamycin, rifampicin, oxacillin, ceftriaxone, penicillin and ampicillin/sulbactam, but less resistant to antibiotics such as levofloxacin and gentamicin. These antibiotics of higher susceptibility will be useful in treatment of infections caused by *S.aureus*.

### References

- Acar, J. F., and Moulin, G. (2012). Antimicrobial resistance: a complex issue. *Revue Scientifique et Technique (Paris)*, 31: 23–31.
- Al-haddad, O.H., Zorgani, A. and Ghenghesh, K.S. (2014). Nasal carriage of multidrug resistant panton-valentine leucocidin-positive methicillin-resistant *Staphylococcus aureus* in children in Tripoli-Libya. *American Journal of Tropical Medicine and Hygiene*, 90: 724-727.
- Alli, O.A., Ogbolu, D.O., Shittu, A.O., Okorie, A.N., Akinola, J.O. and Daniel, J.B. (2015). Association of virulence genes with *mec A* gene in *Staphylococcus* isolates from Tertiary Hospitals in Nigeria. *Indian Journal of Pathology Microbiology*, 58:464-471.
- Ba, X., Harrison, E.M., Edwards, G.F., Holden, M.T.G., Larsen, A.R., Petersen, A., Skov, R.L., Peacock, S.J., Parkhill, J., Paterson, G.K. and Holmes, M.A. (2014). Novel mutations in penicillin-binding protein genes in clinical *Staphylococcus aureus* isolates that are

**Comment [ks5]: ISOLATION AND ANTIBIOTIC RESISTANCE OF STAPHYLOCOCCUS AUREUS ISOLATED FROM NOSOCOMIAL SOURCES**

better check the concordance with the theme of the article and improve the conclusion.

**Comment [ks6]:** please add numerical order.  
May be:  
1,2,3,4,.....

methicillin resistant on susceptibility testing but lack the mec gene. *Journal of Antimicrobial Chemotherapy*, 69: 594-597.

Bangerter, P.D., Sidler, X., Perreten, V., Overesch, G., (2016). Longitudinal study on the colonisation and transmission of methicillin-resistant *Staphylococcus aureus* in pig farms. *Veterinary Microbiology*, 183:125-134.

Bannerman, T. L. (2003). Staphylococci, Micrococcus and other catalase positive cocci that grow aerobically. In: Murray, P.R., Baron, E.J. and Jorgensen, J.M. (Eds), *Manual of Clinical Microbiology*, 8th edition. American Society for Microbiology Press, Washington, 384-404.

Boyce, J. M. (1997). Vancomycin-resistant enterococcus: detection, epidemiology, and control measures. *Infect Dis Clin North Am*, 11:367–384.

Boyle-Vavra, S. and Daum, R.S. (2007). Community acquired methicillin resistant *Staphylococcus aureus*: the role of Panton Valentine Leukocidin. *Laboratory invest*, 87>3-9.

Brown, D. F. (2001). Detection of methicillin/oxacillin resistance in Staphylococci. *Journal of Antimicrobial and Chemotherapy*, 48: 65- 70.

Brown, D.F., Edwards, D.I., Hawkey. P.M., Morrison, D., Ridgway, G.L., Towner, K.J. and Wren, M.W. (2005). Guidelines for the laboratory diagnosis and susceptibility testing of methicillin-resistant *Staphylococcus aureus* (MRSA). *Journal of Antimicrobial Chemotherapy*, 56(6):1000-18.

- Cain C. L, (2013). Antimicrobial resistance in staphylococci in small animals. *Veterinary Clinical North America Small Animal Practice*,43(1):19-40.
- Cennet, R., Mehmet, P, Yasemin, B., Huseyin, G. and Nesrin, C. (2016). Evaluation of Antimicrobial Resistance in *Staphylococcus aureus* Isolates by Years. *Interdisciplinary Perspectives on Infectious Diseases*, 27:20-24.
- Cheesbrough, M. (2006).Biochemical testing of microorganism.*Medical Laboratory Manual forTropical Countries*. Vol. II.
- Cheesebrough, M. (2006). Medical Laboratory Manual for Tropical Countries. Cambridge : Cambridge University Press, Pp. 49-97.
- Christopher, A.J., Hora, S. and Ali, Z. (2013). Investigation of plasmid profile antibiotic susceptibility pattern multiple antibiotic resistance index calculation of *Escherichia coli* isolates obtained from different human clinical specimens at tertiary care hospital in Barcilla-India. *Annals of Tropical Medicine and Public Health*, 6:285-289.
- Clinical and Laboratory Standards Institute (2012).Performance Standards for Antimicrobial Susceptibility Testing. CLSI 22nd Informational Supplement M100-S22. Wayne, PA, USA.
- CLSI. (2009). Analysis and presentation of cumulative antimicrobial susceptibility test data. 3rd edition. Approved guideline M39-A3. Wayne, PA, USA.
- CLSI. (2013). Perfomance standards for antimicrobial susceptibility testing. CLSI document M100-S23. Wayne, PA, USA

CLSI (2014). Performance Standards for Antimicrobial Susceptibility Testing: 24<sup>th</sup> Informational Supplement M100-S24. Wagne, PA: Clinical and Laboratory Standards Institute.

Conceição, T., Coelho, C., Santos-Silva, I., de Lencastre, H. and Aires-de-Sousa, M. (2014). Epidemiology of methicillin-resistant and -susceptible *Staphylococcus aureus* in Luanda, Angola: first description of the spread of the MRSA ST5-IVa clone in the African continent. *Microbial Drug Resistance*, 20(5):441-9.

Dalpe, A. H., Hofko, M. and Zimmerman, S. (2012). Comparison of the BD MAX methicillin resistant *Staphylococcus aureus* (MRSA) and the BD GeneOhm MRSA achromopeptidase assay with direct- and enriched-culture techniques using clinical specimens for detection of MRSA. *Journal of Clinical Microbiology*, 50: 3365-3367.

Daniyan, S.Y., Galadima, M., Ijah, U.J. and Oda, M. (2011): *In vitro* susceptibility profile of Methicillin-resistant *Staphylococcus aureus* isolates from clinical specimens to commonly used Antibiotics in Minna, Nigeria. *Asian Journal of Pharmaceutical Health Sciences*, 1(3): 128-129.

Daum, Robert S. "Skin and soft-tissue infections caused by methicillin-resistant *Staphylococcus aureus*." *New England Journal of Medicine*, Vol. 357, No. 4, 2007, pp. 380-90.

David, M. Z. and Daum, R. S. (2017). Treatment of *Staphylococcus aureus* infections, *Current Top Microbiology Immunology*, 409:325-383.

Dossaji, S., Çelik, Ü., Alhan, E., Yıldızdaş, D. and Ünal, İ. (2008). Infection Markers for Nosocomial Infections. *J PediatrInf*, 2: 12-18.

Dyer, O. (2007). New MRSA strain is not at epidemic level, expert says. *BMJ* 334(7583):10.

Elhassan, M.M., Hani, A.O., Hassan, A.H., Miskelyemen A.E., and Leila, M.A. (2015). Absence of the *mec A* Gene in Methicillin Resistant *Staphylococcus aureus* Isolated from Different Clinical Specimens in Shendi City , Sudan. *BioMed ResearchInternational*, 10:89.

En.wikipedia.org.

Ezeamagu, C., Imanatue, I., Dosunmu, M., Odeseye, A., Baysah, G., Aina, D., Odutayo, F. and Mensah-Agyei, G. (2018). Detection of methicillin resistant and toxin-associated genes in *Staphylococcus aureus*. *Beni-Suef University Journal of Basic and Applied Sciences*, 7:92-97.

Fayomi, O.D., Oyediran, E.I., Adeyemo, A.T., Oyekale, O.T. (2011). Resistance pattern of methicillin-resistance *Staphylococcus aureus* among in-patients at a tertiary health facility in Ido-Ekiti, Nigeria. *Internet Journal Laboratory Medicine*, 4:1 -5.

Ferreira, J. P., Anderson, K. L., Correa, M. T., Lyman, R., Ruffin, F., Reller, L. B., Fowler, V. G. Jr. (2011). Transmission of MRSA between companion animals and infected human patients presenting to outpatient medical care facilities. *PLoS One*, 6: e26978.

Gade, N. D. and Qazi, M. S. (2013). Fluoroquinolone therapy in *Staphylococcus aureus* infections: where do we stand? *Journal of Laboratory Physicians*, 5: 109-112.

Garba, S., Onaolapo, J. A. and Olayinka, B.O. (2017). Antimicrobial susceptibility pattern of *Staphylococcus aureus* from clinical isolates in Zaria Metropolis, Kaduna. *Nigerian Journal of Pharmaceutical and Biomedical Research*, 2(116-120).

- Hamid, S., Bhat, M.A., Mir, I.A., Taku, A., Badroo, G.A., Nazki, S., Malik, A. (2017). Phenotypic and genotypic characterization of methicillin-resistant *Staphylococcus aureus* from bovine mastitis, *Veterinary World*, 10(3): 363367.
- Hare, K. T., Darshan, S. and Malaya, R. S. (2008). High prevalence of multidrug-resistant MRSA in a tertiary care hospital of northern India. *Infection and Drug Resistance*, 1: 57–61.
- Hawraa, W.A., Al-Dulaimi, T. and Al-Marzoqi, A.H. (2014) “Phenotypic detection of resistance in *Staphylococcus aureus* isolates: detection of (*mecA* and *fem A*) gene in methicillin resistant *Staphylococcus aureus* (MRSA) by polymerase chain reaction,” *Journal of Natural Sciences Research*, 4(1):112–118
- Hong, S., Yingmei, L., Jiuxin, Q. and Bin, C. (2016). Comparison of vanA gene mRNA levels between vancomycin-resistant Enterococci presenting the VanA or VanB phenotype with identical Tn1546-like elements. *Journal of Microbiology, Immunology and Infection*, 49 (6): 866-871.
- Ibadin, E.E., Enabulele, I.O. and Muinah, F. (2017).Prevalence of *mecA* gene among staphylococci from clinical samples of a tertiary hospital in Benin City, Nigeria. *African Health Sciences*, 4:1002-1003.
- Jarvis, W.R., Jarvis, A.A. and Chinn, R.Y.(2010).“Nationalprevalence of methicillin-resistant *Staphylococcus aureus* in inpatients at UnitedStateshealthcare facilities,”*TheAmerican Journal of Infection Control*, 3:194–200.

- Jin, M., Rosario, W., Watler, E. and Calhoun, D.H. (2004). Development of a largescale HPLC-based purification for the urease from *Staphylococcus leei* and determination of subunit structure. *Protein Experimental Purification*, 34: 111-117.
- Kapatamoyo, B., Andrews, B. and Bowa, K. (2010). Association of HIV with breast abscess and altered microbial susceptibility patterns. *Medical Journal of Zambia*, 37: 58-63.
- Kaspar, U., Kriegeskorte, A., Schubert, T., Peters, G., Rudack, C., and Pieper, D.H. (2015). The Culturome of the Human Nose Habitats Reveals Individual Bacterial Fingerprint Patterns. *Environmental Microbiology*, 18(7):2130-42
- Kaur, D.C., and Chate, S.S. (2015). Study of Antibiotic Resistance Pattern in Methicillin Resistant *Staphylococcus aureus* with Special Reference to Newer Antibiotics. *Journal of Global Infectious Diseases*, 7(2):78-84.
- Kim, J. W., Chung, G. T., Yoo, S. J., Lee, Y. S. and Yoo, I. J. (2012). Autolytic activity and molecular characteristics of *Staphylococcus haemolyticus* strains with induced vancomycin resistance. *Journal of Medical Microbiology*, 61; 1428–1434.
- Kim, M.W., Greenfield, B.K., Snyder, R.E., Steinmaus, C.M., and Riley, L.W. (2018). The association between community associated *Staphylococcus aureus* colonization and disease: a meta-analysis. *BMC Infectious Diseases*, 18:86
- Oyepola, O.O., Olasupo, N.A., Egwari, L.O., Becker, K., and Schaumburg, F. (2015). Molecular Characterization and Antimicrobial Susceptibility of *Staphylococcus aureus* Isolates from Clinical Infection and Asymptomatic Carriers in Southwest Nigeria. *PloS One*, 10(9):e0137531

Paterson, D. L., Ko, W. C. and Mohapatra, S. (1997). *Klebsiella pneumonia* bacteremia: impact of extended spectrum beta-lactamase (ESBL) production in a global study of 216 patients [abstract]. 37th Interscience Conference on Antimicrobial Agents and Chemotherapy; Sept 28 –Oct 1, 1997; Toronto, Ontario, Canada

UNDER PEER REVIEW