

Original Research Article

Gastrointestinal Parasites of Wild *Thryonomys swinderianus* (Grasscutters) Sold at Eateries in Umuawulu Community, Awka South Local Government Area, Anambra State, Nigeria.

Abstract

Parasitic infections have been shown to affect the growth and economic value of animals. The demand for grasscutter meat is high in most rural and semi-urban communities in Nigeria due to its value as a source of protein. This study investigated the gastrointestinal parasite composition of wild grasscutters (*Thryonomys swinderianus*) sold in restaurants within Umuawulu, Anambra State. Three locations were randomly sampled within the study area and the grasscutters were purchased from three eateries in the sampled locations. The gut and intestinal contents of the animals were collected at the point of slaughter, transferred to a sterilised container and transported to Zoology Laboratory, Nnamdi Azikiwe University, Awka for further analysis. A total of 45 (25 male and 20 female) grasscutters were examined out of which 23(51.1%) were infected. The parasites identified were: *Ancylostoma* sp. 20 (75.0%) and *Ascaris* sp. 5 (55.56%), *Necator americanus* 2 (25.00%) and *Trichuris* sp. (12.50%). The highest parasite prevalence in the grasscutters sampled was found in females 13 (65.0%) while the male has the least 10 (40.0%). Parasite prevalence was not sex-dependent ($\chi^2= 0.95$, $P=0.136$). Also, there is no significant relationship between the parasite distribution and location ($\chi^2= 3.889$, $P>0.05$). It is recommended that further studies should be carried out in other communities since the high prevalence of helminth infection may compromise the productivity of the grasscutters if infected.

Keywords: Gastrointestinal Parasites, Grasscutters, Umuawulu Community, Anambra State, and Nigeria.

I. Introduction

Wildlife has been an important protein source for many West African countries over centuries. The population of these animals (duikers, red Columbus monkeys, chimpanzees and grasscutters etc.) are now dwindling and is inadequate to provide the protein needs of the people of West Africa due to over-dependence in the past years [1]. The meat is eaten locally and it is also exported for income among countries in West Africa [2, 3]. In some developing countries in

Comment [A1]: The present study

Comment [A2]: prevalence of gastrointestinal parasites

Comment [A3]: to

Comment [A4]: had

Comment [A5]: was

Comment [A6]: It thus highlighted the need for screening of parasites in stock grasscutters brought from the wild for domestication as well as properly cooked before consumption. Further studies are required generate more data on it.

Comment [A7]: consumed

Africa, bush meat is the main source of animal protein among rural communities and the demand for it is rapidly increasing irrespective of the fact that zoonotic infections such as Ebola and COVID-19 are associated with the consumption of meat from wild animals [4,5].

Nigeria is well-known for its consumption of bush meat [2]. More than 80% of Nigerians in both rural and urban areas prefer bush meat and would eat it if available [6]. Bush meat is popular and important in most traditional festivities among African communities [7, 8] and among the wild animals is the grasscutter which has gained wider interest among African countries [9] including Nigeria [10].

Comment [A8]: Grasscutters (*Thryonomys swinderianus*)

Most hunters sell grasscutter in the raw form in local markets and on highways though the processed form has better prices. Lack of hygiene and handling practices of processed meat can potentially create a negative impression about the grasscutter industry in Nigeria [11].

Domestication of grasscutters is a relatively novel practice in Nigeria with its potentialities, challenges and opportunities [12]. Grasscutter farming has social acceptability, good meat quality of high biological value (high protein and low fat), inexpensive feeds and responds to captive rearing, good litter size and short generation interval. Despite the challenges of domestication of grasscutter, non-steady supply of meat, air pollution and ecological devastation as a result of bush burning to hunt grasscutter and the threat to the extinction of grasscutter; the domestication and production of grasscutter is another dimension in the livestock industry that has the potential to ensure regular and sustainable animal production in the nation [12, 13].

With the current trend in grasscutter farming and the intensification of management practices, there is bound to be an increased incidence of infections and other challenges. Some challenges to grasscutter production include the irregular supply of breeding stock, environmental issues, poor processing and marketing plan, lack of balanced diet, poor producer training and education, inadequate infrastructural development, poor information dissemination, incidences of diseases and mortality [14]. Grasscutters can transmit parasitic diseases with significant public health and economic consequences [2]. They are capable of transmitting some of the zoonotic gastrointestinal parasites, especially if the meat is not properly cooked [15]. It has been reported [16] that some parasites responsible for zoonoses and communicable diseases to man and

animals such as nematodes (*Ascaris*, *Bunostomum*, *Oesophagostomum*, *Strongyloides*, *Trichostrongylus*, *Toxocara*); trematodes (*Fasciola*, *Schistosoma*); Cestodes (*Taenia*) and acanthocephalan (*Moniliformis*) occur in the grasscutter and their public health significance is worthy of note.

Also, the devastating effect of parasites on grasscutters was brought to notice with the study of [17]), the post-mortem results revealed heavy infection and the prevalence of intestinal helminths in grasscutters from Omagwa district in River State, Nigeria. In this study, the result revealed the presence of seven different species of intestinal helminths (*Ascaris* sp., *Strongyloides* sp., *Trichuris* sp., *Fasciola* sp., *Taenia* sp., *Moniliformis* sp. and Hookworm). These pieces of evidence show that the health of captive grasscutters can thus no longer be taken for granted. Several parasites in wild grasscutters were reported in Oyo State, southwestern Nigeria [18]. In Umuahia, Imo State, Nigeria, there was a 37.2% prevalence of gastrointestinal parasites in grasscutters. Male and female grasscutters had 30.7% and 52.9% parasite infections respectively [19].

Due to the increased demand for quality grasscutter meat in Nigeria [20] and the patronage for grasscutter meat consumption became high in sub-Saharan Africa [21]. The African population growth rate with the lowest protein supply in the world [2] also, calls for more attention to the regular supply of grasscutter meat as a possible alternative to pork, chevon, beef and chicken since there are no religious prohibitions for the consumption of grasscutter meat [22, 23]. However, research and literature have shown that the availability of this valued product is affected by many factors such as parasitic diseases (protozoa and helminths) [24, 25]. The frequency dependence of humans on bush meat as a vital resource poses increased risks of zoonotic pathogens and disease transmission [26] through its handling and consumption [27, 28].

As promising as micro-livestock integration sounds, parasites in grasscutters will threaten the practicability of production and availability of meat to the general public. It is against this background that this study is designed to find out the intestinal parasites of grasscutters sold at eateries in Umuawulu, Anambra State. The objectives of the study are to identify and determine

Comment [A9]: a

the prevalence of intestinal parasites of grasscutters sold at popular eatery joints in the Umuawulu community in Anambra State, Nigeria.

2. Materials and Methods

Area of Study

The study was carried out in Umuawulu, Awka South local government area, Anambra State, Nigeria. The town is located at Latitude: 6° 08' 60.00" N and Longitude: 7° 05' 60.00" E. The estimated population by 2005 is 20,150 living in three villages: Enugwu, Umuenu and Agbana. The study was carried out in three eateries popularly known for the **sailing** of grasscutter meat.

Comment [A10]: Do not bold the letter

Comment [A11]: selling

Study design

The study is a survey aimed at identifying and determining the prevalence of the gastrointestinal parasites of grasscutters.

Ethical clearance certificate

Permission and ethical clearance certificate were obtained from Nnamdi Azikiwe University Animal Research Ethical Committee with reference number NAU/AREC/2021/00057.

Sample Collection and Analysis

The study was carried out from October to December 2021. The study site was visited two times a month; 4 grams of faecal matter were collected from the rectum of the animals before they were sliced by the local chefs. Samples were kept in coolers containing ice packs and were taken to the Zoology Laboratory of Nnamdi Azikiwe University, Awka, Anambra State, Nigeria where they were analysed. A total of forty-five (45) faecal samples were collected from the three different eateries in three locations and were given a unique code number (UET1 – Umuawulu eatery number 1, UET2 - Umuawulu eatery number 2 and UET3 - Umuawulu eatery number 3) for easy identification.

The qualitative faecal examination was done using the Cornell-McMaster dilution egg counting technique [29]. Two grams of faecal samples were measured into a cup and mixed with 28 ml **flotation solution** and allowed to soak for 5 minutes. After 5 minutes the solution is steered and strained into a second cup. With the use of a syringe, McMaster slide chambers were filled with the mixture and set aside for at least 5 minutes to allow parasite eggs to float to the surface. In less than an hour, all eggs inside the grid areas were counted using the **X10** objective lens of the Microscope.

Comment [A12]: If possible elaborate composition

Comment [A13]: 10X

The prevalence of intestinal parasites was determined using the Chi-square (χ^2) test from the contingency tables while Fisher's exact test was used to calculate the significant difference in prevalence for gender, location and parasite species. Analysis was done using SPSS (version 20.0) statistical software. The significant difference was set at $P = 0.05$ and all tests were performed with a 95 % confidence level.

3. Results

The overall prevalence of gastrointestinal parasites in the animals was 51.11%. Table 1 revealed that the grasscutters sold at eateries in Umuawulu had the highest number of *Ancylostoma* sp, 15(75.00%) followed by *Ascaris* sp, 5(55.56%) and *Trichuris* sp. 1(12.50%) recorded the lowest. The prevalence rate was significantly different ($P>0.05$) with species of parasite.

Comment [A14]: Begin result section by stating what is mentioned in table 1, table 2 and table 3.

Comment [A15]: What about *Necator americanus*?

Comment [A16]: Use less than greater sign as per significant or non-significant. For significant results $P<0.05$

Table 1: Prevalence of Gastrointestinal Parasites of Grasscutters in Relation to Parasite Species

Name of Parasite	No. of grasscutters examined	Infected	Prevalence (%)
<i>Ancylostomas</i> p.	20	15	75.00
<i>Ascaris</i> sp.	9	5	55.56
<i>Necatoramericanus</i>	8	2	25.00
<i>Trichuris</i> sp.	8	1	12.50
Total	45	23	51.11

$\chi^2 = 11.595$, $df = 3$, $P = 0.01$

Comment [A17]: These can be mentioned in the columns that can be added on the right side of table.

Table 2 revealed that the female grasscutters had a higher number of infections, 13(65.00%) than the males 10(40.00%). The prevalence rate of infection was not significantly different ($P>0.05$) with gender.

Table 2: Prevalence of Gastrointestinal Parasites of Grasscutters in Relation to Gender

Gender	No. examined	Infected	Prevalence (%)
--------	--------------	----------	----------------

Male	25	10	40.00
Female	20	13	65.00
Total	45	23	51.11

$\chi^2 = 2.779, df = 1, P = 0.09$

Comment [A18]: These can be mentioned in the columns that can be added on the right side of table.

In Table 3 it can be seen that the grasscutters sold at UET1 and UET3 had a higher number of infections, 9(60.00%) than those of UET2 5(33.33%), however, there was no significant difference ($P > 0.05$) with location.

Comment [A19]: Table 3,

Comment [A20]: non

Table 3: Prevalence of Gastrointestinal Parasites of Grasscutters in Relation to Location

Location	No. of grasscutters		Prevalence (%)
	Examined	No Infected	
UET1	15	9	60.00
UET2	15	5	33.33
UET3	15	9	60.00
Total	45	23	51.11

$\chi^2 = 2.846, df = 2, P = 0.241$

Comment [A21]: These can be mentioned in the columns that can be added on the right side of table.

4. Discussion

The results revealed that grasscutters from the wild are heavily infected with intestinal parasites that can be transmitted to their offspring and other mammals, including humans that may come into contact with them if eaten poorly prepared or domesticated. Two types of hookworm were isolated from the grasscutter and *Ancylostoma* sp. has the highest prevalence. This is similar to the study of Futagbi *et al.* [30] who reported a high prevalence of *Ancylostoma* sp. in grasscutter which may have high medical significance. *Ancylostoma* sp. causes chronic loss of blood in animals including humans. Hookworm diseases such as Ancylostomiasis and Necatoriasis are widely endemic in tropical and subtropical countries where sanitary disposal of human feces is low and the soil moisture and temperature conditions favour the development of infective larvae

Comment [A22]: Please check the style of citations from journal format and maintain uniformity throughout manuscript.

[31]. At least 68 hookworm species have been described in 9 orders, 24 families, and 111 species of wild mammals. Black bears, red foxes, and bobcats harboured the highest diversity of hookworm species. *A. pluridentatum*, *A. tubaeforme*, *Uncinariastenocephala* and *Necator americanus* were the hookworm species with the highest host diversity index. Hookworm infections cause anaemia, retarded growth, tissue damage, inflammation and significant mortality in several wildlife species [32]. Wildlife infections with *Necator americanus* are common in many regions of the world [33].

Another parasite in the study with a relatively high prevalence is *Ascaris* sp. In a study by Opara and Fagbemi reported by [16], 14 species of nematodes including *Ascaris* sp. were reported in Imo State, Nigeria. *Trichuris* sp. prevalence in the grasscutters was also moderately high as it is in this study in Anambra State.

An improvement in sanitation in the study area is very important because of the relatively high number of hookworm parasites identified in this region.

Comment [A23]: Reduce gap between two lines

5.2 Conclusion and Recommendations

Ancylostoma sp. was the most prevalent species among the parasites found in the study. It is recommended that stock grasscutters brought from the wild for domestication should be screened for parasites before they are prepared for consumption. Further, the meat should be properly cooked to avoid ingestion of helminth eggs.

Comment [A24]: After first sentence of conclusion name of other parasites can be mentioned in their decreasing order of prevalence.

REFERENCES

1. Aguirre, A. A. (2017). Changing patterns of emerging zoonotic diseases in wildlife, domestic animals, and humans are linked to biodiversity loss and globalization. *ILAR International League against Rheumatism*, 58(3), 315-318
2. Adewumi, A., Famubo, E., Ofuya, E., Wahab, M. (2021). Economic Analysis of Grass Cutter Raised in Captivity in Osun and Oyo State, Nigeria. *Scientific Reports in Life Sciences*, 2(3), 1-7.
3. Tokede, A. M. (2020). Cultural Beliefs on Grasscutter Consumption and Willingness to Adopt its Domestication Technology among Urban Dwellers in Southwest, Nigeria. *Nigeria Agricultural Journal*, 51(3), 10-15.
4. Tiwari, R., Dhama, K., Sharun, K., Yattoo, M. I., Malik, Y. S., Singh, R., Michalak, I., Sah, R., Bonilla-Aldana, D. K., & Rodriguez-Morales, A. J. (2020). COVID-19:

- Animals, veterinary and zoonotic links. *The Veterinary Quarterly*, 40(1), 169-182. <https://doi.org/10.1080/01652176.2020.1766725>
5. Available: https://www.who.int/health-topics/ebola/#tab=tab_1 Ebola virus disease
 6. Aiyelaja, A. A., and Ogunjinmi, A. (2013). Economic aspects of grasscutter farming in southwest Nigeria: Implications for sustainable adoption and conservation. *International Journal of Scientific and Engineering Research (IJSER)*, 4(10), 17-23
 7. Blankson-Arthur, S. (2009). Levels of organochlorine pesticide residue in grasscutter (*Thryonomys swinderianus*) tissues. Bibliographic information available from INIS: http://inis.iaea.org/search/search.aspx?orig_q=RN:41058575; Available from the University of Ghana, School of Nuclear and Allied Sciences, Department of Nuclear and Environmental Protection, P. O. Box AE 1, Legon, Accra, Ghana
 8. Luiselli, L., Hema, E. M., Segniagbeto, G. H., Ouattara, V. A. L. Y., Eniang, E. A., Parfait, G., and Fa, J. E. (2020). Bush meat consumption in large urban centres in West Africa. *Oryx*, 54(5), 731-734
 9. Odunlami, S. S., and Nkata, J. E. (2021). Contributions of grasscutter (*Thryonomys swinderianus*) hunting to the livelihood of communities in Abi local government area, Cross River state, Nigeria. *Journal of Research in Forestry, Wildlife and Environment*, 13(1), 26-35.
 10. Adu E K, Asafu-Adjaye A, Hagan B A and Nyameasem J K 2017: The grasscutter: an untapped resource of Africa's grasslands. *Livestock Research for Rural Development*. Volume 29, Article #47. Retrieved June 28, 2023, from <http://www.lrrd.org/lrrd29/3/jnya29047.html>
 11. Uka, U. U., Mbanasor, J. A., Nto, P. O., and Aigbokie, S. O. (2021). Determinants of Profitability of Grasscutter Farming in South East Nigeria. *JCCR/ Journal of Community and Communication Research*, 6(1), 69-75.
 12. Owen, O. J., and Dike, U. A. (2012). Grasscutter (*Thryonomys swinderianus*) husbandry in Nigeria: A review of the potentialities, opportunities and challenges. *Journal of Environmental Issues and Agriculture in Developing Countries*, 4(1), 104-111.
 13. Okanlawon, Oluwatoyin Mayowa, Adeleke, Oluwaseun Aderonke and Eniola, Phillip Olanrewaju. (2019). Factors influencing Grasscutter (*Thryonomys swinderianus*) Domestication in Rural Communities of Oyo State, Nigeria. *Journal of Agricultural Extension*. 23, 1 DOI: 10.4314/jae.v23i1.3
 14. Akinola, Letorn ,Etela, Ibisime&Emiero, S. (2015). Grasscutter (*Thryonomys swinderianus*) Production in West Africa: Prospects, Challenges and Role in Disease Transmission. *American Journal of Experimental Agriculture*, 6, 196-207. 10.9734/AJEA/2015/14194.
 15. Nwoke, B.E.B. (2001): Urbanization and livestock handling and farming: The public health and parasitological implications. *Nigeria Journal of Parasitology*, 22, 121-128
 16. Opara, Maxwell. (2012). Zoonotic Role of the Grasscutter. Available from: https://www.researchgate.net/publication/224829662_Zoonotic_Role_of_the_Grasscutter
 17. Abara, P.N., Adjero, L. A., Nwachukwu, M.O., Osinomumu, I.D. (2021) Preliminary survey of the intestinal helminths of greater cane rat and antelope (bush meat) in Omagwa Rivers State. *IAA J. Appl. Sci.* 7(1), 49–56.
 18. Okorafor, K. A., Okete, J. A., Andem, A. B., and Eleng, I. E. (2012). Assessment of grasscutters (*Thryonomys Swinderianus*) sellers and hunter's conservation knowledge,

Comment [A25]: Insert gap between two words. Check if there is different version of word document.

rate of hunting and methods of hunting in Oyo State, Nigeria. *European Journal of Zoological Research*, 1(4), 86-92.

19. Onyeabor, A. (2015). Prevalence of Gastrointestinal Helminth Parasites in Grasscutter in Umuahia Area of Abia State. *Journal of Veterinary Advances*, 5, 814-818.
20. Timothy, J., Anthony, A., and Ochani, A. N. (2020). Skills Required by Agricultural Education Graduates in Grass Cutter (*Thryonomys swinderianus*) Farming for Self-Employment in Kaduna State, Nigeria. *Journal Agriculture Sciences and Food Resources*, 11, 275.
21. Opara, M. N., and Fagbemi, B. O. (2008). Occurrence and prevalence of gastrointestinal helminths in the wild grasscutter (*Thryonomys swinderianus*, Temminck) from Southeast Nigeria. *Life Science Journal*, 5(3) 50-56.
22. Odebode, A. V., Awe, F., Famuyide, O. O., Adebayo, O., Ojo, O. B., and Daniel, G. (2011). Households' consumption patterns of grasscutter (*Thryonomys swinderianus*) meat within Ibadan Metropolis, Oyo State, Nigeria. *Continental Journal of Food Science and Technology*, 5(2), 49-57.
23. Ekwochi, U., Osuorah, C.D.I., Ndu, I.K. *et al.* Food taboos and myths in South Eastern Nigeria: The belief and practice of mothers in the region. *J. Ethnobiology Ethnomedicine*, 12, 7 (2016). <https://doi.org/10.1186/s13002-016-0079-x>
24. Opara, M.N. (2010). Grasscutter: The Haematology and Major Parasites. *Research Journal of Parasitology*, 5, 214-223
25. Robinson, M. W., & Dalton, J. P. (2009). Zoonotic helminth infections with particular emphasis on fasciolosis and other trematodiasis. *Philosophical Transactions of the Royal Society of London. Series B, Biological sciences*, 364(1530), 2763–2776. <https://doi.org/10.1098/rstb.2009.0089>
26. Friant S, Paige SB, Goldberg TL (2015) Drivers of Bushmeat Hunting and Perceptions of Zoonoses in Nigerian Hunting Communities. *PLOS Neglected Tropical Diseases* 9(5), e0003792. <https://doi.org/10.1371/journal.pntd.0003792>
27. FAO, IFAD and WFP. 2015. The State of Food Insecurity in the World 2015. Meeting the 2015 international hunger targets: taking stock of uneven progress. Rome, FAO
28. Wiafe, Edward. (2014). Consumer Views of Bushmeat Consumption in Two Ghanaian Markets. *Applied Research Journal*, 2014 (1), 20-27.
29. Available: <https://wcv.m.usask.ca/learnaboutparasites/diagnostics/quantitative-faecal-flotation-mcmaster.php> Quantitative Faecal Flotation - McMaster Egg Counting Technique.
30. Futagbi, G., Agyei, D. O., Aboagye, I. F., Yirenya-Tawiah, D. R., and Edoh, D. A. (2010). Intestinal parasites of the grasscutter (*Thryonomys swinderianus* Temminck 1827) from the Kwaebibirem District of the Eastern Region of Ghana. *West African Journal of Applied Ecology*, 17(1), 1-6.
31. Alemayehu. M. (2004). Communicable Diseases Control: Ethiopia Public Health Training Initiative (EPHTI). Produced in collaboration with the Ethiopia Public Health Training Initiative, The Carter Centre, the Ethiopia Ministry of Health, and the Ethiopia Ministry of Education. Pp 53
32. Seguel M, Gottdenker N. The diversity and impact of hookworm infections in wildlife. *Int J Parasitol Parasites Wildl.* 2017 Apr 4;6(3):177-194. doi: 10.1016/j.ijppaw.2017.03.007.

33. Wiethoelter, Anke, Beltran-Alcrudo, Daniel, Kock, Richard & Mor, Siobhan. (2015). Global trends in infectious diseases at the wildlife–livestock interface. *Proceedings of the National Academy of Sciences of the United States of America*. 112. 10.1073/pnas.1422741112.

Comment [A26]: Please check the format of references as required by the journal and maintain uniformity in the manuscript. In some references abbreviated form of journal name is used whereas at other places full form is used. Recheck formatting as per guidelines of the journal.

UNDER PEER REVIEW