

### ***Abstract***

Chemical characterization of acid sulphate soils of Upper Kuttanad, Kerala was done to study its present nutrient status due to the impact of heavy flood that occurred in 2018. Extreme acidity is the major issue with this type of soil. Among the collected soil samples, 53.33 per cent of soil samples fall under very strongly acidic group (4.5 to 5) and 33.33 per cent comes under strongly acidic category (5.1 to 5.5). Lower values of EC in UpperKuttanad indicate that this area is less affected by saltwater intrusion. More than 90 per cent of the samples were rich in organic carbon content. Among the major nutrients, 40 per cent of the soil samples recorded low available P content ( $<10\text{kg ha}^{-1}$ ). The availability of potassium, calcium, and magnesium increased with flooding. Almost all the samples collected from the acidsulphate soils of UpperKuttanad showed high available sulphur status. It ranged from 20.20 to 3361.50  $\text{mg kg}^{-1}$ . The hike in micronutrient status showed the impact of flooding in these locations. The cation exchange capacity of the collected soil ranged from 5.92 to 30.64  $\text{cmol (+) kg}^{-1}$  and showed a positive correlation with organic carbon (0.779\*\*\*), available potassium (0.571\*) and available iron (0.717\*). This paper helps in understanding the present nutrient status in acidsulphate rice soil of Upper Kuttanad after the flood and adoption of suitable management practices.

**Keywords :** *Acidsulphate soil, Upper Kuttanad, Flood, Available nutrients, Organic carbon, Micronutrients, Cation exchange capacity*

### **Introduction**

Kuttanad, the low-lying tract in Kerala State of south-west India, is a place where rice cultivation is mainly practised below mean sea level. Kuttanad, "the rice bowl of Kerala covers an area of 874  $\text{km}^2$  distributed with in Alappuzha , Kottayam and Pathanamthitta districts and lying 0.6 to 2.2 m below mean sea level. The area extends from  $9^{\circ}17'$  to  $9^{\circ}40'$  N latitude and  $76^{\circ}19'$  to  $76^{\circ}33'$  E longitude on the west coast of Kerala. Based on the flood incidence, salt water intrusion and acidity, Kuttanad has been divided into 7 agro-ecological zones, viz., Upper Kuttanad, Lower Kuttanad, Kayal lands, North Kuttanad, Coastal Kuttanad, Purakkadkari and Vaikamkari.

Soils in these regions are extremely acidic due to the deposit of pyrites or sulphidic material in the subsurface layers of soil. Due to the deposition of sulphidic materials, Kuttanad is included under the great group sulfaquents and the subgroup TypicSulfaquents. Extreme acidity, high organic matter content and presence of Fe and Al ions in the soil affect plant growth by reducing the uptake of elements like phosphorus. Kerala State receives very high rainfall during the monsoon season. In August 2018, the State received a terrific rainfall of nearly 2346.6 mm against the normal value of 1649.5 mm causing much havoc to entire Kerala except Kasargod district. The soils of Kerala, exposed to this devastating flood in this incessant rainfall experienced great damage to the soil environment in different ways. Besides heavy agricultural losses, soil fertility and productivity have been disturbed, the exposed soils were eroded and various kinds of debris accumulated on farm fields. The top soil in the hills and upland areas have been removed in the flash flood and plantations were completely destroyed. Physical, chemical and biological properties of the soil were highly altered. In this circumstances chemical characterisation of acidsulphate soils of Upper Kuttanad is inevitable and this paper will help in understanding the distribution and availability of macro and micronutrients in Upperkuttanad in post-flood scenario.

**Comment [c2]:** Refernces are missing ,write introduction more so that it brings clarity.

## Materials and Method

The present study entitled "Chemical characterization of acidsulphate rice soils of Upper Kuttanad of Kerala" was carried out at College of Agriculture, KAU during 2019. Acid sulphate soils from Upper Kuttanad regions of Kuttanad were collected during the season, Oct to Jan. A total of fifteen soil samples were collected from the acidsulphate soils of Pavookara, Veeyapuram, Payippad, Karuvatta of Alappuzha and Thiruvarp village of Kottayam district. Details of locations are given in (Table 1).

**Table 1: Details of locations**

SI No.	Block /Panchayath/ village	Name of the padav	Latitude	Longitude
	<b>Pavukara</b>			
1		Ariyodipadashekaram (1)	9.323066133074462	76.52112998999655
2		Ariyodipadashekaram(2)	9.323414065875113	76.52080971747637
3		Ariyodipadashekaram (3)	9.32342127431184	76.52046228758991

4		Panthalattipadam (1)	9.324274216778576	76.51800530031323
5		Panthalattipadam (2)	9.324298272840679	76.51756642386317
	<b>Veeyapuram</b>			
6		Kattakuzhi, Maelppadam (1)	9.325750730931759	76.46919714286923
7		Kattakuzhi, Maelppadam (2)	9.3249811020763516	76.4676930103451
8		Veeyapuram state seed farm (1)	9.322019568644464	76.4618010353297
9		Veeyapuram state seed farm (2)	9.322077655233443	76.46131597459316
	<b>Payippad</b>			
10		Prayatterippadam (1)	9.321848535910249	76.4577947370708
11		Prayatterippadam (2)	9.3216404132545	76.45756758749485
12			9.320900542661548	76.45257147029042
	<b>Karuvatta</b>			
13		Vellookaripadam	9.320897608995438	76.43266185186803
	<b>Thiruvarpu</b>			
14		Muppaykaripadam	9.5893772598356	76.47217825055122
15		Malaraikkal, J block 9000	9.557558675296605	76.49141463451087

Soil sampling were done using a core sampler from 0–20 cm depth without disturbing the reduced condition as far as possible. The collected samples were immediately sealed in a polythene cover and labelled with the location details and additional information, if any. The geographic coordinates of sampling sites were recorded using **GPS**.

**Comment [c3]:** Explain the method in more precise way from starting.

### Characterization of soil samples

Soil samples collected from the paddy fields of Upper Kutanad regions were analysed for estimating pH, EC, OC, Available N, P, K, Ca, Mg, S, Fe, Mn, Cu, Zn and CEC. The procedures adopted for the characterization of soil samples are detailed below.

Soil reaction (pH) was determined in a 1:2.5 soil-water suspension potentiometrically using a pH meter (Jackson, 1958). Electrical conductivity (EC) was estimated in the supernatant liquid of the soil-water suspension (1:2.5) used for pH estimation with the help of a conductivity meter (Jackson, 1958). Organic carbon was estimated by wet digestion method (Walkley and Black). Available nitrogen was determined by alkaline potassium permanganate ( $\text{KMnO}_4$ ) method by Subbiah and Asija (1956). Available phosphorus was extracted using Bray No. 1 extractant (Bray and Kurtz, 1945) which was widely used as an index of available P in acid soil. Available potassium, calcium and magnesium in the soil samples were extracted using neutral normal ammonium acetate. The potassium content in the extract was estimated by flame photometry (Jackson, 1958). Calcium and magnesium content were estimated in an atomic absorption spectrophotometer (Model: Perkin Elmer-PinAAcle 500). Available sulphur was extracted by using 0.15%  $\text{CaCl}_2$  (Tabatabai, 1982) and estimated by turbidimetry (Massoumi and Cornfield, 1963) using a spectrophotometer (Model: Systronics 169). Available micronutrients were analysed using 0.1M HCl extractant and concentrations of micronutrients (Fe, Mn, Zn and Cu) were measured using AAS (Sims and Johnson, 1991). Barium chloride extractable cations were determined by following the procedure of Hendershot and Duquette (1986). The cations present in the exchangeable sites in the soil were replaced by Barium. Estimation of cations in the extract (Ca, Mg, Al, Fe, Mn, Cu, Zn, Na and K) was done using AAS. Since the lowland soils are high in water soluble cations, the deduction of water soluble fraction from barium chloride extractable cations was carried out to avoid over estimation. Water-soluble cations were estimated by the modified procedure of Baruah et al. (2011).

## Result and discussion

Chemical characterisation of soil samples from Upper Kuttanad is given in the Table 2. Soil reaction or pH is the single most informative measurement that can be made to determine soil characteristics. The pH of the collected soil varied from 4.63 to 6.49 with a mean of 5.13. The lowest value, 4.63 was reported by Muppaykaripadam of Thiruvarpu village and highest value, 6.49 was reported by Prayatterippadam of Payeppad. Among the collected soil samples, 53.33 per cent of soil samples fall under very strongly acidic group (4.5 to 5) and 33.33 per cent fall under strongly acidic category (5.1 to 5.5) and 6.67 per cent fall under moderately (5.6 to 6) and slightly acidic category (6.1 to 6.5). Nhung and Ponnamperruma (1966), based on experiments with four acid sulphate soils from Philippines, Vietnam and Malaysia suggested that the chief obstacle to the growth of rice in acid sulphate soils might be their strong acidity because even after months of submergence, the pH of these soils did not rise above 6.0. The accumulation of pyrites ( $\text{FeS}_2$ ) on the subsurface layers of these soil profiles followed by its oxidation results in acidification.

The electrical conductivity of the soil samples varied from 0.04 to 0.97 with a mean value of  $0.26 \text{ dSm}^{-1}$ . Soil samples from the state seed farm, Veeyapurampanchayath reported the lowest EC content and J block padashekaram of Thiruvarpu showed highest value for EC. The electrical conductivity of all the collected soil samples were below  $1.00 \text{ dSm}^{-1}$ . Lower values of EC in Upper Kuttanad indicate that this area is less affected by saltwater intrusion. Even though the salt intrusion is low in Upper Kuttanad, this area is vulnerable to flooding (Thampatti, 1997).

The organic carbon content of the collected soil samples varied from 0.42 to 4.33 per cent with a mean content of 2.35 per cent. Almost all the samples were rich in organic carbon content. Pisharody and Brito-Mutunayagam (1967) observed that organic carbon in rice soils of Kerala varied from 0.03 to 2.86 per cent with the highest in Kuttanad soils. The general trend was a decrease in organic carbon with depth. Tampatti (1997) reported the presence of undecomposed organic matter in Upper Kuttanad, which coincides with a high CN ratio that will go above 10. From the correlation matrix (Table 3) it is very clear that organic carbon showed a strong positive correlation with available potassium ( $0.514^*$ ).

Collected soil samples from Upper Kuttanad showed a variation in available nitrogen content. It varied from 170.72 to 480.45  $\text{kg ha}^{-1}$  with a mean value of  $264.11 \text{ kg ha}^{-1}$ . The lowest value was reported in Ariyodipadashekaram of Pavukara village and highest value was reported by Muppaykaripadam of Thiruvarpu. Almost 73.33 per cent of soil samples showed low available nitrogen content ( $<280 \text{ kg ha}^{-1}$ ) whereas 26.67 per cent showed available

**Comment [c4]:** Correlate your result with more updated research that had already done. Add references along with it.

**Comment [c5]:** Use latest refernces in place of these old ones.

**Comment [c6]:** Try to add more new refernces.

nitrogen in medium range (280-560 kg ha<sup>-1</sup>). Kabeerathumma and Patnaik (1980) reported that acid sulphate soils had moderate to high content of total N (0.095 to 0.37 per cent) and organic carbon (1.4 to 8.4 per cent). The carbon and nitrogen contents are rather high and decrease with depth. In the correlation matrix (Table 3), available nitrogen showed a strong positive correlation with organic carbon content (0.562\*).

The available phosphorus content in acid sulphate soil ranged from 2.14 to 118.29 kg ha<sup>-1</sup> with a mean value of 38.73 kg ha<sup>-1</sup>. Ariyodipadashekaram of Pavukara village showed lowest value for available P whereas Panthalattipadam of Pavukara showed the highest value. Gradation in available P status revealed that almost 40 per cent of the soil samples recorded low available P content (<10 kg ha<sup>-1</sup>), 13.33 per cent of the collected soil samples were in the medium range (10-25 kg ha<sup>-1</sup>) and 46.67 per cent of soil samples recorded high value for available P status (>25 kg ha<sup>-1</sup>). Mathews and Jose (1984) reported a low content of available phosphorus in acid rice soils of Kerala. It was only 4.79 ppm in laterite and 3.84 ppm in *kari* soils. Irene (2014) reported that the available phosphorus in Ponnankole is medium and it is significantly and positively correlated with electrical conductivity. Raychaudhuri and Landey (1960) observed that soils of high clay and silt contents retained more phosphorus and showed low availability to plants. Pareek and Mathur (1969) found higher concentrations of available phosphorus in the surface layers. This was attributed to the distribution pattern of organic matter, pH, sesquioxides and clay content in the soils as well as the requirements of the crop and the rooting depth.

Available potassium ranged from 34.11 to 769.25 kg ha<sup>-1</sup> with a mean value of 215.72 kg ha<sup>-1</sup>. Among the collected soil samples, 33.33 per cent were under low range for available potassium (<120 kg ha<sup>-1</sup>) in which Ariyodipadashekaram of Pavukara showed available K content of 34.11 kg ha<sup>-1</sup>. Though the soils were deficient in available K, they are well supplied with total K and nonexchangeable K (Thampatti, 1997). 40 per cent of soil samples fall under the medium range (120-280 kg ha<sup>-1</sup>) and 26.67 per cent of soil samples fall under high for available potassium (>280 kg ha<sup>-1</sup>). Beena and Manorama (2013) reported high content of potassium in the acid sulphate soils of Kuttanad ranging from 142.1 mg kg<sup>-1</sup> to 326.4 mg kg<sup>-1</sup>. In the correlation study (Table 3) potassium showed a strong positive correlation with available nitrogen (0.671\*\*) and cation exchange capacity (0.571\*).

Secondary nutrients like available calcium, magnesium and sulphur also varied among different locations. Available calcium status ranged from 197.10 to 2581.35 mg kg<sup>-1</sup>

<sup>1</sup>with a mean of 714.76 mgkg<sup>-1</sup>. Among the soil samples collected, Ariyodipadashekaram of Pavukara reported very low available Ca content and Prayatterippadam of Payippad reported very high value for available Ca status. Around 20 per cent of soil samples showed low available Ca content (<300 mgkg<sup>-1</sup>) whereas 80 per cent of the collected soil samples showed high Ca content (>300 mgkg<sup>-1</sup>). Available magnesium content ranged from 39.22 to 372.51 mgkg<sup>-1</sup> with a mean value of 139.23 mgkg<sup>-1</sup>. Ariyodipadashekaram of Pavukara showed low available Mg content and J block of Thiruvapur reported high for available Mg status. More than 50 per cent of soil samples showed low for available Mg (<120 mgkg<sup>-1</sup>) and 46.67 per cent showed high levels of available Mg (>120 mgkg<sup>-1</sup>). According to Irene (2014), under acidic conditions, the exchangeable K, Ca and Mg in solid phase are released into the solution phase. Thus, the bioavailability of these cations is increased upon flooding.

Almost all the samples collected from the acid sulphate soils of Upper Kuttanad showed high value for available S status. It ranged from 20.20 to 3361.50 mgkg<sup>-1</sup> with a mean value of 1257.27 mgkg<sup>-1</sup>. The lowest values were reported by the state seed farm of Veeyapuram and the highest values were reported by Kattakuzhi, Maelpadam of Veeyapuram. Money (1961) reported that the sub-soil clay in the Kari soils of Kuttanad contains high amount of organic matter and are rich in sulphur compounds. Surface soils contain total sulphur (1.92 per cent), subsoil (3.13 per cent) and wood fossils (6.10 per cent).

Among the micronutrients, Fe, Mn, Cu and Zn reported very high values in all the collected soil samples. Kattakuzhi, Maelpadam of Veeyapuram showed very high values for available Fe and Mn. Available Fe ranged from 104.74 to 2817.99 mgkg<sup>-1</sup> with a mean content of 933.70 mgkg<sup>-1</sup>. In the case of available Mn, values ranged from 2.38 to 247.07 mg kg<sup>-1</sup> with a mean value of 44.01mg kg<sup>-1</sup>. Vellookaripadam of Karivatta reported very low value for available Cu and Zn. Available Cu ranged from 2.30 to 35.69 mgkg<sup>-1</sup> with a mean value of 11.86 mgkg<sup>-1</sup>. Available Zn ranged from 1.03 to 21.37 with a mean value of 8.64 mgkg<sup>-1</sup>. Micronutrient status during the pre-flooded situation in Upper kuttanad of Kerala reported very low value, which was in accordance with the result of (Irene 2014). But after the flood in 2018, there was a hike in its status. In the correlation analysis (Table 3), available Fe showed positive correlation with available Mn (0.747\*\*), organic carbon (637\*) and available nitrogen (654\*\*). Available Mn showed a positive correlation with available Zn (0.695\*\*). Available Cu showed a positive correlation with available sulphur (0.563\*). Cation exchange capacity of the collected soil ranged from 5.92 to 30.64cmol (+) kg<sup>-1</sup> with a mean value of 18.87 cmol (+) kg<sup>-1</sup>. Soil samples collected from Veeyapuram reported a low CEC of

5.92 cmol (+) kg<sup>-1</sup> and the highest CEC of 30.64 cmol (+) kg<sup>-1</sup>. The correlation matrix (Table 3) showed a strong positive correlation between CEC and organic carbon (0.779\*\*\*) (Fig 1), available potassium (0.571\*), and available iron (0.717). Organic carbon contributes more to CEC. Higher values of CEC in acid sulphate soils of Kuttanad is mainly due to the presence of clay and organic matter (Thampatti, 1996).

### Conclusion

Chemical characterization of the acid sulphate soils of Upper Kuttanad reveals that the distribution of nutrients and its sufficiency range. Most of the soil falls under very strongly acidic group with high content of organic carbon. Almost 40 per cent of the soil samples recorded low in available P content. Among the secondary nutrients, almost all the samples from the different locations showed very high values for available sulphur. The hike in micronutrient status showed the impact of flooding in these locations. This characterization is the basic study that will help to sort out the effect of acidity on the availability of nutrients and their interaction, followed by its management.

### References

1. Baruah BK, Das B, Haque A, Medhi C, Misra AK. Sequential extraction of common metals (Na, K, Ca and Mg) from surface soil. *J. Chem. Pharm. Res.* 2011; 3(5); 565-573.
2. Bray RH, Kurtz LT. Determining total, organic and available forms of phosphate in soils. *Soil Sci.* 1945; 59; 39-45
3. Hendershot WH, Duquette M. A simplified barium chloride method for determining cation exchange capacity and exchangeable cations. *Soil Sci. Soc. Am. J.* 1986; 50; 605-608.
4. Irene EJ. Wet analysis for nutrient prescription in paddy soil. M. Sc. (Ag) thesis, Kerala Agricultural University, Thrissur. 2014: 192.
5. Jackson ML. *Soil Chemical Analysis*. Prentice-Hall Inc., Englewood Cliffs, NJ: 1958: 498 .
6. Kabeerathumma S, Patnaik S. Studies on the change of lime potential of acid sulphate soils of Kerala on submergence. *Agric. Res. J. Kerala.* 1980; 18; 229-235
7. Manorama Thampatti, KC. Morphological, physical and chemical characterization of the soils of North Kuttanad (Doctoral dissertation, Department of Soil Science and Agricultural Chemistry, College of Horticulture, Vellanikkara). 1997:160
8. Massoumi J, Cornfield AH. A rapid method for determination of sulphate in water extracts of soils. *Analyst.* 1963; 88; 321-322.

9. Mathews RP, Jose AI. Effect of submergence on inorganic phosphorus fractions and available phosphorus in two acid rice soils of Kerala. *Agric. Res. J. Kerala.* 1984; 22; 107-112
10. Money NS. Studies on the soils of kuttanad, part I. Toxic factors. *Agric. Res. J. Kerala* 1. 1961: 52-58.
11. Nhung, MT, Ponnampereuma FN. Effects of calcium carbonate, manganese dioxide, ferric hydroxide and prolonged flooding on chemical and electrochemical changes and growth of rice in a flooded acid sulphate soil. *Soil Sci.* 1966: 102; 29-41
12. Pareek BL, Mathur CM. Available and total phosphorus status of soils of Raiastan. *J. Indian Soc. Soil Sci.* 1969: 147-149.
13. Pisharody PN, Brito-Mutunayagam, APA. Manganese status of rice soils of Kerala. *Agric. Res. J. Kerala.* 1967: 5; 39-48
14. Raychaudhuri SP, Landey RJ. Effect of soil reaction on the availability of phosphorus and potash. *J. Ind. Soc. Soil Sci.* 1960: 8; 171-175.
15. Sims JR, Johnson GV. Micronutrient soil tests. In: Mortvedt, J.J., Cox, F.R., human, L.M. and Welch, R.M. *Micronutrient in Agriculture* (2nd ed.), SSSA, Madison, USA. 1991: 427-476.
16. Subbiah BV, Asija GLA. A rapid procedure for the estimation of available nitrogen in soils. *Curr. Sci.* 1956: 25; 259-260.
17. Tabatabai MA. Sulfur. In: Page, A. L., Miller, R. H., and Keeney, D. R. (eds.). *Methods of soil analysis. Part 2- Chemical and microbiological properties.* (2nd Ed.). American Society of Agronomy, Madison, Wisconsin, USA. 1982: 501-538
18. Walkley AJ, Black IA. Estimation of soil organic carbon by chromic acid titration method. *Soil Sci.* 1934: 31; 29-38.

	pH	EC (dSm <sup>-1</sup> )	OC (%)	Av N(kg/ha)	Av P (kg/ha)	Av K (kg/ha)	Av Ca (mg/kg)	Av Mg(mg/kg)	Av S(mg/kg)	Av Fe(mg/kg)	Av Mn(mg/kg)	Av Cu(mg/kg)	Av Zn(mg/kg)	CEC (cmol(+) kg <sup>-1</sup> )
1	4.76	0.09	0.42	170.72	15.67	34.11	197.10	39.22	719.37	721.66	7.12	3.54	5.14	14.14
2	5.08	0.05	1.89	263.20	2.14	86.59	446.13	61.63	725.66	889.25	8.22	7.18	5.22	13.58
3	4.85	0.06	1.38	219.32	97.04	90.58	519.03	84.46	3239.49	759.24	15.28	10.59	7.42	13.49
4	5.95	0.08	2.96	176.72	118.29	103.68	979.43	98.91	1277.78	561.66	27.25	12.91	8.26	15.75
5	5.56	0.13	3.91	402.01	53.76	769.25	729.61	163.55	57.16	1512.48	8.84	3.67	5.73	25.59
6	5.07	0.06	1.16	171.58	3.50	84.32	462.87	94.41	943.43	398.16	44.69	19.85	20.52	5.92
7	4.91	0.31	4.33	368.29	45.12	146.44	850.46	212.77	3361.50	2817.99	247.07	17.04	19.34	30.64
8	5.17	0.04	0.49	208.30	4.58	170.84	410.15	80.63	1219.66	104.74	7.27	9.94	1.38	13.72
9	4.70	0.18	2.47	221.01	2.41	204.01	1024.43	91.77	20.20	569.49	12.96	6.49	5.62	18.72
10	6.49	0.55	2.40	249.05	2.15	309.91	2581.35	192.63	763.68	1030.04	136.63	9.09	11.11	23.30
11	5.22	0.23	2.80	244.43	5.20	231.71	765.00	98.04	1418.05	1290.80	89.68	4.30	21.37	20.66
12	4.91	0.23	3.42	248.24	79.75	170.53	573.58	129.88	2839.15	1056.29	25.19	35.69	7.99	23.31
13	4.85	0.31	2.16	221.04	77.92	150.55	271.33	142.20	81.14	466.58	5.17	2.30	1.03	13.05
14	4.63	0.55	2.32	480.45	18.16	344.60	618.12	225.86	43.58	1608.93	22.34	16.37	5.14	26.68
15	4.74	0.97	3.07	317.30	55.21	338.72	292.88	372.51	2149.19	218.21	2.38	18.92	4.28	24.50
<b>Mean</b>	5.13	0.26	2.35	264.11	38.73	215.72	714.76	139.23	1257.27	933.70	44.01	11.86	8.64	18.87

**Table 2: Chemical characteristics of soil samples from Upper Kuttanad****Table 3: Correlation analysis**

	pH	EC	OC	Av N	Av P	Av K	Av Ca	Av Mg	Av S	Av Fe	Av Mn	Av Cu	Av Zn	CEC
pH	1													
EC	-0.046	1												
OC	0.163	0.363	1											
Av N	-0.157	0.482	0.562*	1										
Av P	0.041	-0.012	0.35	-0.052	1									
Av K	0.251	0.327	0.514*	0.671**	-0.018	1								
Av Ca	0.783***	0.21	0.261	0.036	-0.181	0.216	1							
Av Mg	-0.042	0.929***	0.541*	0.625*	0.13	0.473	0.129	1						
Av S	-0.14	0.066	0.25	-0.06	0.429	-0.318	-0.093	0.174	1					
Av Fe	0.005	0.068	0.637*	0.654**	-0.017	0.259	0.222	0.211	0.273	1				

Av Mn	0.277	0.151	0.462	0.219	-0.139	-0.058	0.506	0.224	0.424	0.747**	1			
Av Cu	-0.165	0.238	0.282	0.12	0.255	-0.136	-0.064	0.322	0.563*	0.101	0.122	1		
Av Zn	0.177	-0.088	0.269	-0.044	-0.211	-0.138	0.254	-0.036	0.346	0.467	0.695**	0.195	1	
CEC	0.054	0.579*	0.779***	0.783***	0.059	0.571*	0.328	0.664**	0.215	0.717**	0.495	0.231	0.09	1

UNDER PEER REVIEW




Fig 1: Relation between CEC AND OC

