

Standardization of recipe for value added Herbal Kinnow and Guava jelly for its better quality and shelf life

Abstract

During the year 2023, a field experiment was carried out at the Post Harvest Laboratory of Horticulture Department, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj to investigate "Standardization of recipe for value-added Herbal Kinnow and Guava jelly for better quality and shelf life." The experiment was carried out using a Completely Randomized Design (CRD), with ten treatments replicated thrice. Total number of treatments were ten viz. (T0 Control + 50% Sugar), (T1 50% Sugar + 2.0 g Ginger Extract), (T2 50% Sugar + 4.0 g Ginger Extract), (T3 50% Sugar + 6.0 g Ginger Extract), (T4 50% Sugar + 2.0 g Cardamom), (T5 50% Sugar + 4.0 g Cardamom), (T6 50% Sugar + 6.0 g Cardamom), (T7 50% Sugar + 2.0 g Mint Extract), (T8 50% Sugar + 4.0 g Mint Extract), (T9 50% Sugar + 6.0 g Mint Extract). The treatment (T6 50% Sugar + 6.0 g Cardamom) was found superior in respect of parameters like TSS(0Brix), pH, acidity(%), ascorbic acid (mg/100g), reducing sugar(%), non-reducing sugar(%), Colour score, texture, flavour, taste, overall acceptability. The benefit-cost ratio was found to be the highest (1:2.20) in the treatment (T6 50% Sugar + 6.0 g Cardamom). Maximum Mean shelf life of 41.66 days was found in (T6 50% Sugar + 6.0 g Cardamom).

Keywords: Herbal, Kinnow -Guava jelly, Sugar, Ginger, Cardamom, Mint

1. Introduction

In India, citrus ranks third in terms of output and acreage after mango and banana. Mandarin occupies a prominent position and makes up more than 50% of the area covered by citrus fruits [1]. The most desirable among the new exotic cultivars cultivated in India is beyond any doubt Kinnow, a cross between "King" and "Willow leaf mandarins." It has developed significantly and is currently commonly cultivated in Northern India's irrigated regions [2]. Numerous producers have started cultivating it on a commercial basis since it yields fruit with high dessert quality, is less prone to citrus decline, and has a low incidence of granulation [3]. Additionally, it possesses a broad range of paedoclimatic adaptation and early bearing. A fruit high in nutrients, kinnow has total

soluble solids of 14%, acidity of 0.79%, reducing sugar of 4.83%, pectin of 0.97%, and ascorbic acid of 26.50 mg/l 00 ml [4]. The fruits have a distinctively attractive fragrance and generally juicy with a pleasant flavor.

Guava (*Psidium guajava* Linn.) is well-known for its culinary and dietary benefits all over the world. Quercetin, Guaijaverin, Isoflavonoids, Gallic Acid, Catechin, Epicatechin, Rutin, Naringenin, Kaempferol Flavonoids, and Galactose-Specific Lecithins are only a few of the plant-derived compounds that have demonstrated potential efficacy. studies investigating the effects of various leaf, seed, pulp, skin, and fruit extracts in varying concentrations on the prevention of cancer, management of blood pressure, and treatment of diarrhea in mice and other animal

models as well as in controlled human research. Scientific research has verified several of the traditional applications. The plant's principal constituents have been thoroughly investigated for their pharmacological action, and the results demonstrate antioxidant, antipyretic, antifungal, antibacterial, hypotensive, analgesic & anti-inflammatory effects [5,6].

Ginger is a spice that may be used both in cooking and as a delicacy or medication. It is the subterranean stem of the *Zingiber officinale* plant, which produces ginger. Zingerone, shogaols, and gingerols are a combination of volatile oils that make up one to three percent of the weight of the freshly harvested ginger root and give it its distinctive flavor and aroma. The digestive tract's motility is enhanced in experimental mice by gingerols, which also have analgesic, sedative, antipyretic, and antibacterial activities. The nonvolatile phenylpropanoid-derived substances, notably gingerols, and shogaols, which are formed from gingerols when ginger is dried or cooked, are what give ginger its pungent flavor. During this process, gingerols are also converted into zingerone, a less fragrant substance with a spicy sweet aroma taste [7,8].

Mint has delicate stems and soft leaves. It is better to add them either uncooked or just before serving. They retain their delicate flavour and texture thanks to this. Mint is a sustainable approach to flavour food since it is reasonably simple to grow and can be grown at home. A little under a third of a cup, about 14 grams, of spearmint Calories: 6 grams Fibre: one gram. 12% of the RDI for vitamin A 9% of the RDI is iron. 8% of the RDI for manganese 4% of the RDI for folate. Irritable bowel syndrome (IBS) is a prevalent condition that affects the digestive

system. Digestion-related symptoms such as gas, bloating, stomach discomfort, and changes in bowel habits are its hallmarks [9].

In South Asia, green cardamom is frequently used to treat digestive disorders and to prevent infections of the teeth and gums, as well as to prevent and cure throat problems, lung congestion, pulmonary tuberculosis, inflammation of the eyelids, and infections of the teeth and gums. It was reputedly used as an antidote for both snake and scorpion venom, as well as to dissolve kidney and gall stones. In systems such as traditional Chinese medicine in China, Ayurveda in India, Japan, Korea, and Vietnam, amomum is utilized as a spice and an ingredient. Traditional Indian medicine utilizes species from the genus *Amomum* [10].

2. Materials and Methods

The Experimental was conducted in Completely Randomized Design (CRD) with 10 treatments of different level of Guava, Kinnow and sugar with three replications in the Post Harvest Laboratory of Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during 2022. Total number of treatments were nine viz. T0 (Control + 50% Sugar), T1 (50% Sugar + 2.0 g Ginger Extract), T2 (50% Sugar + 4.0 g Ginger Extract), T3 (50% Sugar + 6.0 g Ginger Extract), T4 (50% Sugar + 2.0 g Cardamom), T5 (50% Sugar + 4.0 g Cardamom), T6 (50% Sugar + 6.0 g Cardamom), T7 (50% Sugar + 2.0 g Mint Extract), T8 (50% Sugar + 4.0 g Mint Extract), T9 (50% Sugar + 6.0 g Mint Extract). The parameters were recorded like Physiochemical properties of various products i.e. T.S.S, acidity, reducing sugar, non-reducing sugar, pH, Vitamin C and sensory evaluation as

well as economics of different treatments were recorded.

3. Results and Discussion

The present investigation entitled “Standardization of recipe for value added Herbal Kinnow and Guava jelly for its better quality and shelf life” was carried out during 2022 in Post Harvest Laboratory of Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) India. The results of the present investigation, regarding the Standardization of value-added Herbal Kinnow and Guava jelly, have been discussed and interpreted in the light of previous research work done in India and abroad. The experiment was conducted in Completely Randomized design with 10 treatments, and three replications.

3.1 Quality parameters

It is evident that the table 1 was influenced by different treatments at successive stage 45 days of storage. There was significant differences between the treatments. The quality parameters on TSS (°brix), was recorded in T6(500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Cardamom) with (69.91) followed by T7 (500 ml Kinnow Juice + 500 ml Guava Juice + 2.0 g Mint Extract) with T7 (500 ml Kinnow Juice + 500 ml Guava Juice + 2.0 g Mint Extract) with (69.89) and minimum was T0 (Control). However T0 (Control) was recorded (0.63) highest acidity (%), followed by T₂(500 ml Kinnow Juice + 500 ml Guava Juice + 4.0 g Ginger Extract) and minimum T9 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Mint Extract) with (0.32). in case of Reducing Sugar (%), treatments T6(500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Cardamom) with (37.75),

followed by T7 (500 ml Kinnow Juice + 500 ml Guava Juice + 2.0 g Mint Extract) and minimum was T0 (Control) with (30.97). similarly in Non Reducing Sugar (%), treatment used T8(500 ml Kinnow Juice + 500 ml Guava Juice + 4.0 g Mint Extract) with (35.60), followed by T7 (500 ml Kinnow Juice + 500 ml Guava Juice + 2.0 g Mint Extract) and minimum T6 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Cardamom) with (23.90). Among the treatment used T8(500 ml Kinnow Juice + 500 ml Guava Juice + 4.0 g Mint Extract) with (3.76), followed by T7 (500 ml Kinnow Juice + 500 ml Guava Juice + 2.0 g Mint Extract) and minimum was T6 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Cardamom) with (3.29) in pH and Ascorbic Acid treatment T6(500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Cardamom) with (42.33), followed by T7 (500 ml Kinnow Juice + 500 ml Guava Juice + 2.0 g Mint Extract) and minimum ascorbic acid T0-Control with (39.45). A slight increase in total soluble solids during storage might be due to conversion of polysaccharides (present in fruits) into sugars during hydrolysis process. This finding agreed with the finding of Vikram and Prasad [11]. compositional changes in value added Kinnow-Aonla RTS revealed that there was increase in the level of TSS during the storage period (six months) and in aonla RTS beverage [12]. Degradation of pectin substances into soluble solids might have contributed towards increase the level of acidity in the during storage period of aonla jam. [13] The increase in reducing sugar was slightly higher in storage condition that could be attributed to more rapid hydrolysis of polysaccharides and their subsequent conversion into sugars. Nath and Yadav and Deka et al. reported similar finding with lime aonla blended RTS [14,15]. The present findings are in

agreement with those of Khan et al. [16] who observed decrease in non-reducing sugar in strawberry jam during storage of 60 days at ambient temperature. [17] also reported a similar trend in jam, squash and RTS beverage of local mango stored for 100 days. Variations in pH during storage may be due to change in chemical properties which are affected by storage conditions. This finding agreed with the finding of Vikram and Prasad and Rayguru et al. [18,19] also reported similar trend in apple jam. The decrease in Ascorbic Acid was slightly higher in storage condition that could be attributed to more rapid hydrolysis of polysaccharides and their subsequent conversion into sugars. Nath and Yadav and Deka et al. reported similar finding with lime aonla blended RTS [16,17].

3.2 Sensory Evaluation

It is evident that the table 2 was influenced by different treatments at successive stage 45 days of storage. There was significant differences between the treatments. The Sensory Evaluation on Colour / Appearance score, treatment T6 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Cardamom) with (8.08) have highest Colour / Appearance, followed by followed by T3 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Ginger Extract) and minimum was T0 (Control) with (6.17), however in Flavour, treatment T6 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Cardamom) with (7.60) found highest Flavour, followed by T3 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Ginger Extract) and minimum was T0 (Control) with (5.50), similarly in Taste,

treatment T6 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Cardamom) with (7.60) were significantly superior, followed by T3 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Ginger Extract) with (7.30) and minimum was T0 (Control) with (5.70). in case of Aroma , treatment T6 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Cardamom) with (7.80), followed by T3 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Ginger Extract) and minimum was T0 (Control) with (5.50) and in overall acceptability the highest score in T6 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Cardamom) with (7.74), followed by T3 (500 ml Kinnow Juice + 500 ml Guava Juice + 6.0 g Ginger Extract) and minimum was T0 (Control) with (5.63). Deterioration of colour due to enzymatic and nonenzymatic reactions on pigment during storage of fruit products impair the quality of the products. It could be attributed to non-enzymatic reactions, which occur between nitrogenous compounds and sugars or organic acid and organic acids with sugars. Similar results were reported in sweet orange based products [16]. The decreasing trend was observed for flavour, taste and texture with increase storage period. This might be due to degradation of volatile substance and flavor constituents. Similar result was reported in aonla segments in syrup, prepared from stored fruits and in aonla RTS beverages [17,12]. Overall acceptability scores were decreased in all the treatments during storage due to decline in colour, consistency and flavour scores. Similar results were reported in aonla beverages [18,19,11].

Table 1: Effect of different treatments on quality parameters of Kinnow-Guava Jelly

Treatments	TSS (°brix)	Acidity (%)	Reducing Sugar (%)	Non Reducing Sugar (%)	pH	Ascorbic Acid
T0	68.96	0.63	30.97	25.97	3.29	39.45
T1	69.14	0.41	32.14	27.57	3.46	40.23
T2	69.09	0.58	32.84	26.87	3.45	40.41
T3	69.25	0.32	32.94	27.14	3.55	41.55
T4	69.33	0.36	34.37	24.27	3.6	41.31
T5	69.74	0.46	33.51	30.4	3.55	41.01
T6	69.91	0.32	37.75	23.9	3.42	42.33
T7	69.89	0.35	35.71	34.64	3.62	41.49
T8	69.49	0.43	33.07	35.6	3.76	40.53
T9	69.66	0.32	33.24	30.57	3.63	41.01
Result	S	S	S	S	S	S
	0.3	0.117	0.353	0.394	0.104	0.208
C.D. at 5%	0.636	0.249	0.749	0.836	0.221	0.442
CV	16.434	6.428	19.349	21.607	5.715	11.409




Figure 1: Effect of different treatments on quality parameters of Kinnow-Guava Jelly

UNDER PEER REVIEW

Table 2: Effect of different treatments on Sensory Evaluation of Kinnow-Guava Jelly

Treatments	Colour	Flavour	Taste	Aroma	Overall Acceptability
T0	6.43	6	6.2	6	6.09
T1	6.77	6.6	6.8	6.6	6.67
T2	7.43	7.2	7.4	7.4	7.37
T3	8.17	7.7	8	8	7.93
T4	6.6	6.2	6.4	6.2	6.32
T5	7.17	7	7.3	7.2	7.17
T6	8.5	8.2	8.2	8.4	8.28
T7	7	6.8	7.1	7.1	7
T8	7.6	7.5	7.6	7.6	7.56
T9	7.8	7.6	7.8	7.8	7.74
Result	S	S	S	S	S
S. Ed. (\pm)	0.317	0.266	0.244	0.251	0.256
C.D. at 5%	0.672	0.564	0.518	0.533	0.543
CV	17.367	14.575	13.383	13.77	17.367




Figure 2: Effect of different treatments on Sensory Evaluation of Kinnow-Guava Jelly

Treatments	Common expenditure (Rs)(1924.526÷8)	Selling price in market (Rs/bottle)	Gross return for 3 bottle (Rs)	Net return for 3 Bottle(Rs)	B : Cratio
T0-Control -500 ml Kinnow Juice + 500 ml Guava Juice + 50% Sugar	265.57	140	420	154.43	1:1.58
T1-500 ml Kinnow Juice + 500 ml Guava Juice + 50% Sugar + 2.0 g Ginger Extract	265.57	160	480	214.43	1:1.81
T2-500 ml Kinnow Juice + 500 ml Guava Juice + 50% Sugar + 4.0 g Ginger Extract	265.57	175	525	259.43	1:1.98
T3-500 ml Kinnow Juice + 500 ml Guava Juice + 50% Sugar + 6.0 g Ginger Extract	265.57	195	585	319.43	1:2.20
T4-500 ml Kinnow Juice + 500 ml Guava Juice + 50% Sugar + 2.0 g Cardamom	265.57	160	480	214.43	1:1.81
T5-500 ml Kinnow Juice + 500 ml Guava Juice + 50% Sugar + 4.0 g Cardamom	265.57	175	525	259.43	1:1.98
T6-500 ml Kinnow Juice + 500 ml Guava Juice + 50% Sugar + 6.0 g Cardamom	265.57	195	585	319.43	1:2.20
T7-500 ml Kinnow Juice + 500 ml Guava Juice + 50% Sugar + 2.0 g Mint Extract	265.57	160	480	214.43	1:1.81
T8-500 ml Kinnow Juice + 500 ml Guava Juice + 50% Sugar + 4.0 g Mint Extract	265.57	175	525	259.43	1:1.98
T9-500 ml Kinnow Juice + 500 ml Guava Juice+ 50% Sugar + 6.0 g Mint Extract	265.57	195	585	319.43	1:2.20

Table No. 3 Cost of Production

4. CONCLUSION

Based on the data, it is stated that kinnow is ideal for making jelly when combined with guava. Because of its physicochemical properties, Organoleptic qualities, maximum Mean shelf life is of 41.66 days, maximum net return of 106.47 Rs/ Bottle, and best Benefit Cost Ratio of 1:2.20, the blend of Kinnow 50% + Guava 50% treated with 6.0gm cardamom extract i.e. T6 was found to be the best recipe for Herbal Kinnow-Guava jelly.

References

- Food and Agriculture Organization (FAO). Statistical Yearbook. World Food and Agriculture. Food and Agriculture Organization of the United Nations, Rome, Italy. 2021; 86–368.
- Bangladesh Bureau of Statistics (BBS). Yearbook of Agricultural Statistics of Bangladesh. Bangladesh Bureau of Statistics, Statistics and Information Division, Ministry of Planning, Government of the Peoples Republic of Bangladesh, Dhaka. 2022; 210–33.
- Bhat SA, Singh ER. Extraction and characterization of pectin from guava fruit peel. *Int J Adv Res Technol.* 2014;2:1–7.
- Kuchi VS, Gupta R, Tamang S. Standardization of recipe for preparation of guava jelly bar. *J Crop Weed.* 2014;10:77–81.
- Jain PK, Asati VK. Evaluation of guava cultivars for pulp preparation. *J Food Sci Technol.* 2004;41:6846.
- Paull RE, Duarte O. *Tropical Fruits*, 2nd Ed. CAB International. 2011:327–65.
- Coppens d'Eeckenbrugge G, Sanewski GM, Smith MK, Duval MF, Leal F. Ananas In: Kole C. (ed). *Wild crop relatives: genomic and breeding resources: tropical and subtropical fruits.* Springer. 2011:21–41.
- Joy PP. *Benefits and Uses of Pineapple*, Pineapple Research Station (Kerala Agricultural University), Vazhakulam. 2010 ([http:// prsvkm. kau. in/ book/importance](http://prsvkm.kau.in/book/importance)).
- Khan MA. Bitter truth: slow poisoning continues unabated. *The Daily Star*, 17 May, Bangladesh. 2004.
- Anand SP, Sati N. Artificial preservatives and their harmful effects: looking toward nature for safer alternatives. *Int J Pharm Sci Res.* 2013;4:2496–501.
- Molla MM. Effect of foxtail millet diet on liver injury and blood lipid profile induced by D-galactoseamine in mice. PhD dissertation,

College of Food Science and Nutritional Engineering, China Agricultural University, Beijing-100083, China. 2016;113p.

Amir RM, Randhawa MA, Sajid MW, Nadeem M, Ahmad A, Watto FM. Evaluation of various soaking agents a novel tool for heavy metal residues mitigation from spinach. *Food Sci Technol.* 2020;39:176–80.

Wang Z, Jackson LS, Jablonski JE. Factors affecting the level of heavy metals in juices processed with filter aids. *J Food Prot.* 2017;80:892–902.

Abu-Almaaly RA. Effect of cooking method on the content of heavy metals in rice that available in local market. *Plant Arch.* 2020;20:2976–81.

Inobeme A, Ajai AI, Eziukwu C, Obigwa PA, Okonkwo S, Ekwoba LM. Effect of cooking methods on heavy metals content of food. *J Xidian Univ.* 2020;14:704–14.

Shinta YC, Zaman B, Sumiyati S. Citric acid and EDTA as chelating agents in phytoremediation of heavy metal in polluted soil: a review. *IOP Conf Series: Earth Environ Sci.* 2021;896:1–8.

Belitz H-D, Grosch W, Schieberle P. *Food Chemistry*. In: Burghagen M (3rd ed.) Translation from the Vth German edition 3rd ed. Springer, Germany. 2004.

Bang LM, Bunting C, Molan PC. The effect of dilution on the rate of hydrogen peroxide production in honey and its implications for wound healing. *J Altern Complement Med.* 2003;9:267–73.

Badawy OFH, Shafil SSA, Tharwat EE, Kamal AM. Antibacterial activity of bee honey and its therapeutic usefulness against *Escherichia coli* O157:H7 and *Salmonella typhimurium* infection. *Rev Sci Technol Int Epiz.* 2004;23:1011–22. Mundo MA, Padilla-Zakour OI, Worobo RW. Growth inhibition of foodborne pathogens and food spoilage organisms by select raw honeys. *Int J Food Microbiol.* 2004;97:1–8.

Lusby PE, Coombes AL, Wilkinson JM. Bactericidal activity of different honeys against pathogenic bacteria. *Arch Med Res.* 2005;36:464–7.

Adeleke OE, Olaitan JO, Okepeke EI. Comparative antibacterial activity of honey and gentamicin against *Escherichia coli* and *Pseu-domonas aeruginosa*. *Ann Burn Fire Disasters.* 2006;19:2014.

Basualdo C, Sgroy V, Finola MS, Juam M. Comparison of the antibacterial activity of honey from different provenance against bacteria usually isolated from skin wounds. *Vet Microbiol.* 2007;124:375–81.

Al-Mamar y M, Al-Meerri A, Al-Habori M. Antioxidant activities and total phenolics of different types of honey. *Nutr Res.* 2002;22:1041–7.

Alvarez-Suarez JM, Tulipani S, Romandini S, Bertoli E, Battino M. Contribution of honey in nutrition and human health: a review. *Mediterr J Nutr Metab.* 2010;3:15–23.

Johnston JE, Sepe HA, Miano CL, Brannan RG, Alderton AL. Honey inhibits lipid oxidation in ready-to-eat ground beef patties. *Meat Sci.* 2005;70:627–31.

Turkmen N, Sari F, Poyrazoglu ES, Velioglu YS. Effects of prolonged heating on antioxidant activity and colour of honey. *Food Chem.* 2006;95:653–7.

Rakha MK, Nabil ZI, Hussein AA. Cardioactive and vasoactive effects of natural wild honey against cardiac malperformance induced by hyperadrenergic activity. *J Med Food.* 2008;11:91–8.

Eteraf-OskoueïT NM. Traditional and modern uses of natural honey in human diseases: A Review. *Iran J Basic Med Sci.* 2013;16:731–42.

Daily Bangladesh. Good news to honey farmers. *Daily Bangladesh news* 17 February. 2020.

UNDER PEER REVIEW