

## Original Research Article

### **Comparative study on Pasting and Functional properties of Composite flour produced from Acha (*Digitaria exilis*) starch, Chickpea (*Cicer arietinum*) and wheat (*Triticum aestivum*) flour blends.**

#### **Abstract**

Introduction: Composite flour can relatively be any wheat flour composition with a particular substitution percentage of any type of flour produced from any other raw material. This is done to increase the nutritional profile, functional properties or further utilization of the said raw materials.

**Objective:** The functional properties of flour blends play an important part in the processing of food products. Functional properties determine the usability of the blends in bakery products where hydration aid handling in products such as ground meat, baked foods and pancakes where oil and absorption properties is of utmost importance. Functional properties plays significant role in the physicochemical properties in determining the complex interaction between the composition, structure, and molecular conformation, therefore the objective of the study was to produce composite flour from blends of acha starch, chickpea and wheat flour and determine the pasting and functional properties of the blends for food uses.

**Methodology:** Composite flour was produced in different blends using D-Optimal experimental runs. Pasting and functional characteristics is essential in estimating the properties of food paste and products during and after it is cooked.

**Results:** The pasting properties of the blends of the flour, the peak 1(RVU), trough 1(RVU), breakdown(RVU), final viscosity(RVU), set back(RVU), peak time(min) and pasting temperature( $^{\circ}$ C) ranged between 1722.50 and 2483.5, 931.0 and 1723.00, 659.00 and 824.50, 1949.5.00 and 3267.0, 1095.50 and 1544.00, 5.44 and 6.00, 76.28 and 88.83 respectively unit. Results for functional analysis of the flours blends, water absorption capacity(g/ml), oil absorption capacity(g/ml), bulk density(g/ml), swelling power(%), solubility index ranged between 175 and 258, 186 and 232, 0.5588 and 0.68, 4.52 and 6.08, 3.50% and 7.50% respectively.

**Conclusion:** The flour blends of Acha starch, chickpea flour and wheat flour showed that it can be good raw materials for food that requires high water and oil absorption properties, bulk density, swelling power, set viscosity and final viscosity. The best sample blend was sample (wheat flour 52.50%, acha flour 42.50% and 5% chickpea flour).

**Keywords; Composite flour, pasting properties, Functional properties**

## 1. Introduction

Composite flour can relatively be any wheat flour composition with a particular substitution percentage of any type of flour produced from any other raw material. This is done to increase the nutritional profile, functional properties or further utilization of the said raw materials. In another words, it is a flour made by blending or mixing varying proportion of more than one non-wheat flour with or without wheat flour and used for production food products that are conventionally made from wheat flour and increase the essential nutrients in human diet. Functional properties are the fundamental Physico-chemical properties that reflect the complex interaction between the composition, structure, molecular conformation and physico-chemical properties of food components together with the nature of environment in which these are associated and measured (1). Pasting characteristic is important in predicting the behaviour of food paste during and after cooking. Attainment of the pasting temperature is essential in ensuring swelling, gelatinization and subsequent gel formulation during processing (2). Acha(*Digitariaexilis*) is an ancient cereal grain that is cultivated in most West Africa countries which is also known as Fonio, it is the tiniest grain in the millet family and a main source of protein (3). Nutritionally, Acha contains 67.1 to 91% carbohydrate, 7.0 to 11.5% protein, 11.30% crude fibre, 1.3 to 5.2% fat and 4619KcalGE/KG energy value. (4). Chickpea (*Cicerarietinum*) is an annual legume crop which has a generous supply of carbohydrates, proteins, fibers lipids, and mineral contents that belongs to Fabaceae family Chickpea contains essentials vitamins such as niacin, riboflavin, thiamine, folate and vitamin A. It has several health benefits and in combination with other pulses and

cereals helps in conditions such as Diabetes type 2, digestive diseases, cardiovascular diseases, and some cancers (5).

## **2. Materials and methods**

### **2.1 Collection of materials**

Acha grain was procured from Lupe market, FCT Abuja ,Nigeria. While the chickpea grains and wheat flour was purchased from Bazaar supermarket at Ilupeju in Lagos state, Nigeria.

### **2.2 Production of the flour blends**

#### **2.2.1 Acha starch**

The acha starch was obtained using the method of (6). The acha grains was cleaned, steeped in water for 24 hours and wet milled. After which it was sieved and the starch was allowed to settle and the water was decanted so as to obtain the starch granules. The starch granule was dried using hot air oven and milled into fine powder. The starch obtained was stored in a cool dry place in an airtight container prior to usage.

#### **2.2.2 Chickpea and wheat flour**

The chickpea grains were screened for broken seeds, dust and stones. The seeds was washed with portable water and then soaked for 17hours, decanted and replaced with clean water in the ratio 1:10 (w/v) to the chickpea. The grains was boiled for one hour at 100°C in an insulated kettle, decanted, dried for 8hours at 38-40°C. The grain was grinded into powder and sieved with a 80-mesh stainless steel sieve (250um grain size distribution) to obtain fine flour. The obtained chickpea flour was packed in an airtight bag prior to usage (7). Wheat flour, Acha starch and Chickpea was mixed in different proportions with low and high limits set at (45: 60%), (30: 50%), and (5:20%) respectively. D-optimal design was employed because it puts into consideration the design constraints of the flour blend mixture as shown in Table 1. This resulted into fourteen (14) experimental runs as shown in Table 2.

## **3. Determination of pasting properties**

Brabender visco-amylograph (BVA), rheometer and rapid visco analyzer (RVA) were used to examine the pasting characteristics which includes breakdown, setback, peak, trough, final viscosity, peak temperature and peak time.

#### **4. Determination of Functional properties**

Swelling power solubility profile of starch was determined according to the method of (8). Water absorption capacity of the starch was determined according to the method of (9). Bulk density was determined using method of (10). Oil absorption capacity was determined using method of (11).

#### **5. Statistical Analysis**

All the data generated from the experiment analyzed using analysis of variance (ANOVA) using SPSS version 22.0 and the difference between significant mean values were separated at  $p < 0.05$  probability level with Duncan's Multiple Range Test.

#### **6. Results and Discussion**

##### **6.1 Pasting properties**

Pasting properties of the composite flour from the Wheat- Chickpea flour and Acha starch is found on Table 3. Peak viscosity varied from 1722.50 to 2483.50RVU. Peak viscosity describes swelling of starch or water-binding capability during the process of heating. It is also an indication of the ability of the products to swell freely before their physical breakdown (12). The trough viscosity ranged from 931 to 1723RVU. High trough viscosity observed in this study revealed that these combinations can survive high-heat treatments during processing. High trough values may represent low cooking losses and superior eating quality (13). The breakdown ranged between 659 and 824.50RVU. Breakdown viscosity is a measure of degree of starch disintegration or the hot paste stability of the starch (14). The breakdown viscosity in the study was low revealing that the paste stability of the flour blends is low. The final viscosity ranged between 1949.50 and 3267.00 RVU. Final viscosities are essential in predicting the strength of the flour sample to form gel during heat processing. The final viscosity is the most commonly used parameter to determine a particular

sample quality and indicates the stability to form various paste or gel after cooling (15). The final viscosities studied shows high strength of the flours to gelatinize during heating process. Setback viscosity indicates gel stability and potential for retrogradation (16). The setback ranged between 1018.50 and 1544.00RVU. The composite flour could form a great paste that could be useful in the confectionaries applications. Peak time of the composite flour blends varied between 5.44 and 6.00mins. The pasting temperature ranged between 84.80 and 88.83 °C. Pasting temperature is the lowest temperature needed for gel to be formed, as well as the temperature at which the viscosity during the heating process goes rises. The granules begin to swell when the temperature is higher than that at which gelatinization occurs (17). The samples shows a high possibility of resistance against swelling in the ingredient, which could be correlated to the amount amylose and amylopectin in the samples.

## **6.2 Functional Properties wheat-chickpea flour and acha starch**

The mean value for functional properties of wheat-chickpea flour and acha starch for different experimental runs is as shown in Table 4. Significant ( $p < 0.05$ ) difference was seen in the functional properties of the composite flour. The water absorption capacity of wheat-chickpea flour and acha starch ranged from 175.00 to 258.00%. Water Absorption Capacity (WAC) is the ability of a product to associate with water under a condition where water is limiting (18). High water absorption capacity could be attributed to the alteration in starch polymer structure while low value indicates compactness of the structure (19). The values obtained in this study were higher than the values (87-111%) reported (20) for blends of wheat, acha, cowpea and *Moringaoleifera* powder but agrees with the values(164-252%) reported (21) which means that the blends has a good water holding capacity which a desirable feature in dough formation which correlated with an increase in the amylose leaching and solubility and loss of crystalline starch structure. The bulk density ranged from 0.5588 to 0.7031g/ml. Bulk density is a determinant of flour expansion and an indicator of the porosity of food products (22). Low bulk density is desired in flour blends as it's contributes to lower dietary bulk, ease of packaging and transportation (23). The low bulk density of the samples

show the flour can be an advantage in the formulation of weaning foods with lower dietary bulk and can reduce the cost of transportation. The swelling power of wheat-chickpea flour and acha starch ranged between 4.52 and 6.08%. Swelling capacity actually describes the firmness of the bonds in the crystalline part of the starch granule which ultimately shows the easiness of cooking the flour or starch. Usually, the starch granules with more crystalline areas and stronger bonds swell less in cold or hot water as it forms a low viscosity paste with higher retrogradation (24). The oil absorption capacity of composite flour from wheat-chickpea flour and acha starch ranged between 186.00 and 232.00%. The ability of flours to absorb and retain oil may enhance flavour retention and improve mouth feel (25). The oil absorption capacity of the blends were relatively high which indicates ease emulsification and fat absorption of the of flours are affected by the extent to which water is taken up and held down within starch the granules and the increase in solubility figures could be traced to increase leaching of solubilized amylose molecules from expanded starch granules promoted by disruption of the starches. Factors capable of influencing the solubility of flours include flour composition and particle size, density and pH, processing conditions and storage conditions (26). The solubility of wheat-chickpea flour and acha starch ranged from 3.50 to 7.50%.

## **7. Conclusion**

The flour blends of Acha starch, chickpea flour and wheat flour showed that it can be good raw materials for food that requires high water and oil absorption properties, bulk density, swelling power, set viscosity and final viscosity. The best blend was (wheat flour 52.50%, acha flour 42.50% and 5% chickpea flour). This research work has shown significant improvement in the functional and pasting properties of the flour resulting from the substitution of wheat with Acha starch and Chickpea flour in different formulations. The trend observed showed improvement in the functional properties of the product, especially in the water absorption and swelling power due to the addition of acha starch.

## References

1. Kinsella, J. E., & Melachouris, N. (1976). Functional properties of proteins in foods: a survey. *Critical Reviews in Food Science & Nutrition*, 7(3), 219-280.
2. Akinsola, A. O., Idowu, M. A., Babajide, J. M., Oguntona, C. R. B., & Shittu, T. A. (2018). Production and functional property of maize-millet based complementary food blended with soybean. *African Journal of Food Science*, 12(12), 360-366.
3. Adoukonou-Sagbadja, H., Dansi, A., Vodouhè, R., & Akpagana, K. (2006). Indigenous knowledge and traditional conservation of fonio millet (*Digitaria exilis*, *Digitaria iburua*) in Togo. *Biodiversity & Conservation*, 15, 2379-2395.
4. Idris, R. M., Yawas, D. S., Jolaiya, S. O., & Abdulrahman, M. A. PERFORMANCE EVALUATION OF AN IMPROVED ACHA (*DIGITERIA EXILIS*) DEHUSKING MACHINE.
5. Fitness, R. W. Healthy living made easy with tasty recipes, efficient workouts and evidence-based guidance on nutrition, weight loss and more.
6. Musa, H., Muazu, J., Bhatia, P. G., & Mshelbwala, K. (2008). Investigation into the use of fonio (*Digitaria exilis*) starch as a tablet disintegrant. *Nigerian Journal of Pharmaceutical Sciences*, 7(1), 67-78.
7. Rachwa-Rosiak, D., Nebesny, E., & Budryn, G. (2015). Chickpeas—composition, nutritional value, health benefits, application to bread and snacks: a review. *Critical reviews in food science and nutrition*, 55(8), 1137-1145.
8. Adebowale, A. A., Sanni, L. O., & Awonorin, S. O. (2005). Effect of texture modifiers on the physicochemical and sensory properties of dried fufu. *Food Science and Technology International*, 11(5), 373-382.
9. Azeez, A. T., Adegunwa, M. O., Sobukola, O. P., Onabanjo, O. O., & Adebowale, A. A. (2015). Evaluation of some quality attributes of noodles from unripe plantain and defatted sesame flour blends. *Journal of Culinary Science & Technology*, 13(4), 303-329.
10. Bezerra, C. V., Amante, E. R., de Oliveira, D. C., Rodrigues, A. M., & da Silva, L. H. M. (2013). Green banana (*Musa cavendishii*) flour obtained in spouted bed—Effect of drying on physico-chemical, functional and morphological characteristics of the starch. *Industrial crops and products*, 41, 241-249.

11. Adepeju, A. B., Gbadamosi, S. O., Omobuwajo, T. O., & Abiodun, O. A. (2014). Functional and physico-chemical properties of complementary diets produced from breadfruit (*Artocarpus altilis*). *African Journal of Food Science and Technology*, 5(4), 105-113.
12. Akinsola, A. O., Idowu, M. A., Babajide, J. M., Oguntona, C. R. B., & Shittu, T. A. (2018). Production and functional property of maize-millet based complementary food blended with soybean. *African Journal of Food Science*, 12(12), 360-366.
13. Asaam, E. S., Adubofuor, J., Amoah, I., & Apeku, O. J. D. (2018). Functional and pasting properties of yellow maize–soya bean–pumpkin composite flours and acceptability study on their breakfast cereals. *Cogent Food & Agriculture*, 4(1), 1501932.
14. Bakare, A. H., Osundahunsi, O. F., & Olusanya, J. O. (2016). Rheological, baking, and sensory properties of composite bread dough with breadfruit (*Artocarpus communis* Forst) and wheat flours. *Food Science & Nutrition*, 4(4), 573-587.
15. Ikegwu, O. J., Nwobasi, V. N., Odoh, M. O., & Oledinma, N. U. (2009). Evaluation of the pasting and some functional properties of starch isolated from some improved cassava varieties in Nigeria. *African Journal of Biotechnology*, 8(10).
16. Niba, L. L., Bokanga, M. M., Jackson, F. L., Schlimme, D. S., & Li, B. W. (2002). Physicochemical properties and starch granular characteristics of flour from various *Manihot esculenta* (cassava) genotypes. *Journal of food science*, 67(5), 1701-1705.
17. Otegbayo, B. O., Samuel, F. O., & Alalade, T. (2013). Functional properties of soy-enriched tapioca. *African Journal of Biotechnology*, 12(22).
18. Abiodun, O. A., Odedeji, J. O., Adepeju, A. B., & Oladapo, A. S. (2018). Physicochemical properties of modified trifoliolate yam starches. *Ife Journal of Science*, 20(1), 189-198.
19. Ubbor, S. C., Arukwe, D. C., Ezeocha, V. C., Nwoso, O. N., Iguh, B. N., & Nwibo, O. G. (2022). PRODUCTION AND QUALITY EVALUATION OF READY TO EAT EXTRUDED SNACKS FROM FLOUR BLENDS OF ACHA-COWPEA AND SWEET POTATO STARCH. *FUDMA JOURNAL OF SCIENCES*, 6(4), 245-153.
20. Orisa, C. A., & Udofia, S. U. (2020). Functional and pasting properties of composite flours from *Triticum durum*, *Digitaria exilis*, *Vigna unguiculata* and *Moringa oleifera* Powder. *Asian Food Science Journal*, 19(2), 40-49.
21. Moutaleb, O. H., Amadou, I., Amza, T., & Zhang, M. (2017). Physico-functional and sensory properties of cowpea flour based recipes (akara) and enriched with sweet potato. *J Nutr Health Food Eng*, 7(4), 00243.
22. Kraithong, S., Lee, S., & Rawdkuen, S. (2018). Physicochemical and functional properties of Thai organic rice flour. *Journal of Cereal Science*, 79, 259-266.

23. Aluge, O. O., Akinola, S. A., & Osundahunsi, O. F. (2016). Effect of malted sorghum on quality characteristics of wheat-sorghum-soybean flour for potential use in confectionaries. *Food and Nutrition Sciences*, 7(13), 1241-1252.
24. Olurin, T. O., Abbo, E. S., & Oladiboye, O. F. (2021). Production and evaluation of breakfast meal using blends of sorghum, bambara nut and date palm fruit flour. *Agro-Science*, 20(3), 30-36.
25. Kaur, M., Kaushal, P., & Sandhu, K. S. (2013). Studies on physicochemical and pasting properties of Taro (*Colocasia esculenta* L.) flour in comparison with a cereal, tuber and legume flour. *Journal of Food Science and Technology*, 50, 94-100.
26. Mirhosseini, H., & Amid, B. T. (2013). Effect of different drying techniques on flowability characteristics and chemical properties of natural carbohydrate-protein Gum from durian fruit seed. *Chemistry Central Journal*, 7(1), 1-14.

**Table 1. Blends formulation based on optimal of mixture design, design Constraints**

Constraints	Low limits	High limits
A	45	60
B	30	50
C	5	20

A=Wheat flour, B=Acha starch, C=Chickpea flour

**Table 2. Formulation of wheat, Acha starch and chickpea flour blends using D-optimal mixture design**

Runs	Sample	WF	AS	CF
1	CAD	60.00	30.00	10.00
2	FDA	45.00	42.50	12.50
3	DEB	52.50	42.50	5.00
4	ORD	55.00	30.00	15.00
5	DOP	45.00	35.00	15.00
6	XAD	52.00	36.00	12.00
7	FEM	48.50	35.50	16.00
8	SNM	50.00	30.00	20.00

10	KLM	60.00	35.00	5.00
11	DEB	52.50	42.50	5.00
12	GTK	45.00	50.00	5.00
13	KLM	60.00	35.00	5.00
14	GTK	45.00	50.00	5.00

WF: Wheat Flour, AS: Acha starch, CF: Chickpea flour

**Table 3. Pasting properties of Wheat- Chickpea flour and Acha starch**

E R	WF	AS	CF	PV (RVU)	T (RVU)	BDV (RVU)	FV (RVU)	SB (RVU)	P Γ (Min)	PT(°C)
1	60.0	30.0	10.0	2035.50±47	1211.00±4.	824.50±43.	2606.50±58.	1395.50±54	5.70±0.	86.00±0.
	0	0	0	.37 <sup>e</sup>	24 <sup>e</sup>	13 <sup>k</sup>	68 <sup>e</sup>	.44 <sup>f</sup>	14 <sup>d</sup>	49 <sup>d</sup>
2	45.0	42.5	12.5	1722.50±38	931.00±65.	791.50±26.	1949.50±26.	1018.50±10	6.00±0.	88.83±0.
	0	0	0	.89 <sup>a</sup>	05 <sup>a</sup>	16 <sup>h</sup>	16 <sup>a</sup>	.61 <sup>a</sup>	18 <sup>g</sup>	03 <sup>h</sup>
3	52.5	42.5	5.00	2385.50±58	1614.00±0.	771.50±r58.	3149.00±97.	1535.00±97	5.87±0.	86.83±1.
	0	0		.68 <sup>j</sup>	00 <sup>j</sup>	68 <sup>g</sup>	58 <sup>j</sup>	.58 <sup>j</sup>	09 <sup>e</sup>	73 <sup>f</sup>
4	55.0	30.0	15.0	1920.00±42	1155.00±19	765.00±22.	2520.00±38.	1365.00±18	5.64±0.	85.63±0.
	0	0	0	.43 <sup>d</sup>	.79 <sup>d</sup>	62 <sup>f</sup>	18 <sup>d</sup>	.38 <sup>e</sup>	05 <sup>c</sup>	03 <sup>c</sup>

5	45.0	35.0	15.0	2131.50±89	1388.50±38	743.00±50.	2879.00±82.	1490.50±43	5.57±0.	85.60±0.
	0	0	0	.80 <sup>h</sup>	.89 <sup>h</sup>	91 <sup>d</sup>	02 <sup>h</sup>	.13 <sup>h</sup>	05 <sup>b</sup>	07 <sup>c</sup>
6	52.0	36.0	12.0	1898.50±94	1089.50±94	809.00±0.0	2185.00±103	1095.50±9.	5.44±0.	76.28±0.
	0	0	0	.05 <sup>c</sup>	.05 <sup>b</sup>	0 <sup>i</sup>	.24 <sup>b</sup>	19 <sup>b</sup>	05 <sup>a</sup>	53 <sup>a</sup>
7	48.5	35.5	16.0	2058.50±6.	1359.50±20	699.00±26.	2702.50±53.	1343.00±73	5.74±0.	86.35±0.
	0	0	0	36 <sup>g</sup>	.51 <sup>g</sup>	87 <sup>b</sup>	03 <sup>f</sup>	.54 <sup>d</sup>	09 <sup>d</sup>	07 <sup>e</sup>
8	50.0	30.0	20.0	1774.50±47	1115.50±43	659.00±4.2	2458.00±43.	1342.50±0.	5.64±0.	88.10±0.
	0	0	0	.37 <sup>b</sup>	.13 <sup>c</sup>	4 <sup>a</sup>	84 <sup>c</sup>	71 <sup>c</sup>	05 <sup>c</sup>	00 <sup>g</sup>
9	45.0	35.0	20.0	2006.50±33	1292.00±9.	714.50±23.	2715.00±19.	1423.00±9.	5.57±0.	85.98±0.
	0	0	0	.23 <sup>e</sup>	89 <sup>f</sup>	33 <sup>c</sup>	79 <sup>g</sup>	89 <sup>g</sup>	05 <sup>b</sup>	60 <sup>d</sup>
10	60.0	35.0	5.00	2294.00±41	1481.50±23	812.50±17.	2990.50±40.	1509.00±16	5.93±0.	86.38±0.
	0	0	0	.01 <sup>i</sup>	.33 <sup>i</sup>	68 <sup>j</sup>	31 <sup>i</sup>	.97 <sup>i</sup>	00 <sup>f</sup>	04 <sup>e</sup>
11	52.5	42.5	5.00	2385.50±58	1614.00±0.	771.50±58.	3149.00±97.	1535.00±97	5.87±0.	86.83±1.
	1	0	0	.68 <sup>j</sup>	00 <sup>j</sup>	68 <sup>g</sup>	58 <sup>j</sup>	.58 <sup>j</sup>	09 <sup>e</sup>	73 <sup>f</sup>
12	45.0	5.00	5.00	2483.50±16	1723.00±39	760.50±23.	3267.00±36.	1544.00±76	5.90±0.	84.80±1.
	2	0	0	.26 <sup>k</sup>	.59 <sup>k</sup>	33 <sup>e</sup>	77 <sup>k</sup>	.37 <sup>k</sup>	04 <sup>f</sup>	06 <sup>b</sup>
13	60.0	35.0	5.00	2294.00±41	1481.50±23	812.50±17.	2990.50±40.	1509.00±16	5.93±0.	86.38±0.
	3	0	0	.01 <sup>i</sup>	.33 <sup>i</sup>	68 <sup>j</sup>	31 <sup>i</sup>	.97 <sup>i</sup>	00 <sup>f</sup>	04 <sup>e</sup>
14	45.0	5.00	5.00	2483.50±16	1723.00±39	760.50±23.	3267.00±36.	1544.00±76	5.90±0.	84.80±1.
	4	0	0	.26 <sup>k</sup>	.59 <sup>k</sup>	33 <sup>e</sup>	77 <sup>k</sup>	.37 <sup>k</sup>	04 <sup>f</sup>	06 <sup>b</sup>

Mean values with different superscripts within the same column are significantly different (p>0.05); CPF: Chickpea flour

**N.B:** ER- Experimental runs, AS- Acha starch, CF- chickpea flour, PV- peak viscosity, T-Trough, BDV- Breakdown viscosity, F-Final viscosity, SB- Setback viscosity, PT- Pasting time, PT- Pasting Temperature

**Table 4: Functional Properties of Wheat-Acha starch -Chickpea Composite Flour**

Experimen tal Runs	WF	AS	CP F	WAC (g/ml)	BD(g/ml)	S Pr (%)	OAC (g/ml)	SI(g/ml)
1	60.0	30.0	10.0	183.00±0.0	0.7031±0.0	4.73±0.0	216.00±0.0	4.00±0.0
	0	0	0	2 <sup>b</sup>	1 <sup>k</sup>	8 <sup>c</sup>	7 <sup>bc</sup>	0 <sup>b</sup>
2	45.0	42.5	12.5	258.00±0.0	0.5588±0.0	5.17±0.1	186.00±0.0	7.50±0.7
	0	0	0	3 <sup>d</sup>	0 <sup>a</sup>	5 <sup>g</sup>	2 <sup>a</sup>	1 <sup>c</sup>
3	52.5	42.5	5.00	182.00±0.0	0.5775±0.3	5.00±0.0	232.00±0.0	4.00±0.0
	0	0		1 <sup>b</sup>	5 <sup>c</sup>	1 <sup>ef</sup>	3 <sup>e</sup>	0 <sup>b</sup>
4	55.0	30.0	15.0	192.00±0.0	0.6667±0.0	4.52±0.0	213.00±0.0	3.50±0.7
	0	0	0	2 <sup>c</sup>	0 <sup>d</sup>	2 <sup>a</sup>	2 <sup>b</sup>	1 <sup>a</sup>
5	45.0	35.0	15.0	190.00±0.0	0.6792±0.0	4.77±0.0	226.00±0.0	4.00±0.0
	0	0	0	2 <sup>c</sup>	3 <sup>i</sup>	0 <sup>c</sup>	8 <sup>d</sup>	0 <sup>b</sup>
6	52.0	36.0	12.0	181.00±0.0	0.6712±0.0	4.97±0.0	219.00±0.0	4.00±0.0
	0	0	0	1 <sup>ab</sup>	2 <sup>f</sup>	6 <sup>e</sup>	6 <sup>c</sup>	0 <sup>b</sup>
7	48.5	35.5	16.0	180.00±0.0	0.6733±0.0	4.64±0.0	221.00±0.0	3.50±0.7
	0	0	0	1 <sup>a</sup>	0 <sup>g</sup>	3 <sup>b</sup>	5 <sup>c</sup>	1 <sup>a</sup>
8	50.0	30.0	20.0	189.00±0.0	0.6800±0.0	4.93±0.0	212.00±0.0	3.50±0.7
	0	0	0	1 <sup>bc</sup>	0 <sup>j</sup>	5 <sup>e</sup>	3 <sup>b</sup>	1 <sup>a</sup>
9	45.0	35.0	20.0	193.00±0.0	0.6684±0.0	4.83±0.0	220.00±0.0	4.00±0.0
	0	0	0	1 <sup>c</sup>	5 <sup>e</sup>	3 <sup>d</sup>	2 <sup>c</sup>	0 <sup>b</sup>
10	60.0	35.0	5.00	175.00±0.0	0.6756±0.0	6.08±0.2	209.00±0.0	4.00±0.0
	0	0		1 <sup>a</sup>	0 <sup>h</sup>	2 <sup>h</sup>	2 <sup>b</sup>	0 <sup>b</sup>
11	52.5	42.5	5.00	182.00±0.0	0.5775±0.3	5.00±0.0	232.00±0.0	4.00±0.0
	0	0		1 <sup>b</sup>	5 <sup>c</sup>	1 <sup>ef</sup>	3 <sup>e</sup>	0 <sup>b</sup>
12	45.0	50.0	5.00	183.00±0.0	0.5608±0.1	5.17±0.0	229.00±0.0	4.00±0.0
	0	0		2 <sup>b</sup>	<sup>b</sup>	1 <sup>g</sup>	3 <sup>d</sup>	0 <sup>b</sup>
13	60.0	35.0	5.00	175.00±0.0	0.6756±0.0	6.08±0.2	209.00±0.0	4.00±0.0
	0	0		1 <sup>a</sup>	0 <sup>h</sup>	2 <sup>h</sup>	2 <sup>b</sup>	0 <sup>b</sup>
14	45.0	50.0	5.00	183.00±0.0	0.5608±0.1	5.17±0.0	229.00±0.0	4.00±0.0
	0	0		2 <sup>b</sup>	<sup>b</sup>	1 <sup>g</sup>	3 <sup>d</sup>	0 <sup>b</sup>

Mean values with different superscripts within the same column are significantly different (p<0.05); CPF: Chickpea flour, WAC: Water absorption capacity, OAC: Oil absorption capacity

WF-Wheat Flour, AS- Acha Flour, CPF, Chickpea flour, WAC – Water absorption capacity, BD-Bulk Density, S P-Swelling power, OAC- Oil absorption capacity, SI- Solubility Index

UNDER PEER REVIEW